#### UNIVERSITY OF CAMBRIDGE INTERNATIONAL EXAMINATIONS

International General Certificate of Secondary Education

#### MARK SCHEME for the June 2005 question paper

#### 0620 CHEMISTRY

0620/03

Paper 3 (Extended Theory), maximum mark 80

This mark scheme is published as an aid to teachers and students, to indicate the requirements of the examination. It shows the basis on which Examiners were initially instructed to award marks. It does not indicate the details of the discussions that took place at an Examiners' meeting before marking began. Any substantial changes to the mark scheme that arose from these discussions will be recorded in the published *Report on the Examination*.

All Examiners are instructed that alternative correct answers and unexpected approaches in candidates' scripts must be given marks that fairly reflect the relevant knowledge and skills demonstrated.

Mark schemes must be read in conjunction with the question papers and the Report on the Examination.

• CIE will not enter into discussion or correspondence in connection with these mark schemes.

CIE is publishing the mark schemes for the June 2005 question papers for most IGCSE and GCE Advanced Level and Advanced Subsidiary Level syllabuses and some Ordinary Level syllabuses.



### Grade thresholds for Syllabus 0620 (Chemistry) in the June 2005 examination.

	maximum	minimum mark required for grade:			
	mark available	А	С	E	F
Component 3	80	58	30	16	11

The threshold (minimum mark) for B is set halfway between those for Grades A and C. The threshold (minimum mark) for D is set halfway between those for Grades C and E. The threshold (minimum mark) for G is set as many marks below the F threshold as the E threshold is above it.

Grade A\* does not exist at the level of an individual component.

June 2005


## **IGCSE**

# MARK SCHEME

**MAXIMUM MARK: 80** 

**SYLLABUS/COMPONENT: 0620/03** 

**CHEMISTRY Extended Theory** 



Page 1	Mark Scheme	Syllabus	Paper
	IGCSE – JUNE 2005	0620	3

1	(a)	(i) darker or actual colours chlorine yellow, yellow/green bromine orange, brown, brownish red iodine black grey, purple					
		(ii) gas, liquid, solid all three needed					
		(iii)	colourless <b>or</b> (pale) yellow gas	[1] [1]			
	(b)	Must have a correct reagent otherwise wc = 0					
		l chlorine water <b>or</b> bubble in chlorine gas ow <b>or</b> orange <b>or</b> brown k brown <b>or</b> grey crystals	[1] [1]				
			cept colour that is darker than for bromide)	[1]			
		off yell	add (acidified) silver nitrate(aq) white <b>or</b> pale yellow <b>or</b> cream <u>precipitate</u> <b>or</b> soluble in aqueous ammonia ow <u>precipitate</u> insoluble in aqueous ammonia cipitate essential then either colour <b>or</b> solubility in aqueous ammonia	[1] [1] [1]			
		<b>OR</b> add lead nitrate(aq) pale yellow <b>or</b> off white <b>or</b> cream <u>precipitate</u> yellow <u>precipitate</u> insoluble in aqueous ammonia					
			cept any test that could work – electrolysis, iron(III) salt mine, potassium dichromate, potassium manganate(VII) etc.				
	(c)		$3Cl_2 = 2ICl_3$ having either reactants <b>or</b> products correct ONLY [1]	[2]			
	(d)		orine ND lower M <sub>r</sub> or lower density or lighter molecules or molecules move faster	[1] [2]			
		OR	lighter <b>or</b> based on A <sub>r</sub> MAX [1] smaller with no additional comment <b>or</b> sieve idea [0] <b>N.B.</b> a total of [3] not [2]				
			TOTAL =	12			
2	(a)		+ I <sub>2</sub> = Zn <sup>2+</sup> + 2I <sup>-</sup> having either reactants <b>or</b> products correct ONLY [1]	[2]			
	(b)		zinc and sodium hydroxide white precipitate solves in excess (only if precipitate mentioned)	[1] [1]			
		Mai	zinc and ammonia same results rk either first (sodium hydroxide <b>or</b> aqueous ammonia), if completely correct, then litional [1] can be awarded for stating that the other has the same results.	[1] an			

Paper

**Syllabus** 

			IGCSE – JUNE 2005	0620	3
	(c)	(i)	zinc <u>and</u> a reason Do not mark conseq to iodine in excess		[1]
		(ii)	final mass of zinc bigger <b>or</b> the level section higher <b>or</b> legradient less steep <b>or</b> longer time <b>or</b> falls more slowly	ess zinc used	up [1]
		(iii)	steeper gradient same loss of mass of zinc		[1] [1]
					TOTAL = 10
3	(a)	(i)	$CH_3$ - $CH$ == $CH_2$		[1]
		(ii)	conseq to (i) correct repeat unit COND evidence of continuation		[1] [1]
		(iii)	monomer COND because it has a double bond <b>or</b> unsaturated <b>or NOT</b> addition	alkene	[1] [1]
	(b)	(i)	to remove fibres <b>or</b> remove solid <b>NOT</b> precipitate, <b>NOT</b> impurities, <b>NOT</b> to obtain a filtrate	Э	[1]
		(ii)	because silver atoms have <u>lost electrons</u> <b>OR</b> oxidation number increased		[1]
		(iii)	silver chloride		[1]
	(c)	(i)	name of an ester formula of an ester if they do not correspond MAX [1] <b>Accept</b> name - terylene for formula ester linkage and continuation If a 'fat' complete structure must be correct e.g. C <sub>17</sub> H <sub>35</sub> of Mark for formula only - [1]	etc.	[1] [1]
		(ii)	alcohol <b>or</b> alkanol <b>NOT</b> a named alcohol		[1]
	(d)	(i)	acid loses a proton base accepts a proton		[2] [1]
			OR same explanation but acid loses a hydrogen ion ( and base gains hydrogen ion (1)	(1)	
		(ii)	only partially ionised <b>or</b> poor hydrogen ion donor <b>or</b> poor <b>NOT</b> does not form many hydrogen ions in water <b>or</b> low ions <b>NOT</b> pH	•	

Mark Scheme

Page 2

Page 3	Mark Scheme	Syllabus	Paper
	IGCSE – JUNE 2005	0620	3

4	(a)	(i)	correct word equation (carbon dioxide and water)  Accept correct symbol equation	[1]
		(ii)	Must have a correct reagent otherwise wc = 0 add (acidified) barium chloride(aq) <b>or</b> nitrate <b>or</b> add barium ions <b>COND</b> white precipitate <b>NOT</b> lead(II) compounds	[1] [1]
		(iii)	low pH <b>or</b> universal indicator turns red(aq) pH 3 <b>or</b> less	[1]
	(b)	(i)	$H_2S + 2O_2 = H_2SO_4$ unbalanced [1]	[2]
		(ii)	unpleasant smell <b>or</b> it is poisonous <b>or</b> when burnt forms acid rain <b>or</b> forms suldioxide <b>or</b> forms sulphuric acid <b>NOT</b> it is a pollutant	lphur [1]
		(iii)	2H to 1S  COND 8e around sulphur atom 2e per hydrogen atom  THREE correct  TWO from above [1] lonic structure = [0]	[2]
	(c)	(i)	vanadium oxide ${f or}$ vanadium(V) oxide ${f or}$ vanadium pentoxide or $V_2O_5$ Must be correct oxidation state if one given	[1]
		(ii)	400 to 500° C	[1]
		(iii)	add to (concentrated) sulphuric acid <b>NOT</b> dilute <b>COND</b> (upon sulphuric acid) above then add water	[1] [1]
	(d)	mol mol	ass of one mole of $CaSO_4 = 136$ les of $CaSO_4$ in 79.1g = 0.58 accept 0.6 les of $H_2O$ in 20.9 g = 1.16 accept 1.2 aseq x = 2 x given as an integer	[1] [1] [1]
			TOTAL	= 16
5	(a)	(i)	A is glutamic acid B is alanine	[1] [1]
		(ii)	because acids are colourless <b>or</b> to make them visible <b>or</b> to show positions of the samples <b>or</b> distance travelled	[1]
		(iii)	compare with known acids <b>or</b> reference samples <b>or</b> standards <b>Accept</b> from colours of samples	[1]
		(iv)	amide linkage  COND different monomers continuation  Accept hydrocarbon part of chain as boxes  If nylon 6 then only one monomer [1] NOT different monomers	[1] [1] [1]

**Syllabus** 

Paper

			IGCSE – JUNE 2005		0620	3
	(b)	corr	ect structure as syllabus (box representation ect linkageO tinuation	)		[1] [1]
	(c)	(i)	$C_6H_{12}O_6 = 2C_2H_5OH + 2CO_2$ not balanced [1] <b>Accept</b> $C_2H_6O$			[2]
		(ii)	gives out <u>energy</u> <b>or</b> equivalent <b>NOT</b> heat N.B. a total of [1] not [2]			[1]
		(iii)	glucose used up <b>or</b> yeast 'killed' by ethanol <b>NOT</b> yeast used up <b>NOT</b> r	eactant use	ed up	[1]
		(iv)	oxidise alcohol to acid <b>or</b> to ethanoic acid <b>or</b> to carbon dioxide and water <b>or</b> if oxygen present aerobic respiration <b>or</b> cannot have anaerobic respiration in presence it is anaerobic respiration, must be add	•	•	[1]
		(v)	fractional distillation			[1]
						TOTAL = 15
6	(a)	(i)	bauxite			[1]
		(ii)	to reduce melting point <b>or</b> improve conductivor as a solvent <b>or</b> reduce the working temper			[1]
		(iii)	carbon dioxide or monoxide or fluorine			[1]
	(b)	(i)	aluminium			[1]
		(ii)	solution goes colourless <b>or</b> copper formed <b>or</b> a <u>brown solid</u> forms <b>or</b> blue colour disapp <b>or</b> bubbles <b>NOT</b> goes clear <b>or</b> copper formed	pears		[1]
		(iii)	covered with an oxide layer			[1]
	(c)	reac	ction no reaction reaction	า		[1] [1]
	(d)	(i)	$2Al(OH)_3 = Al_2O_3 + 3H_2O$ Not balanced [1]			[2]
		(ii)	Aluminium nitrate = aluminium oxide + nitronly TWO correct products [1]	ogen dioxid	e + oxygen	[2]
						TOTAL = 12

Mark Scheme

Page 4

6