CAMBRIDGE INTERNATIONAL EXAMINATIONS

International General Certificate of Secondary Education

MARK SCHEME for the May/June 2014 series

0620 CHEMISTRY

0620/31

Paper 3 (Extended Theory), maximum raw mark 80

This mark scheme is published as an aid to teachers and candidates, to indicate the requirements of the examination. It shows the basis on which Examiners were instructed to award marks. It does not indicate the details of the discussions that took place at an Examiners' meeting before marking began, which would have considered the acceptability of alternative answers.

Mark schemes should be read in conjunction with the question paper and the Principal Examiner Report for Teachers.

Cambridge will not enter into discussions about these mark schemes.

Cambridge is publishing the mark schemes for the May/June 2014 series for most IGCSE, GCE Advanced Level and Advanced Subsidiary Level components and some Ordinary Level components.



Page 2	Mark Scheme	Syllabus	Paper
	IGCSE – May/June 2014	0620	31

1 (a) $\underline{A}, \underline{D}, \underline{E}$ (1)

same number of protons and electrons/electrically neutral (1) [2]

(b) C (1)

more electrons than protons/36e⁻ and 34p⁺/it has gained electrons (1) [2]

(c) B, F (1)

(d) they have same number of protons (1)

different number of neutrons/neutron number (1) [2]

[Total: 7]

2 (a) (i) filtration (1)

chlorination (1) [2]

(ii) Any two from: [2]

- manufacture of ethanol
- used in the manufacture of sulfuric acid **or** in the Contact process
- manufacture of hydrogen or ammonia or for the Haber process

(iii) Any **two** from: [2]

- cooking
- washing or laundry
- drinking
- toilets
- watering plants
- (domestic) heating
- **(b)** boiling or turning to steam (1)

then condensing/condensation (1) [2]

[Total: 7]

(a) (i) (particles) spread to fill total available volume/move from high concentration to low concentration/moves down a concentration gradient (1) [1]

[1]

(ii) mass or M_r (1)

(b) (i) helium atoms/molecules are lighter than molecules in air or N_2 and O_2 or helium is less dense than air or N_2 and O_2 .

or helium diffuses (through the porous barrier) faster than air or N_2 and O_2 . (1)

[1]

Page 3	Mark Scheme	Syllabus	Paper
	IGCSE – May/June 2014	0620	31

(ii) faster rate of diffusion/molecules move faster (at high temperatures). (1) [1]

(c) (i)
$$CH_4 + 2O_2 \rightarrow CO_2 + 2H_2O$$
 (1) [1]

(ii) would get a mixture of helium and carbon dioxide
or would get a mixture of gases

or waste of methane/natural gas/fossil fuel (1)

(iii) <u>fractional</u> distillation (1) [1]

[Total: 7]

[1]

4 (a) (i)

Group number	I	II	III	IV	V	VI	VII
symbol	Na	Mg	Al	Si	Р	S	Cl
number of valency electrons	1	2	3	4	5	6	7
valency	1	2	3	4	3	2	1

(1) for each line [2]

- (ii) number of valency electrons = the group number (1) [1]
- (iii) for Na to Al

the valency is the same as the number of valency (outer) electrons (1)

(because) this is the number of electrons **lost** (for full energy level) (1)

for P to C1

the valency is 8 – [number of valency (outer) electrons] **or** valency + valency electrons = 8 (1)

(because) this is number of electrons **needed** (or to be **gained**) (for full energy level) (1)

(b) (i) Assume change is from L to R unless clearly stated: basic to amphoteric to acidic (2)

[2]

(ii) ionic (metal) chlorides on the left (1) covalent (non-metal) chlorides on the right (1)

[2]

[Total: 11]

Page 4	Mark Scheme	Syllabus	Paper
	IGCSE – May/June 2014	0620	31

5 (a) M1: (zinc sulfide) heated/roasted/burnt in air (1)

M2: zinc oxide formed (1)

M3: zinc oxide **reduced** (1)

M4: (by adding) coke or carbon (1)

M5: Balanced equation (any one of) (1) [5]

- (b) Any two from:
 - (making) brass or alloys (1)
 - galvanising (1)
 - sacrificial protection (1)
 - batteries (1)

[Total: 7]

[2]

- 6 (a) (i) rate at t_2 less than at t_1 or the rate decreases (1)
 - rate at t₃ zero/reaction stopped (1)

[2]

[2]

[2]

- (ii) rate at t_2 less than at t_1 because **concentration** of hydrogen peroxide is less at t_2 **or concentration** of hydrogen peroxide is decreasing. (1)
- (rate at t₃ zero/reaction stopped because) hydrogen peroxide is used up (1) [2]
- (b) (i) steeper and must come from the origin (1) final volumes the same (1)
 - (ii) Any **two** from:
 steeper curve because of a faster rate
 faster rate because of increased surface area
 same amount/volume/mass/no of mol of hydrogen peroxide
 ecf for M1 for a shallower curve because of slower rate.

Page 5	Mark Scheme	Syllabus	Paper	
	IGCSE – May/June 2014	0620	31	

(c) filter (and rinse/wash) (1)

dry manganese (IV) oxide (1)

weigh/measure mass manganese(IV) oxide after reaction (1)

the mass should be 0.1 g or unchanged. (1)

[4]

[3]

(d) number of moles of O_2 formed = 0.096/24 = 0.004 (1) number of moles of H_2O_2 in 40 cm³ of solution = 0.004 × 2 = 0.008 (1)

concentration of the hydrogen peroxide in $mol/dm^3 = 0.008/0.04 = 0.2$ (1)

[Total:15]

7 (a) (i)

aqueous solution	lead Pb	magnesium Mg	zinc Zn	silver Ag
lead (II) nitrate				
magnesium nitrate	Χ×		*	×
zinc nitrate	×	✓		×
silver(I) nitrate	✓	✓	✓	

each horizontal line correct (1)

[3]

(ii) Zn (1)

An arrow from $Zn \text{ to } Zn^{2+}$ (1)

[2]

(iii)
$$Zn + 2Ag^+ \rightarrow Zn^{2+} + 2Ag$$
 (1)

[1]

(b) (i) correct direction from zinc to lead (1)

[1]

(ii) metals react by losing electrons (1)

the more reactive metal/zinc will lose electrons more readily (making the electrode negatively charged). (1)

[2]

[Total: 14]

	Page 6		Mark Scheme	Syllabus	Paper		
			IGCSE – May/June 2014	0620	31		
	(iii)	ii) manganese and zinc are more reactive than lead (and/or copper) (1)					
		lead	[2]				
	(iv)		the polarity of a Mn/Zn (cell)				
		or th	ne voltages of Zn/Pb and Mn/Pb (cells) (1)		[1]		
					[Total: 12]		
8	(a) (i)	CH ₃	-CH=CH-CH₃ (1)		[1]		
	(ii)	one					
		corr	ect sequencing of a second amide link and monome	ers (1)			
			correct amide links and rest of structure correct nomers if seen) and correct continuation bonds (1)	(including addition	onal [3]		
		-	-c	3 marks			
	(iii)	prot	ein or polypeptide or named protein (1)		[1]		
	(iv)	addi	ition: only the polymer or one product is formed (1)				
		cond	densation: the polymer and a small molecule/water	/HC l is formed (1) [2]		
	(b) (i)	does	s not break down or rot or decompose (1)				
		by n	nicrobes or fungi or bacteria or by living organisms	(1)	[2]		
	(ii)	•	three from: al pollution (1)		[3]		
		(sho	ortage of) landfill sites (1)				
		dan	ger to wildlife/animals (including at sea) (1)				
		toxic	c gases when burnt or greenhouse gases produced	when burned (1)			
		sistant	to corrosion/unreactive to water/more durable (1)		[2]		
	ligl	lighter/less dense (1)					

easier to manufacture/can be moulded (1)

good insulator/keeps the water cold (1)