

5 2

UNIVERSITY OF CAMBRIDGE INTERNATIONAL EXAMINATIONS International General Certificate of Secondary Education

CANDIDATE NAME		
CENTRE NUMBER		CANDIDATE NUMBER
CHEMISTRY		0620/31
Paper 3 (Exten	ded)	October/November 2009
		1 hour 15 minutes
Candidates and	swer on the Question Paper.	

No Additional Materials are required.

READ THESE INSTRUCTIONS FIRST

Write your Centre number, candidate number and name on all the work you hand in. Write in dark blue or black pen. You may use a pencil for any diagrams, graphs or rough working. Do not use staples, paper clips, highlighters, glue or correction fluid.

DO NOT WRITE IN ANY BARCODES.

Answer all questions.

A copy of the Periodic Table is printed on page 16.

At the end of the examination, fasten all your work securely together.	For Exam	iner's Use
The number of marks is given in brackets [] at the end of each question or part questions.	1	
	2	
	3	
	4	
	5	
	6	
	7	
	Total	

This document consists of 14 printed pages and 2 blank pages.



1 (a) The	e major gases in unpolluted air are 79% nitrogen and 20% oxygen.
(i)	Name another gaseous element in unpolluted air.
	[1]
(ii)	Name two compounds in unpolluted air.
	[2]
(b) Tw	o common pollutants in air are carbon monoxide and the oxides of nitrogen.
(i)	Name another pollutant in air.
	[1]
(ii)	Describe how carbon monoxide is formed.
	[2]
(iii)	How are the oxides of nitrogen formed?
	[2]
(iv)	Explain how a catalytic converter reduces the emission of these two gases.
	[2]
	[Total: 10]



3	(a)	An	important ore of zinc is zinc blende, ZnS.	For
		(i)	How is zinc blende changed into zinc oxide?	Use
			[1]	
		(ii)	Write a balanced equation for the reduction of zinc oxide to zinc by carbon.	
			[2]	
	(b)	A m Thi	najor use of zinc is galvanizing; steel objects are coated with a thin layer of zinc. s protects the steel from rusting even when the layer of zinc is broken.	
			thin layer steel exposed to	
			of zinc oxygen and water	
			steel	
			Explain, by mentioning ions and electrons, why the exposed steel does not rust.	
			[3]	

For

Use

(c) Zinc electrodes have been used in cells for many years, one of the first was the Daniel cell in 1831. Examiner's voltmeter. copper electrode zinc electrode zinc sulfate(aq) copper(II) sulfate(aq) porous pot - stops solutions from mixing (i) Give an explanation for the following in terms of atoms and ions. observation at zinc electrode - the electrode becomes smaller explanation [1] observation at copper electrode – the electrode becomes bigger explanation [1] (ii) When a current flows, charged particles move around the circuit. What type of particle moves through the electrolytes? [1] Which particle moves through the wires and the voltmeter? [1] [Total: 10]

- 4 The distinctive smell of the seaside was thought to be caused by ozone, O₃. Ozone is a form of the element oxygen.
 - (a) A mixture of oxygen and ozone is formed by passing electric sparks through oxygen.

 $3O_2 \rightleftharpoons 2O_3$

Suggest a technique that might separate this mixture. Explain why this method separates the two forms of oxygen.

technique explanation______

(b) Ozone is an oxidant. It can oxidise an iodide to iodine.

$$2I^{\scriptscriptstyle -} \ \textbf{+} \ \textbf{O}_3 \ \textbf{+} \ 2\textbf{H}^{\scriptscriptstyle +} \ \rightarrow \ I_2 \ \textbf{+} \ \textbf{O}_2 \ \textbf{+} \ \textbf{H}_2\textbf{O}$$

(i) What would you see when ozone is bubbled through aqueous acidified potassium iodide?

(ii) Explain in terms of electron transfer why the change from iodide ions to iodine molecules is oxidation.
 (iii) Explain, using your answer to b(ii), why ozone is the oxidant in this reaction.
 [1]

(c)	lt is (C⊦	s now known that the smell of the seaside is due to the chemical dimethyl sulfide, $J_3)_2S.$	For Examiner's Use
	(i)	Draw a diagram that shows the arrangement of the valency electrons in one molecule of this covalent compound. Use x to represent an electron from a carbon atom. Use o to represent an electron from a hydrogen atom. Use • to represent an electron from a sulfur atom.	
	(ii)	[3] Name the three compounds formed when dimethyl sulfide is burnt in excess oxygen.	
		[2] [Total: 11]	

For

Examiner's Use

- 5 The first three elements in Group IV are carbon, silicon and germanium. The elements and their compounds have similar properties.
 - (a) The compound, silicon carbide, has a macromolecular structure similar to that of diamond.
 - (i) A major use of silicon carbide is to reinforce aluminium alloys which are used in the construction of spacecraft. Suggest **three** of its physical properties.

(ii) Complete the following description of the structure of silicon carbide.
 Each carbon atom is bonded to four ______ atoms.
 Each silicon atom is bonded to ______ carbon atoms. [2]

(b) Germanium(IV) oxide, GeO₂, has the same macromolecular structure as silicon(IV) oxide. Draw the structural formula of germanium(IV) oxide.

[3]

For Examiner's Use

(c) Germanium forms a series of hydrides comparable to the alkane	€S.
---	-----

(i) Draw the structural formula of the hydride which contains four germanium atoms per molecule.

(ii)	Predict the products of the complete combustion of this hydride.	[1]
		[2]

[Total: 11]

(a) Su	Ifuric acid is made by the Contact process.	For
	$2SO_2 + O_2 \rightleftharpoons 2SO_3$	Use
Thi	s is carried out in the presence of a catalyst at 450 $^\circ$ C and 2 atmospheres pressure.	
(i)	How is the sulfur dioxide made?	
	[1]	
(ii)	Give another use of sulfur dioxide.	
	[1]	
(iii)	Name the catalyst used.	
	[1]	
(iv)	If the temperature is decreased to 300 °C, the yield of sulfur trioxide increases. Explain why this lower temperature is not used.	
	[1]	
(v)	Sulfur trioxide is dissolved in concentrated sulfuric acid. This is added to water to make more sulfuric acid. Why is sulfur trioxide not added directly to water?	
	[1]	

6

For Examiner's Use

(b)	Sulfuric acid was first made in the Middle East by heating the mineral, green vitriol,
	$FeSO_4.7H_2O$. The gases formed were cooled.

	FeS gree	$O_4.7H_2O(s) \rightarrow FeSO_4(s)$ en crystals yellow powder	+ 7H ₂ O(g)	
	2Fe	$\text{SO}_4(s) \rightarrow \text{Fe}_2\text{O}_3(s) + \text{SO}_2(s)$	(g) + SO ₃ (g)	
	On	cooling		
	SO3 SO2	$H_3 + H_2O \rightarrow H_2SO_4$ sulfuric as $H_2 + H_2O \rightarrow H_2SO_3$ sulfurous	cid acid	
	(i)	How could you show that the fir	st reaction is reversible?	
				[2]
			-	
	(ii)	Sulfurous acid is a reductant. W manganate(VII) is added to a se	/hat would you see when acidified potassium olution containing this acid?	
				[2]
	/:::)	Suggest an explanation why su	Ilfurous acid in contact with air changes into sulfu	ric
	()	acid.		
(c)	9.12 forn	2g of anhydrous iron(II) sulfate v ned and the volume of sulfur trio:	vas heated. Calculate the mass of iron(III) oxide xide, at r.t.p., formed.	[1]
	2Fe	$sSO_4(s) \rightarrow Fe_2O_3(s) + SO_2(g)$	+ SO ₃ (g)	
	ma	ass of one mole of $FeSO_4 = 152$	2g	
	nu	mber of moles of FeSO ₄ used	=	
	nu	mber of moles of Fe_2O_3		
	for	med	=	
	ma	ass of one mole of Fe_2O_3	= g	
	ma	ass of iron(III) oxide formed	= g	
	nu	mber of moles of SO_3 formed	=	
	vo	lume of sulfur trioxide formed	= dm ³	
				[6]

[Total: 16]

7	Butan-1 Butan-1	-ol is used as a solvent for paints and varnishes, to make esters and as a fuel. -ol can be manufactured from but-1-ene, which is made from petroleum.	
	Biobuta biomas	nol is a fuel of the future. It can be made by the fermentation of almost any form o s - grain, straw, leaves etc.	of
	(a) But	-1-ene can be obtained from alkanes such as decane, $C_{10}H_{22}$, by cracking.	
	(i)	Give the reaction conditions.	
			[2]
	(ii)	Complete an equation for the cracking of decane, $C_{10}H_{22}$, to give but-1-ene.	
		$C_{10}H_{22} \rightarrow$	[2]
	(iii)	Name the reagent that reacts with but-1-ene to form butan-1-ol.	

- [1]
- (b) (i) Balance the equation for the complete combustion of butan-1-ol.

	$\label{eq:c4H9OH} \begin{array}{cccccccccccccccccccccccccccccccccccc$	[2]
(ii)	Write a word equation for the preparation of the ester butyl methanoate.	
		[2]

(c)	The biol	The fermentation of biomass by bacteria produces a mixture of products which include biobutanol, propanol, hydrogen and propanoic acid.									
	(i)	Draw the structural formula of propanol and of propanoic acid. Show all the bonds.									
		propanol									
		propanoic acid									
		[2]									
	(ii)	Why is it important to develop these fuels, such as biobutanol, as alternatives to petroleum?									
		[1]									
(d)	Ho\ che	w could you show that butanol made from petroleum and biobutanol are the same mical?									
		[1]									
		[Total: 13]									

BLANK PAGE

BLANK PAGE

Permission to reproduce items where third-party owned material protected by copyright is included has been sought and cleared where possible. Every reasonable effort has been made by the publisher (UCLES) to trace copyright holders, but if any items requiring clearance have unwittingly been included, the publisher will be pleased to make amends at the earliest possible opportunity.

University of Cambridge International Examinations is part of the Cambridge Assessment Group. Cambridge Assessment is the brand name of University of Cambridge Local Examinations Syndicate (UCLES), which is itself a department of the University of Cambridge.

	0	4 Helium	9 20 ine Neon 10	.5 40 . 7 Ar	0 84 Kr	ine Krypton 36	. 131	ne Xenon 54	t Rn	tine Radon 86		b 175 b Lu vium 71	o Lawrencium
	>		و الم	35. Chlor 17	∞ ¤	35 Brom	12	53 lodi	A	Astat 85		71 70 70	Ž
	N		16 Oxygen 8	32 Sultur 16	62 CO	Selenium 34	128	Tellurium 52	Ро	Polonium 84		169 Tan Thulium	Mendeleviur
	>		14 Nitrogen 7	31 Phosphorus 15	75 A c	Arsenic 33	122	SD Antimony 51	209 Bi	Bismuth 83		167 Er Erbium 68	F Fermium
	2		12 Carbon 6	28 Silicon	73	Germanium 32	119	20 20	207 Pb	Lead 82		165 Ho Holmium 67	Einsteinium
	■		5 Boron 1	27 A1 Auminium 13	20 20	Gallium 31	115	Indium 49	204 T 1	Thallium 81		162 Dysprosium 66	Cf Californium
					65 7 5	Zinc 30	112	Cadmium 48	201 Hg	Mercury 80		159 Tb Terbium 65	BK Berkelium
					64	Copper 29	108	Ag Silver 47	197 Au	Gold 79		157 Gd Gadolinium 64	Curium Curium
dno					59 Ni	Nickel 28	106	Palladium 46	195 Pt	Platinum 78		152 Eu Europium 63	Am
Gre					20 20	Cobalt 27	103	Rhodium 45	192 Ir	Iridium 77		150 Sam arium 62	Plutonium
		Hydrogen			56 F.A	lron 26	101	Ruthenium 44	190 OS	Osmium 76		Promethium 61	Neptunium
					55 Min	Manganese 25	ŀ	Technetium 43	186 Re	Rhenium 75		144 Neodymium 60	238 Uranium
					52 C	Chromium 24	96	Molybdenum 42	184 V	Tungsten 74		141 Pr Fraseodymium 59	Pa Protactinium
					51	Vanadium 23	93	Niobium 41	181 Ta	Tantalum 73		140 Cer Cerium 58	232 Thorium
					48	Titanium 22	9 19	Zirconium 40	178 Hf	Hafnium 72			nic mass bol number
				I	45 CC	Scandium 21	88	Yttrium 39	139 La	Lanthanum 57 *	227 Actinium 89	l series series	 = relative ator = atomic sym = nrotom (atom
	=		9 Beryllium	24 Mg Magnesium 12	40 2	Calcium 20	88	Strontium 38	137 Ba	Barium 56	226 Raa 88	anthanoic Actinoid s	X 70
	_		3 Lithium	23 Na Sodium	39	Potassium 19	85	Rubidium 37	133 Cs	Caesium 55	Fr Francium 87	58-71 Lá 90-103 /	ey