

First Variant Question Paper



UNIVERSITY OF CAMBRIDGE INTERNATIONAL EXAMINATIONS
International General Certificate of Secondary Education

CANDIDATE
NAME

CENTRE
NUMBER

--	--	--	--	--

CANDIDATE
NUMBER

--	--	--	--

CHEMISTRY

0620/31

Paper 3 (Extended)

October/November 2008

1 hour 15 minutes

Candidates answer on the Question Paper.

No Additional Materials are required.

READ THESE INSTRUCTIONS FIRST

Write your Centre number, candidate number and name on **all** the work you hand in.

Write in dark blue or black pen.

You may use a pencil for any diagrams, graphs or rough working.

Do not use staples, paper clips, highlighters, glue or correction fluid.

DO NOT WRITE IN ANY BARCODES

Answer **all** questions.

A copy of the Periodic Table is printed on page 12.

At the end of the examination, fasten **all** your work securely together.

The number of marks is given in brackets [] at the end of each question or part questions.

For Examiner's Use	
1	
2	
3	
4	
5	
6	
7	
Total	

This document consists of **12** printed pages.



1 Complete the following table.

gas	test for gas
ammonia	
	bleaches damp litmus paper
hydrogen	
	relights a glowing splint
	turns limewater milky

For
Examiner's
Use

[Total: 5]

2 There are three types of giant structure – ionic, metallic and macromolecular.

- (a) Sodium nitride is an ionic compound. Draw a diagram that shows the formula of the compound, the charges on the ions and the arrangement of the valency electrons around the negative ion.

Use x to represent an electron from a sodium atom.
Use o to represent an electron from a nitrogen atom.

For
Examiner's
Use

[3]

- (b) (i) Describe metallic bonding.

.....
..... [3]

- (ii) Use the above ideas to explain why
metals are good conductors of electricity,

..... [1]

metals are malleable.

..... [2]

- (c) Silicon(IV) oxide has a macromolecular structure.

- (i) **Describe** the structure of silicon(IV) oxide (a diagram is not acceptable).

.....
.....
..... [3]

- (ii) Diamond has a similar structure and consequently similar properties.
Give **two** physical properties common to both diamond and silicon(IV) oxide.

.....
..... [2]

[Total: 14]

3 Steel is an alloy made from impure iron.

(a) Both iron and steel rust. The formula for rust is $\text{Fe}_2\text{O}_3 \cdot 2\text{H}_2\text{O}$.
It is hydrated iron(III) oxide.

(i) Name the **two** substances that must be present for rusting to occur.

..... [2]

(ii) Painting and coating with grease are two methods of preventing iron or steel from rusting. Give **two** other methods.

.....
..... [2]

(b) (i) Name a reagent that can reduce iron(III) oxide to iron.

..... [1]

(ii) Write a symbol equation for the reduction of iron(III) oxide, Fe_2O_3 , to iron.

..... [2]

(c) (i) Calculate the mass of one mole of $\text{Fe}_2\text{O}_3 \cdot 2\text{H}_2\text{O}$.

..... [1]

(ii) Use your answer to (i) to calculate the percentage of iron in rust.

.....
..... [2]

(d) Iron from the blast furnace is impure. Two of the impurities are carbon and silicon. These are removed by blowing oxygen through the molten iron and adding calcium oxide.

(i) Explain how the addition of oxygen removes carbon.

.....
..... [1]


(ii) Explain how the addition of oxygen and calcium oxide removes silicon.

.....
..... [2]

[Total: 13]

- 4 Across the world, food safety agencies are investigating the presence of minute traces of the toxic hydrocarbon, benzene, in soft drinks. It is formed by the reduction of sodium benzoate by vitamin C.

For
Examiner's
Use



- (a) Sodium benzoate is a salt, it has the formula C_6H_5COONa . It can be made by the neutralisation of benzoic acid by sodium hydroxide.

(i) Deduce the formula of benzoic acid.

..... [1]

(ii) Write a word equation for the reaction between benzoic acid and sodium hydroxide.

..... [1]

(iii) Name **two** other compounds that would react with benzoic acid to form sodium benzoate.

..... [2]

- (b) Benzene contains 92.3% of carbon and its relative molecular mass is 78.

(i) What is the percentage of hydrogen in benzene?

..... [1]

(ii) Calculate the ratio of moles of C atoms: moles of H atoms in benzene.

.....
..... [2]


(iii) Calculate its empirical formula and **then** its molecular formula.

The empirical formula of benzene is

The molecular formula of benzene is [2]

6

(c) The structural formula of Vitamin C is drawn below.



For
Examiner's
Use

(i) What is its molecular formula?

..... [1]

(ii) Name the two functional groups which are circled.

..... [2]

[Total: 12]

5 The electrolysis of concentrated aqueous sodium chloride produces three commercially important chemicals hydrogen, chlorine and sodium hydroxide.

For
Examiner's
Use

(a) The ions present are $\text{Na}^+(\text{aq})$, $\text{H}^+(\text{aq})$, $\text{Cl}^-(\text{aq})$ and $\text{OH}^-(\text{aq})$.

(i) Complete the ionic equation for the reaction at the negative electrode (cathode).



(ii) Complete the ionic equation for the reaction at the positive electrode (anode).



(iii) Explain why the solution changes from sodium chloride to sodium hydroxide.

..... [1]

(b) (i) Why does the water supply industry use chlorine?

..... [1]

(ii) Name an important chemical that is made from hydrogen.

..... [1]


(iii) How is sodium hydroxide used to make soap?

..... [2]

[Total: 7]

6 The reactivity series lists metals in order of reactivity.

- (a) To find out which is the more reactive metal, zinc or tin, the following experiment could be carried out.



This experiment could be carried out with other metals and the results recorded in a table. Then the order of reactivity can be deduced.

- (i) The order was found to be:
 manganese most reactive
 zinc
 tin
 silver least reactive

Complete the table of results from which this order was determined.

aqueous solution	tin Sn	manganese Mn	silver Ag	zinc Zn
tin(II) nitrate		R	NR	R
manganese(II) nitrate				
silver(I) nitrate				
zinc nitrate				

[3]

- (ii) Write the ionic equation for the reaction between tin atoms and silver(I) ions.

.....

[2]

For
Examiner's
Use

- (iii) The following is a redox reaction.



Indicate on the equation the change which is oxidation.
Give a reason for your choice.

..... [2]

- (iv) Explain why experiments of this type cannot be used to find the position of aluminium in the reactivity series.

..... [2]

- (b) Potassium and calcium are very reactive metals at the top of the series. Because their ions have different charges, K^+ and Ca^{2+} , their compounds behave differently when heated.

- (i) Explain why the ions have different charges.

..... [2]

- (ii) Their hydroxides are heated.
If the compound decomposes, complete the word equation.
If it does not decompose, write "no reaction".

Potassium hydroxide \longrightarrow

Calcium hydroxide \longrightarrow [2]

- (iii) Complete the equations for the decomposition of their nitrates.

$2\text{KNO}_3 \longrightarrow$ +

$2\text{Ca}(\text{NO}_3)_2 \longrightarrow$ + + [4]

[Total: 17]


For
Examiner's
Use

7 The alkanes are generally unreactive. Their reactions include combustion, substitution and cracking.

For
Examiner's
Use

(a) The complete combustion of an alkane gives carbon dioxide and water.

(i) 10 cm³ of butane is mixed with 100 cm³ of oxygen, which is an excess. The mixture is ignited. What is the volume of unreacted oxygen left and what is the volume of carbon dioxide formed?



Volume of oxygen left = cm³

Volume of carbon dioxide formed = cm³ [2]

(ii) Why is the incomplete combustion of any alkane dangerous, particularly in an enclosed space?

.....
..... [2]

(b) The equation for a substitution reaction of butane is given below.



(i) Name the organic product.

..... [1]


(ii) This reaction does not need increased temperature or pressure. What is the essential reaction condition?

..... [1]

(iii) Write a different equation for a substitution reaction between butane and chlorine.

..... [1]

- (c) Alkenes are more reactive and industrially more useful than alkanes. They are made by cracking alkanes.



- (i) Draw the structural formula of the polymer poly(propene).

[2]

- (ii) Give the structural formula and name of the alcohol formed when but-1-ene reacts with steam.

name

[1]

structural formula

[1]

- (iii) Deduce the structural formula of the product formed when propene reacts with hydrogen chloride.

[1]

[Total: 12]

For
Examiner's
Use

DATA SHEET
The Periodic Table of the Elements

		Group																																																																																																				
I	II	III	IV	V	VI	VII	0					0																																																																																										
7 Li Lithium 3	9 Be Beryllium 4	1 H Hydrogen 1	11 B Boron 5	12 C Carbon 6	14 N Nitrogen 7	16 O Oxygen 8	19 F Fluorine 9	20 Ne Neon 10	13 Al Aluminium 13	14 Si Silicon 14	15 P Phosphorus 15	16 S Sulphur 16	17 Cl Chlorine 17	18 Ar Argon 18	27 Co Cobalt 27	28 Ni Nickel 28	29 Cu Copper 29	30 Zn Zinc 30	31 Ga Gallium 31	32 Ge Germanium 32	33 As Arsenic 33	34 Se Selenium 34	35 Br Bromine 35	36 Kr Krypton 36	41 Nb Niobium 41	42 Mo Molybdenum 42	43 Tc Technetium 43	44 Ru Ruthenium 44	45 Rh Rhodium 45	46 Pd Palladium 46	47 Ag Silver 47	48 Cd Cadmium 48	49 In Indium 49	50 Sn Tin 50	51 Sb Antimony 51	52 Te Tellurium 52	53 I Iodine 53	54 Xe Xenon 54	55 Rb Rubidium 37	56 Sr Strontium 38	57 Y Yttrium 39	58 Zr Zirconium 40	59 Ni Nickel 28	60 Co Cobalt 27	61 Fe Iron 26	62 Mn Manganese 25	63 Cr Chromium 24	64 V Vanadium 23	65 Ti Titanium 22	66 Sc Scandium 21	67 Ca Calcium 20	68 K Potassium 19	69 Ba Barium 56	70 La Lanthanum 57	71 Ce Cerium 58	72 Pr Praseodymium 59	73 Nd Neodymium 60	74 Pm Promethium 61	75 Sm Samarium 62	76 Eu Europium 63	77 Gd Gadolinium 64	78 Tb Terbium 65	79 Dy Dysprosium 66	80 Ho Holmium 67	81 Er Erbium 68	82 Tm Thulium 69	83 Yb Ytterbium 70	84 Lu Lutetium 71	85 Fr Francium 87	86 Ra Radium 88	87 Ac Actinium 89	88 Fr Francium 87	89 Ra Radium 88	90 Th Thorium 90	91 Pa Protactinium 91	92 U Uranium 92	93 Np Neptunium 93	94 Pu Plutonium 94	95 Am Americium 95	96 Cm Curium 96	97 Bk Berkelium 97	98 Cf Californium 98	99 Es Einsteinium 99	100 Fm Fermium 100	101 Md Mendelevium 101	102 No Nobelium 102	103 Lr Lawrencium 103	104 Rf Rutherfordium 104	105 Db Dubnium 105	106 Sg Seaborgium 106	107 Bh Bohrium 107	108 Hs Hassium 108	109 Mt Meitnerium 109	110 Ds Darmstadtium 110	111 Rg Roentgenium 111	112 Cn Copernicium 112	113 Nh Nihonium 113	114 Fl Flerovium 114	115 Mc Moscovium 115	116 Lv Livermorium 116	117 Ts Tennessine 117	118 Og Oganesson 118

*58-71 Lanthanoid series
†90-103 Actinoid series

Key

a	X	a = relative atomic mass
	X	X = atomic symbol
b		b = proton (atomic) number

The volume of one mole of any gas is 24 dm³ at room temperature and pressure (r.t.p.).

Permission to reproduce items where third-party owned material protected by copyright is included has been sought and cleared where possible. Every reasonable effort has been made by the publisher (UCLES) to trace copyright holders, but if any items requiring clearance have unwittingly been included, the publisher will be pleased to make amends at the earliest possible opportunity.

University of Cambridge International Examinations is part of the Cambridge Assessment Group. Cambridge Assessment is the brand name of University of Cambridge Local Examinations Syndicate (UCLES), which is itself a department of the University of Cambridge.