

# **Cambridge IGCSE**<sup>™</sup>

CANDIDATE NAME					
CENTRE NUMBER			CANDIDATE NUMBER		

**CHEMISTRY** 0620/41

Paper 4 Theory (Extended)

May/June 2022

1 hour 15 minutes

You must answer on the question paper.

No additional materials are needed.

#### **INSTRUCTIONS**

- Answer all questions.
- Use a black or dark blue pen. You may use an HB pencil for any diagrams or graphs.
- Write your name, centre number and candidate number in the boxes at the top of the page.
- Write your answer to each question in the space provided.
- Do not use an erasable pen or correction fluid.
- Do not write on any bar codes.
- You may use a calculator.
- You should show all your working and use appropriate units.

#### **INFORMATION**

- The total mark for this paper is 80.
- The number of marks for each question or part question is shown in brackets [ ].
- The Periodic Table is printed in the question paper.

chlorine

diamond

ethanol

carbon dioxide

1 A list of substances is shown.

aluminium oxide

	glucose	iron(III) oxide	limestone	nitrogen	oxygen						
Ans	Answer the questions using the list of substances.										
Eac	Each substance may be used once, more than once or not at all.										
Sta	State which of the substances:										
(a)	is a reactant in p	photosynthesis									
					[1]						
(I-)	is the series come	ditue at at harmite									
(D)	is the main cons	stituent of bauxite			[4]						
					[1]						
(c)	are <b>two</b> product	s of fermentation									
			and		[2]						
(-I\	in word on a fivel										
(a)	is used as a fuel				[4]						
					[1]						
(e)	is a gas used to	convert iron into steel									
					[1]						
<b>(£</b> )	is a greenhouse	900									
(1)	is a greenhouse	yas			[4]						
					[1]						
(g)	is a gas that is a	approximately 78% of cle	an, dry air								
					[1]						
(L.)	: f f										
(n)	is a form of carb	oon.			F.43						
					[1]						
					[Total: 9]						

2	(a)	Atoms are made of protons, neutrons and electrons. Atoms of the same element are known as isotopes.

(i) Complete the table.

particle	relative charge	relative mass
electron		1 1840
neutron		
proton	+1	

[2]

(ii)  $^{24}_{12}$ Mg and  $^{25}_{12}$ Mg are isotopes of magnesium.

Complete the table to show the numbers of electrons, neutrons and protons in these isotopes of magnesium.

isotope	number of electrons	number of neutrons	number of protons
<sup>24</sup> Mg			
<sup>25</sup> Mg			

[2]

	(iii)	Explain why magnesium ions have a charge of 2+.	
			[1]
(b)	Mg	<sup>2+</sup> ions have the electronic structure 2,8.	
	Giv ion	ve the formula of the following particles which have the same electronic structure as ${\tt N}$ s.	∕lg²⁺
	•	a cation (positive ion)	
	•	an anion (negative ion)	
	•	an atom	
			[3]

[Total: 8]

			4									
	This	que	estion is about sodium and compounds of sodium.									
(	(a)	(i)	Describe the bonding in a metallic element such as sodium.									
			You may include a diagram as part of your answer.									
				[3]								
	(	(ii)	Describe how solid sodium conducts electricity.									
				[1]								
	(b)	Sor	ne properties of sodium chloride are shown:									
		•	melting point of 801 °C									
		•	non-conductor of electricity when solid conductor of electricity when molten									
		•	soluble in water.									
		(i)	Name the type of bonding in sodium chloride.									
				[1]								
	(	(ii)	Explain why sodium chloride conducts electricity when molten.									

......[1]

(c)			nt determines the concentration of a solution of dilute sulfuric acid, $\rm H_2SO_4$ , by titration leous sodium hydroxide, NaOH.							
	ste	p 1	25.0 cm³ of 0.200 mol/dm³ NaOH is transferred into a conical flask.							
	ste	<b>p</b> 2	Three drops of methyl orange indicator are added to the conical flask.							
	ste	р 3	A burette is filled with H <sub>2</sub> SO <sub>4</sub> .							
	ste	p 4	The acid in the burette is added to the conical flask until the indicator changes colour. The volume of acid is recorded. This process is known as titration.							
	ste	p 5	The titration is repeated several times until a suitable number of results is obtained.							
	(i)		ne the piece of apparatus used to measure exactly 25.0 cm³ of 0.200 mol/dm³ NaOH tep 1.							
			[1]							
	(ii)	Sta	te the colour change of the methyl orange indicator in <b>step 4</b> .							
		fron	n to [1]							
(	(iii)	Sta	te how the student decides that a suitable number of results have been obtained.							
			[1]							
(	iv)	20.0	0 cm <sup>3</sup> of H <sub>2</sub> SO <sub>4</sub> reacts with 25.0 cm <sup>3</sup> of 0.200 mol/dm <sup>3</sup> NaOH.							
		The	equation for the reaction is shown.							
			$H_2SO_4 + 2NaOH \rightarrow Na_2SO_4 + 2H_2O$							
		Cal	culate the concentration of H <sub>2</sub> SO <sub>4</sub> using the following steps.							
		•	Calculate the number of moles in 25.0 cm <sup>3</sup> of 0.200 mol/dm <sup>3</sup> NaOH.							
			mol							
		•	Determine the number of moles of H <sub>2</sub> SO <sub>4</sub> that react with the NaOH.							
			mol							
		•	Calculate the concentration of H <sub>2</sub> SO <sub>4</sub> .							
			mol/dm³ [3]							
			[Total: 12]							

<b>4</b> T	his	question	is	about	com	pounds	of	sulfur
------------	-----	----------	----	-------	-----	--------	----	--------

(a)	Sulfuric action four stage	, , ,	is manufactured	using the	Contact process.	This manufacture	involves
	otogo 1	Maltan ault	fur burne in eir te	produce	ulfur diovido		

**stage 1** Molten sulfur burns in air to produce sulfur dioxide.

stage 2 Sulfur dioxide reacts with oxygen to form sulfur trioxide.

stage 3 Sulfur trioxide combines with concentrated sulfuric acid to form oleum, H<sub>2</sub>S<sub>2</sub>O<sub>7</sub>.

stage 4 Oleum reacts to form concentrated sulfuric acid.

(i)	Write a chemical equation for the reaction occurring in <b>stage 1</b> .	
		[1]

(ii) State the essential conditions that are necessary for stage 2. Write an equation for the chemical reaction that occurs.


(iii) Write a chemical equation for the reaction occurring in **stage 3**.

 	 	 		 	 	[1	J
 		 _	_				

(iv) Name the substance that reacts with oleum in stage 4.

**(b)** Hydrogen sulfide has the formula H<sub>2</sub>S.

(i) Complete the dot-and-cross diagram to show the electron arrangement in a molecule of hydrogen sulfide. Show outer shell electrons only.



[2]

(ii) Balance the chemical equation for the reaction of hydrogen sulfide with sulfur dioxide shown.

....
$$H_2S + SO_2 \rightarrow .....S + .....H_2O$$
 [1]

[Total: 10]

[2]

5	Eth	anoic acid is manufactured	by the reaction of methanol w	vith carbon monoxide.	
	An	equilibrium mixture is produ	ced.		
		CH <sub>3</sub>	$OH(g) + CO(g) \rightleftharpoons CH_3CO^{-1}$	OH(g)	
	(a)	State <b>two</b> characteristics o	f an equilibrium.		
		1			
		2			
					[2]
	(b)	The purpose of the industriof reaction.	ial process is to produce a hiç	gh yield of ethanoic acid at a	high rate
		The manufacture is carried	out at a temperature of 300°	°C.	
		The forward reaction is exc	othermic.		
		Use this information to stat	e why the manufacture is <b>not</b>	t carried out at temperatures:	
		• below 300 °C			
		• above 300 °C.			
					[2]
					[-]
	(c)	Complete the table using of	only the words increases, deci	reases or no change.	
			effect on the rate of the forward reaction	effect on the equilibrium yield of CH <sub>3</sub> COOH(g)	
		adding a catalyst		no change	
		decreasing the pressure			
					[3]
	(d)	Suggest which of the follow your answer.	ving metals is a suitable catal	lyst for the reaction. Give a re	ason for
		aluminium calo	cium cobalt magne	esium potassium	
		suitable catalyst			

State t	he general formula of this homologous series.
	[1]
	he structure of the carboxylic acid containing three carbon atoms. Show all of the atoms of the bonds.
	[2]
(g) When	carboxylic acids react with alcohols, esters are produced.
The for	rmula of ester <b>X</b> is CH <sub>3</sub> CH <sub>2</sub> COOCH <sub>3</sub> .
<b>(i)</b> Na	ame ester X.
	[1]
(ii) Gi	ve the name of the carboxylic acid and the alcohol that react together to produce ester ${f X}$ .
ca	ırboxylic acid
ald	cohol
	[2]
(h) Ester	f has the following composition by mass:
C, 48.6	65%; H, 8.11%; O, 43.24%.
Calcula	ate the empirical formula of ester <b>Y</b> .
	empirical formula =[3]

(i)	Ester <b>Z</b> has the empirical formula C <sub>2</sub> H <sub>4</sub> O and a relative molecular mass of 88.

Determine the molecular formula of ester **Z**.

molecular formula = ..... [1]

[Total: 19]

6 This question is about zinc and its com	pounds.
---	---------

- 1	~ I	7ing is oversate	d from ita	ara which ic	moinly zino	aulfida 7nC
١	a	Zinc is extracted	1 110111 ILS	OLE MILICITIES	illallly Zillo	Sullide, ZIIS.

The steps for this extraction are shown.

**step 1** Zinc sulfide is converted into zinc oxide.

**step 2** The zinc oxide is then reduced to zinc in a furnace. The zinc formed becomes a gas.

**step 3** The zinc gas is cooled to form molten zinc.

(i) Name the ore of zinc, which is mainly zinc sulfide.

.....[1]

(ii) Describe how zinc sulfide is converted into zinc oxide in step 1.

......[1]

(iii) Name the reducing agent used in step 2.

.....[1]

(iv) Explain why the zinc forms a gas in **step 2** inside the furnace.

.....[1]

(v) State the name of the physical change occurring when zinc gas is converted into molten zinc.

\_\_\_\_\_\_[1

**(b)** Zinc sulfate crystals, ZnSO<sub>4</sub>•7H<sub>2</sub>O, are hydrated.

Zinc sulfate crystals are made by reacting zinc carbonate with dilute sulfuric acid.

The equation for the overall process is shown.

$$ZnCO_3 + H_2SO_4 + 6H_2O \rightarrow ZnSO_4 \cdot 7H_2O + CO_2$$

**step 1** Large pieces of solid zinc carbonate are added to dilute sulfuric acid until the zinc carbonate is in excess. This forms aqueous zinc sulfate.

**step 2** The excess zinc carbonate is separated from the aqueous zinc sulfate.

**step 3** The aqueous zinc sulfate is heated until a saturated solution is formed.

**step 4** The saturated solution is allowed to cool and crystallise.

**step 5** The crystals are removed and dried.

In <b>step 1</b> , zinc carbonate is in excess when no more zinc carbonate dissolves.
State one <b>other</b> observation that indicates the zinc carbonate is in excess in <b>step 1</b> .
[1]
Name a different substance, other than zinc carbonate, that can be added to dilute sulfuric acid to produce aqueous zinc sulfate in <b>step 1</b> .
[1]
Step 1 is repeated using powdered zinc carbonate instead of large pieces.
All other conditions are kept the same.
The rate of reaction increases.
Give a reason why the rate of reaction increases. Explain your answer in terms of particles.
[2]
Suggest what is observed when the solution is saturated in step 3.
[1]
The formula of zinc sulfate crystals is ZnSO <sub>4</sub> •7H <sub>2</sub> O.
Give the formula of the solid formed if the crystals are heated to dryness in <b>step 3</b> .
[1]
[Total: 11]

7 The Periodic Table can be used to classify elements
---

- (a) Group I elements react with cold water to form alkaline solutions.
  - (i) Place the Group I elements caesium, lithium, potassium, rubidium and sodium in their order of reactivity with water.

Put the most reactive element first.

mo	st reactive   ▶ least reactive
	[1]
	(ii) Name the alkaline solution formed when caesium reacts with cold water.
	[1]
(b)	Group I elements have lower melting points than transition elements.
	Describe one <b>other</b> difference in the <b>physical</b> properties of Group I elements and transition elements.
	[1]
(c)	Group VII elements are known as the halogens.
	Astatine is below iodine in Group VII.
	Predict the physical state of astatine at room temperature and pressure.
	[1]
(d)	Some Group VII elements react with aqueous solutions containing halide ions.
	When aqueous chlorine is added to aqueous potassium bromide a reaction occurs.
	The ionic half-equations for the reaction are shown.
	$\mathrm{C}l_2(\mathrm{aq})$ + 2e $^  o$ 2C $l^-(\mathrm{aq})$
	$2Br^{-}(aq) \rightarrow Br_{2}(aq) + 2e^{-}$

[2]

original colour of potassium bromide solution .....

final colour of reaction mixture .....

© UCLES 2022 0620/41/M/J/22

(i) Describe the colour change of the solution.

	(ii)	Identify the specie	es that is oxid	ised.							
		Explain your decis	sion.								
		species oxidised									
explanation											
		Oxpidilation			[2]						
(e)		omine monochloride emical equation is s		ade by the reaction between	en bromine and chlorine. The						
			$Br_2(g)$	$+ Cl_2(g) \rightarrow 2BrCl(g)$							
			bond	bond energy in kJ/mol							
			Br–Br	190							
			Cl-Cl	242							
			Br–C1	218							
	Cal	culate the overall e	energy change	e for the reaction using bond	d energies.						
	Use	e the following step	ıs.								
				a compared to the column	hands in Amela of Davis, and						
	•	1 mole of $Cl_2(g)$ .	amount of e	nergy required to break the	bonds in 1 mole of Br <sub>2</sub> (g) and						
					kJ						
			l = == =								
	•	formed.	ii amount of e	energy released when the b	onds in 2 moles of BrC l(g) are						
					kJ						
		Calculate the ave	roll on oran colo	and for the reaction	•						
	•	Calculate the ove	rall energy ch	ange for the reaction.							
					kJ/mol [3]						
					[Total: 11]						

14

## **BLANK PAGE**

15

### **BLANK PAGE**

Permission to reproduce items where third-party owned material protected by copyright is included has been sought and cleared where possible. Every reasonable effort has been made by the publisher (UCLES) to trace copyright holders, but if any items requiring clearance have unwittingly been included, the publisher will be pleased to make amends at the earliest possible opportunity.

To avoid the issue of disclosure of answer-related information to candidates, all copyright acknowledgements are reproduced online in the Cambridge Assessment International Education Copyright Acknowledgements Booklet. This is produced for each series of examinations and is freely available to download at www.cambridgeinternational.org after the live examination series.

Cambridge Assessment International Education is part of Cambridge Assessment. Cambridge Assessment is the brand name of the University of Cambridge Local Examinations Syndicate (UCLES), which is a department of the University of Cambridge.

The Periodic Table of Elements

	=	2	우	elium 4	10	Je Je	neon 20	18	٩Ľ	argon 40	36	궃	ypton 84	54	×e	enon 131	98	٦	adon -			
			_																			
	₹				6	ш	fluorine 19	17	Cl	chlorine 35.5	35	Ŗ	bromine 80	53	Ι	iodine 127	85	Ą	astatine _			
	5				80	0	oxygen 16	16	ഗ	sulfur 32	34	Se	selenium 79	52	<u>a</u>	tellurium 128	84	Ъ	moloulum —	116	_	livermorium -
	>				7	z	nitrogen 14	15	₾	phosphorus 31	33	As	arsenic 75	51	Sb	antimony 122	83	<u>B</u>	bismuth 209			
	2				9	ပ	carbon 12	14	: <u>S</u>	silicon 28	32	Ge	germanium 73	20	Sn	tin 119	82	Pb	lead 207	114	Εl	flerovium
	=				22	Δ	boron 11	13	Αl	aluminium 27	31	Ga	gallium 70	49	In	indium 115	84	11	thallium 204			
											30	Zu	zinc 65	48	g	cadmium 112	80	Нg	mercury 201	112	S	copernicium
											29	ŋ	copper 64	47	Ag	silver 108	79	Au	gold 197	111	Rg	roentgenium
dn											28	Z	nickel 59	46	Pd	palladium 106	78	Ŧ	platinum 195	110	Ds	darmstadtium -
Group											27	ပိ	cobalt 59	45	뫈	rhodium 103	11	Ir	iridium 192	109	¥	meitnerium -
		-	I	hydrogen 1							26	Ь	iron 56	44	Ru	ruthenium 101	92	SO	osmium 190	108	Η	hassium
					J						25	Mn	manganese 55	43	ပ	technetium -	75	Re	rhenium 186	107	Bh	bohrium
						Го	ss				24	ပ်	chromium 52	42	Mo	molybdenum 96	74	>	tungsten 184	106	Sg	seaborgium
				Key	atomic number	atomic symbo	name relative atomic mass				23	>	vanadium 51	41	q	niobium 93	73	<u>ra</u>	tantalum 181	105	o O	dubnium
					at	ator	relati				22	j	titanium 48	40	Zr	zirconium 91	72	士	hafnium 178	104	¥	rutherfordium -
								J			21	Sc	scandium 45	39	>	yttrium 89	57-71	lanthanoids		89–103	actinoids	
	=				4	Be	beryllium 9	12	Mg	magnesium 24	20	Ca	calcium 40	38	ഗ്	strontium 88	56	Ва	barium 137	88	Ra	radium
	_				8	:=	lithium 7	7	Na	sodium 23	19	¥	potassium 39	37	Rb	rubidium 85	55	S	caesium 133	87	ь Б	francium

71 Lu	lutetium 175	103	۲	lawrencium -	
70 Yb				_	
e9 Tm	thulium 169	101	Md	mendelevium -	
89 Er	erbium 167	100	Fm	fermium -	
67 79	holmium 165	66	Es	einsteinium –	
66 Dy	dysprosium 163	86	ŭ	californium -	
65 Tb	terbium 159	26	Ř	berkelium -	
Gd Gd	gadolinium 157	96	Cm	curium	
® Eu	europium 152	92	Am	americium	
Sm	samarium 150	94	Pu	plutonium	
61 Pm	promethium —	93	dN	neptunium	
9 <b>P</b>	neodymium 144	92	$\supset$	uranium 238	2007
59 Pr	praseodymium 141	91	Ра	protactinium 23.1	- 03
Ce Ce	cerium 140	06	Ч	thorium 232	202
57 <b>La</b>	lanthanum 139	88	Ac	actinium	1

lanthanoids

actinoids

The volume of one mole of any gas is 24 dm<sup>3</sup> at room temperature and pressure (r.t.p.).