Cambridge IGCSE[™]

CANDIDATE NAME					
CENTRE NUMBER			CANDIDATE NUMBER		

CHEMISTRY 0620/43

Paper 4 Theory (Extended)

October/November 2022

1 hour 15 minutes

You must answer on the question paper.

No additional materials are needed.

INSTRUCTIONS

- Answer all questions.
- Use a black or dark blue pen. You may use an HB pencil for any diagrams or graphs.
- Write your name, centre number and candidate number in the boxes at the top of the page.
- Write your answer to each question in the space provided.
- Do not use an erasable pen or correction fluid.
- Do not write on any bar codes.
- You may use a calculator.
- You should show all your working and use appropriate units.

INFORMATION

- The total mark for this paper is 80.
- The number of marks for each question or part question is shown in brackets [].
- The Periodic Table is printed in the question paper.

- 1 Atoms and ions are made from small particles called electrons, neutrons and protons.
 - (a) Complete the table.

particle	relative charge	relative mass
electron	-1	<u>1</u> 1840
neutron		
proton		

[2]

(b) Information about atoms and ions, A, B and C, is shown in the table.

Complete the table.

atom or ion	number of electrons	number of neutrons	number of protons	symbol
Α	18		20	⁴² Ca ²⁺
В		18		³⁵ C <i>l</i>
С	18	16	16	

[6]

[Total: 8]

2 The table shows the melting points, boiling points and electrical conductivities of six substances, $\bf D$, $\bf E$, $\bf F$, $\bf G$, $\bf H$ and $\bf I$.

substance	melting point /°C	boiling point /°C	conducts electricity when solid	conducts electricity when liquid
D	1083	2567	yes	yes
Е	-117	79	no	no
F	3550	4827	no	no
G	119	445	no	no
Н	-210	– 196	no	no
i	801	1413	no	yes

(a)	Identify the substance, D, E, F, G, H or I, which is:					
	(i)	a liquid at 25 °C	[1]			
	(ii)	a gas at 25°C	[1]			
	(iii)	a solid consisting of simple molecules at 25 °C	[1]			
(b)		ntify the substance, D , E , F , G , H or I , which is a metal. Give a reason for your choice.				
		stance				
	rea	son[[2]			
(c)		ntify the substance, \mathbf{D} , \mathbf{E} , \mathbf{F} , \mathbf{G} , \mathbf{H} or \mathbf{I} , which has a macromolecular structure. Give \mathbf{tw} sons for your choice.	/ O			
	sub	stance				
	rea	son 1				
	rea	son 2	 [3]			
		·	•			
(d)	Ide	ntify the substance, D , E , F , G , H or I , which is an ionic solid. Give a reason for your choic	e.			
	sub	stance				
	rea	son				
			 [2]			

[Total: 10]

3

Alur	miniu	um is extracted from its ore by electrolysis.	
(a)	Nar	ne the ore of aluminium which consists mainly of aluminium oxide.	
			[1]
(b)	Sta	te what is meant by the term <i>electrolysis</i> .	
			[2]
(c)	Ele	ctrolysis is carried out on aluminium oxide dissolved in molten cryolite.	
		waste gases	
n	egai	positive electrode aluminium oxide dissolved in molten cryolite aluminium	
	(i)	Give two reasons why the electrolysis is carried out on aluminium oxide dissolved molten cryolite instead of electrolysing molten aluminium oxide only.	l in
		1	
		2	 [2]
	(ii)	Write the ionic half-equation for the reaction occurring at the negative electrode.	
			[2]
(iii)	The positive electrodes are made of carbon.	
		Explain why the positive carbon electrodes are replaced regularly.	

(d)	Aluı	Aluminium is more reactive than copper.					
	When aluminium is added to aqueous $copper(II)$ sulfate, no immediate reaction is seen.						
	Exp	plain why.					
		[1]					
(e)	Aluı	minium reacts with oxygen to form an amphoteric oxide.					
	(i)	State what is meant by the term amphoteric.					
		[1]					
	(ii)	The reaction between aluminium oxide and aqueous sodium hydroxide forms a salt containing the negative ion AlO_2^- . The only other product is water.					
		Write a chemical equation for the reaction between aluminium oxide and aqueous sodium hydroxide.					
		[2]					
(f)	Gal	lium is in the same group as aluminium and forms similar compounds.					
	Pre	dict the formulae of:					
	gall	ium(III) chloride					
	gall	ium(III) sulfate.					
		[2]					
		[Total: 15]					

- 4 This question is about compounds of phosphorus.
 - (a) Gaseous phosphorus(V) chloride decomposes into gaseous phosphorus(III) chloride and gaseous chlorine.

When the three gases are present in a closed container the system reaches equilibrium.

$$PCl_5(g) \rightleftharpoons PCl_3(g) + Cl_2(g)$$

(i) Complete the table using only the words *increases*, *decreases* or *no change*.

	effect on the rate of the forward reaction	effect on the equilibrium yield (PC l_3 (g) and C l_2 (g))
increasing the temperature		increases
decreasing the pressure		
adding a catalyst		no change


[4]

(ii)	The table shows that when the temperature increases, the equilibrium yields of $PCl_3(g)$
	and $Cl_2(g)$ increase.

State what conclusion can be made from this.

•	C 4 T
	ı 1 I

(b) Complete the dot-and-cross diagram to show the electron arrangement in a molecule of phosphorus(III) chloride, PCl_3 . Show outer shell electrons only.



[2]

(c)	Phosphorus oxychloride has the formula $POCl_3$.
	Phosphorus oxychloride is the only product of the reaction between phosphorus(V) chloride, PCl_5 , and phosphorus(V) oxide, P_4O_{10} .
	Write a chemical equation for the reaction between phosphorus $\!\!(V)$ chloride and phosphorus $\!\!(V)$ oxide.
	[2]
(d)	Compound X has the following composition by mass.
	H, 3.66%; P, 37.80%; O, 58.54%
	Calculate the empirical formula of compound X .
	empirical formula = [2]
(e)	Compound Y has the empirical formula H ₃ PO ₄ and a relative molecular mass of 98.
	Deduce the molecular formula of compound Y.
	made avilor formavilo
	molecular formula =[1]
	[Total: 12]

8


BLANK PAGE

5	This q	uestion	is about sulfuric acid, H ₂ SO ₄ , and salts that can be made from sulfuric acid.
	(a) S	ulfuric a	cid is manufactured by the Contact process.
	st	tage 1	Molten sulfur burns in air to produce sulfur dioxide.
	st	tage 2	Sulfur dioxide reacts with oxygen to form sulfur trioxide, SO ₃ .
	st	tage 3	Sulfur trioxide reacts with concentrated sulfuric acid to form oleum, $H_2S_2O_7$.
	st	tage 4	Oleum is converted into sulfuric acid.
	(i)) The	equation for the reaction in stage 2 is shown.
			$2SO_2 + O_2 \rightleftharpoons 2SO_3$
		State	e the temperature and pressure used in stage 2.
		Nam	e the catalyst used in stage 2 .
		temp	erature°C
		press	sure atn
		catal	yst
	(ii)) Write	e the chemical equation for the reaction in stage 3 .
	(iii)) Nam	e the substance that reacts with oleum in stage 4 .
			[1
		ame the	e black solid that is produced when concentrated sulfuric acid is added to sugar

(c) Dilute sulfuric acid and aqueous potassium hydroxide are used to make aqueous potassium sulfate.

$$H_2SO_4 + 2KOH \rightarrow K_2SO_4 + 2H_2O$$

The method includes use of the following apparatus.



(i) Calculate the volume of $0.0625\,\mathrm{mol/dm^3}$ dilute sulfuric acid, $\mathrm{H_2SO_4}$, that completely reacts with $25.0\,\mathrm{cm^3}$ of $0.100\,\mathrm{mol/dm^3}$ potassium hydroxide, KOH, to produce aqueous potassium sulfate.

Use the following steps.

• Calculate the number of moles of KOH in 25.0 cm³ of 0.100 mol/dm³ KOH.

= mol

• Deduce the number of moles of H₂SO₄ that react with KOH.

= mol

•	Calculate	the volume	of H ₂ SO ₄	required.
---	-----------	------------	-----------------------------------	-----------

volume =	cm ³ [3]
(ii) The experiment is repeated using the same volume and concentration of potal hydroxide and the same concentration of dilute sulfuric acid. In this second experiment the product is aqueous potassium hydrogensulfate, KHSO ₄ .	
$H_2SO_4 + KOH \rightarrow KHSO_4 + H_2O$	
Use your answer to (c)(i) and the equation to deduce the volume of H ₂ SO ₄ required	
volume = ci	n³ [1]
(d) Aqueous potassium hydrogensulfate, KHSO ₄ (aq), contains the ions K ⁺ (aq), H ⁺ (aq SO ₄ ²⁻ (aq).) and
Describe the observations in the following tests.	
(i) A flame test is carried out on aqueous potassium hydrogensulfate.	
	[1]
(ii) Solid copper(II) carbonate is added to aqueous potassium hydrogensulfate.	
	[2]
(iii) An acidic solution containing aqueous barium ions, Ba ²⁺ (aq), is added to aq potassium hydrogensulfate.	ueous
	[1]
(e) Write the ionic equation for the reaction in (d)(iii).	
Include state symbols.	
	[3]
[Tot	al: 17]

6 (a) Chloroethene ($CH_2=CHCl$) can be manufactured from 1,2-dichloroethane (CH_2ClCH_2Cl).

The equation can be represented as shown.

$$\begin{array}{c|ccccc} Cl & H & & Cl & H \\ & & & & & | & & | \\ H - C - C - Cl & \rightarrow & C = C & + & H - Cl \\ & & & & | & & | & | \\ & & & & & H & H & H \end{array}$$

(i) Some bond energies are given.

bond	bond energy in kJ/mol
C–C	350
C=C	610
C-C1	340
C–H	410
H–C1	430

Use the bond energies in the table to calculate the energy change, in kJ/mol, of the reaction.

Use the following steps.

Calculate the energy needed to break bonds.

• Calculate the energy released when bonds form.

 Ca 	alculate the	e energy	change	of the	reaction
------------------------	--------------	----------	--------	--------	----------

	energy change of the reaction =kJ/	mol [3]
(ii)	Deduce whether the energy change for this reaction is exothermic or endothermic.	
	Give a reason for your answer.	
		[1]
Pai	rt of a synthetic polymer is shown.	
	$\begin{array}{cccccccccccccccccccccccccccccccccccc$	

(i)	State the number	of	monomer	units	that	are	needed	to	make	the	part	of	the	polym	ner
	shown.														

[1]

(ii) Name and draw the structure of the monomer used to make this polymer. Show all of the atoms and all of the bonds.

name	 	 	 	

structure

(b)

(iii) State the empirical formula of the polymer.



(c)	Pro	teins are natural polymers.
		teins are broken down into amino acids. The process is similar to how complex bohydrates are broken down to give simple sugars.
	(i)	Name the type of reaction in which proteins are broken down into amino acids.
		[1]
	(ii)	Name two types of substance that are used to break down proteins into amino acids.
		1
		2[2]
	(iii)	Amino acids are colourless.
		A sample containing a mixture of amino acids is separated. Each amino acid is detected and identified.
		Name the process used to separate the amino acids.
		Name the type of substance used to detect the amino acids.
		Give the symbol of the value used to determine the identity of each amino acid after separation and detection.
		[3]
(d)	Pro	teins are natural polymers. Proteins contain amide linkages.
	Syr	nthetic polyamides also contain amide linkages.
	(i)	Name a synthetic polyamide.
		[1]
	(ii)	Identify the two functional groups present in the monomers used to produce synthetic polyamides.
		1
		2
		[2]
		[Total: 18]

15

BLANK PAGE

Permission to reproduce items where third-party owned material protected by copyright is included has been sought and cleared where possible. Every reasonable effort has been made by the publisher (UCLES) to trace copyright holders, but if any items requiring clearance have unwittingly been included, the publisher will be pleased to make amends at the earliest possible opportunity.

To avoid the issue of disclosure of answer-related information to candidates, all copyright acknowledgements are reproduced online in the Cambridge Assessment International Education Copyright Acknowledgements Booklet. This is produced for each series of examinations and is freely available to download at www.cambridgeinternational.org after the live examination series.

Cambridge Assessment International Education is part of Cambridge Assessment. Cambridge Assessment is the brand name of the University of Cambridge Local Examinations Syndicate (UCLES), which is a department of the University of Cambridge.

The Periodic Table of Elements

	=	2	우	elium 4	10	Je Je	neon 20	18	٩Ľ	argon 40	36	궃	ypton 84	54	×e	enon 131	98	٦	adon -			
			_																			
	₹				6	ш	fluorine 19	17	Cl	chlorine 35.5	35	Ŗ	bromine 80	53	Ι	iodine 127	85	Ą	astatine _			
	5				80	0	oxygen 16	16	ഗ	sulfur 32	34	Se	selenium 79	52	<u>a</u>	tellurium 128	84	Ъ	moloulum —	116	_	livermorium -
	>				7	z	nitrogen 14	15	₾	phosphorus 31	33	As	arsenic 75	51	Sb	antimony 122	83	<u>B</u>	bismuth 209			
	2				9	O	carbon 12	14	: <u>S</u>	silicon 28	32	Ge	germanium 73	90	Sn	tin 119	82	Pb	lead 207	114	Εl	flerovium
	=				22	Δ	boron 11	13	Αl	aluminium 27	31	Ga	gallium 70	49	In	indium 115	84	11	thallium 204			
											30	Zu	zinc 65	48	g	cadmium 112	80	Нg	mercury 201	112	S	copernicium
											29	n	copper 64	47	Ag	silver 108	79	Αn	gold 197	111	Rg	roentgenium
dn											28	Z	nickel 59	46	Pd	palladium 106	78	Ŧ	platinum 195	110	Ds	darmstadtium -
Group											27	ပိ	cobalt 59	45	뫈	rhodium 103	11	Ir	indium 192	109	¥	meitnerium -
		-	I	hydrogen 1							26	Ь	iron 56	44	Ru	ruthenium 101	92	SO	osmium 190	108	Η	hassium
					J						25	Mn	manganese 55	43	ပ	technetium -	75	Re	rhenium 186	107	Bh	bohrium
						Го	ss				24	ပ်	chromium 52	42	Mo	molybdenum 96	74	>	tungsten 184	106	Sg	seaborgium
				Key	atomic number	atomic symbo	name relative atomic mass				23	>	vanadium 51	41	q	niobium 93	73	<u>ra</u>	tantalum 181	105	o O	dubnium
					at	ator	relati				22	j	titanium 48	40	Zr	zirconium 91	72	士	hafnium 178	104	¥	rutherfordium -
								J			21	Sc	scandium 45	39	>	yttrium 89	57-71	lanthanoids		89–103	actinoids	
	=				4	Be	beryllium 9	12	Mg	magnesium 24	20	Ca	calcium 40	38	ഗ്	strontium 88	56	Ва	barium 137	88	Ra	radium
	_				8	:=	lithium 7	7	Na	sodium 23	19	¥	potassium 39	37	Rb	rubidium 85	55	S	caesium 133	87	ь Б	francium

71 Lu	lutetium 175	103	۲	lawrendum -
70 Yb				_
°9 Tm	thulium 169	101	Md	mendelevium -
68 Er	erbium 167	100	Fm	fermium -
67 H	holmium 165	66	Es	einsteinium
。 O	dysprosium 163	86	₽	californium -
65 Tb	terbium 159	26	B	berkelium
² Od	gadolinium 157	96	Cm	curium
ы Еu	europium 152	92	Am	americium -
62 Sm	samarium 150	94	Pu	plutonium -
Pm	promethium	93	ď	neptunium _
9 9 8	neodymium 144	92	\supset	uranium 238
59 P	praseodymium 141	91	Ра	protactinium 231
Çe O	cerium 140	06	드	thorium 232
57 La	lanthanum 139	88	Ac	actinium

lanthanoids

actinoids

The volume of one mole of any gas is 24 dm³ at room temperature and pressure (r.t.p.).