



## Cambridge International Examinations

Cambridge International General Certificate of Secondary Education

CANDIDATE NAME				
CENTRE NUMBER		CANDIDATE NUMBER		

CHEMISTRY 0620/42

Paper 4 Theory (Extended)

May/June 2017

1 hour 15 minutes

Candidates answer on the Question Paper.

No Additional Materials are required.

## **READ THESE INSTRUCTIONS FIRST**

Write your Centre number, candidate number and name on all the work you hand in.

Write in dark blue or black pen.

You may use an HB pencil for any diagrams or graphs.

Do not use staples, paper clips, glue or correction fluid.

DO NOT WRITE IN ANY BARCODES.

Answer all questions.

Electronic calculators may be used.

A copy of the Periodic Table is printed on page 16.

You may lose marks if you do not show your working or if you do not use appropriate units.

At the end of the examination, fasten all your work securely together.

The number of marks is given in brackets [ ] at the end of each question or part question.

The syllabus is approved for use in England, Wales and Northern Ireland as a Cambridge International Level 1/Level 2 Certificate.



1	(a) Sta	te the name of the process that is used to
	(i)	separate oxygen from liquid air,
	<i>a</i>	[1]
	(ii)	separate the individual dyes in ink,[1]
	(iii)	produce ethanol from simple sugars,
		[1]
	(iv)	obtain water from aqueous sodium chloride,
		[1]
	(v)	separate the precipitate formed when aqueous silver nitrate is added to aqueous sodium chloride.
		[1]
	(h) Sta	te what is meant by the terms
	(i)	element,
		[1]
	(ii)	compound,
		[1]
	(iii)	ion.
		[1]
		[Total: 8]

(a) D	efine the term isotopes.			
 (b) C	omplete the following table which give		it carbon atoms and si	
(6)	simplete the following table which give	carbon	silicon	
	proton number	Carbon	Silicon	
	electronic structure			
	nucleon number	12	28	
	number of neutrons in one atom			
				[3]
( <b>c)</b> Si	licon has a giant structure which is si	milar to the structur	e of diamond.	
(i)	Name the type of bond which is pro	esent between silic	on atoms in silicon.	
				[1]
(ii)	Suggest <b>two</b> physical properties of Use your knowledge of structure a properties.		lain why silicon has th	hese physical
	property 1			
	reason 1			
	property 2			
	reason 2			
(4) <i>C</i>	amples of air taken from industri	al areas are foli	nd to contain small	[4]
	amples of all taken from industri- arbon monoxide.	ai aieas aie ioui	nd to contain smail	amounts of
(i)	Explain how this carbon monoxide	is formed.		
				[2]
(ii)	State why carbon monoxide should	d <b>not</b> be inhaled.		
				[1]

(e)		rbon dioxide, $CO_2$ , is a gas at room temperature and pressure, whereas silicon(IV) oxide, $O_2$ , is a solid.
	(i)	Name the type of structure which the following compounds have.
		carbon dioxide[1]
		silicon(IV) oxide[1]
	(ii)	Use your knowledge of structure and bonding to explain why carbon dioxide is a gas at room temperature and pressure, whereas silicon(IV) oxide is a solid.
		[3]
(f)		con(IV) oxide is an acidic oxide. When silicon(IV) oxide reacts with alkalis, the salts formed stain the ion ${\rm SiO_3^{2-}}$ .
		te a chemical equation for the reaction between $\operatorname{silicon}(\operatorname{IV})$ oxide and aqueous lium hydroxide.
		[2]
		[Total: 20]

3 This question is about nitrogen and some of its compound
--

(a)	Nitrogen in the air can be converted into ammonia by the Haber process. The chemical equation
	for the reaction is shown

$$N_2 + 3H_2 \rightleftharpoons 2NH_3$$

(i)	State the temperature and pressure used in the Haber process.			
	temperature			

(ii)	Name the catalyst used in the Haber process.

- **(b)** The ammonia produced in the Haber process can be oxidised to nitrogen(II) oxide at 900 °C. The reaction is exothermic.
  - (i) Balance the chemical equation for this reaction.

$$4NH_3 + ....O_2 \rightleftharpoons ....NO + ....H_2O$$
 [2]

(ii) Suggest a reason, other than cost, why a temperature greater than 900 °C is **not** used.

(iii) Suggest a reason why a temperature less than 900 °C is **not** used.

(c) Nitrogen(II) oxide can be reacted with oxygen and water to produce nitric acid as the only product.

Write a chemical equation for this reaction.

d) Describe how you would prepare a pure dry sample of copper(II) nitrate crystals in the laboratory using dilute nitric acid and solid copper(II) carbonate. Include a series of key steps in your answer. You should include a chemical equation for the reaction.
[6]
[Total: 15]

Question 4 starts on the next page.

4	Nickel, copper and zinc are three consecutive elements in the Periodic Table.				
	(a)	Nickel and copper	are transition elements.		
		State three chemic	al properties of transition elements.		
	(b)	Copper(II) oxide is insoluble in water.	s a basic oxide but zinc oxide is an a	amphoteric oxide. Both oxides are	
			vith a mixture of solid copper(II) oxide a sample of copper(II) oxide from this mi		
				[3]	
	(c)	Three cells are set	up each using two metals.		
		cell 1	cell 2	cell 3	
_ zinc		1.10 V + copp	0.51 V  er zinc nickel	copper	
		electrolyte	electrolyte	electrolyte	
		(i) Write the ionic	half-equation for the reaction occurring	at the zinc electrode in <b>cell 1</b> .	

(ii)	Put the <b>three</b> metals, copper, nickel and zinc, in order of reactivity.			
	most reactive			
	$\downarrow$			
	least reactive		[1]	
(iii)	•	labelling in <b>cell 3</b> by writing the polarity (+/–) of each electrode in the g the reading on the voltmeter.	circles [2]	
		[To	otal: 11]	

- 5 (a) The elements in Group VII are known as the halogens. Some halogens react with aqueous solutions of halides.
  - (i) Complete the table by adding a ✓ to indicate when a reaction occurs and a X to indicate when no reaction occurs.

	aqueous potassium chloride	aqueous potassium bromide	aqueous potassium iodide
chlorine	X	✓	
bromine		X	
iodine			X

[3]

	(ii)	Write a chemical equation for the reaction between chlorine and aqueous potassium bromide.
		[1]
(b)	silv	ample of vanadium chloride was weighed and dissolved in water. An excess of aqueous er nitrate, acidified with dilute nitric acid, was added. A precipitate of silver chloride was ned. The ionic equation for this reaction is shown.
		$Ag^{+}(aq) + Cl^{-}(aq) \rightarrow AgCl(s)$
	The	e mass of silver chloride formed was 2.87 g.
	(i)	State the colour of the precipitate of silver chloride.
		[1]

(ii) The relative formula mass of silver chloride, AgC *l*, is 143.5.

Calculate the number of moles in 2.87 g of AgCl.

moles of AgCl = ...... mol [1]

(iii) Use your answer to (b)(ii) and the ionic equation to deduce the number of moles of chloride ions, Cl<sup>-</sup>, that produced 2.87 g of AgCl.

moles of  $Cl^- = ..... mol [1]$ 

(iv) The amount of vanadium chloride in the sample was 0.01 moles.

Use this and your answer to (b)(iii) to deduce the whole number ratio of moles of vanadium chloride: moles of chloride ions.

Deduce the formula of vanadium chloride.

moles of vanadium chloride: moles of chloride ions .....:

formula of vanadium chloride ..... [2]

(c)	(c) Astatine is at the bottom of Group VII. Use your knowledge of the properties of the halog										
	(i)	predict the physical state of astatine at room temperature and pressure,  [1]									
	(ii)			the reaction between sodiur							
					[2]						
(d)	lodi	ne reacts with chlor	ine. The ch	emical equation is shown.							
			I	$_2$ + $Cl_2 \rightarrow 2ICl$							
	Use	e the bond energies	to answer	the questions.							
			bond	bond energy in kJ/mol							
			1–1	151							
			Cl-Cl	242							
			I–Cl	208							
	(ii)	1 mole of $\mathrm{C}l_2$ . Calculate the total formed.	amount of	f energy given out when th	kJ [1] ne bonds in 2 moles of IC <i>l</i> are						
(	(iii)			and <b>(d)(ii)</b> to calculate the $_2$ + $Cl_2  o 2ICl$	kJ [1] overall energy change for the						
					kJ/mol [1] [Total: 15]						

6	(a)	Anl	nomologous series is a 'family' of organic compounds whose names have the same endin	ıg.
		(i)	Name the homologous series for which the names of the organic compounds end in -el and -oic acid.	ne
			-ene[	[1]
			-oic acid	[1]
		(ii)	State <b>two</b> characteristics of an homologous series.	
			[	[2]
	(b)		pan-1-ol is a member of the homologous series of alcohols. It reacts in the same way a anol with acidified potassium manganate(VII) and with carboxylic acids.	as
		Nar	me the <b>type</b> of compound that is formed when propan-1-ol is heated with	
		acio	dified potassium manganate(VII),	
		etha	anoic acid and a suitable catalyst.	
				[2]

(c) The structure of prop-2-enoic (acrylic) acid is shown.



(i)	What would you see if prop-2-enoic acid were added to	
	aqueous bromine,	
	a solution of sodium carbonate.	
		[2

(ii)	Prop-2-enoic acid can be polymerised to form poly(acrylic acid).	
	Suggest the type of polymerisation that occurs and draw one repeat unit of the polymerisation	er.
	type of polymerisation	
	repeat unit	

[3]

[Total: 11]

## **BLANK PAGE**

## **BLANK PAGE**

Permission to reproduce items where third-party owned material protected by copyright is included has been sought and cleared where possible. Every reasonable effort has been made by the publisher (UCLES) to trace copyright holders, but if any items requiring clearance have unwittingly been included, the publisher will be pleased to make amends at the earliest possible opportunity.

To avoid the issue of disclosure of answer-related information to candidates, all copyright acknowledgements are reproduced online in the Cambridge International Examinations Copyright Acknowledgements Booklet. This is produced for each series of examinations and is freely available to download at www.cie.org.uk after the live examination series.

Cambridge International Examinations is part of the Cambridge Assessment Group. Cambridge Assessment is the brand name of University of Cambridge Local Examinations Syndicate (UCLES), which is itself a department of the University of Cambridge.

The Periodic Table of Elements

	<b> </b>	2 He	helium 4	10	Se	neon 20	18	Ar	argon 40	36	궃	krypton 84	54	Xe	xenon 131	98	Rn	radon			
	<b>=</b>			6	ட	fluorine 19	17	Cl	chlorine 35.5	35	南	bromine 80	53	П	iodine 127	85	At	astatine -			
	5			80	0	oxygen 16	16	ഗ	sulfur 32	34	Se	selenium 79	52	<u>e</u>	tellurium 128	84	Ъ	nolonium	116		/ermorium -
	>								shosphorus 31												=
	2								silicon pt										114	Εl	erovium
									lluminium 27												=
									alt									mercury th	112	ت ت	copernicium —
																		m gold m			
													$\vdash$					platinum 195			Ē
Group																					
				]														iridium 192			Ε
		- I	hydroger 1							26	Fe	iron 56	44	Ru	rutheniun 101	92	Os	osmium 190	108	H	hassium
							1			25	Mn	manganese 55	43	ည	technetium -	75	Re	rhenium 186	107	Bh	bohrium –
					pol	ass				24	ပ်	chromium 52	42	Mo	molybdenum 96	74	≥	tungsten 184	106	Sg	seaborgium -
			Key	atomic number	atomic symbo	name relative atomic mass				23	>	vanadium 51	41	g	niobium 93	73	ā	tantalum 181	105	<u>6</u>	dubnium
					ato	rela				22	ı=	titanium 48	40	Zr	zirconium 91	72	茔	hafnium 178	104	፟ጟ	rutherfordium -
							•			21	Sc	scandium 45	39	>	yttrium 89	57-71	lanthanoids		89–103	actinoids	
	=			4	Be	beryllium 9	12	Mg	magnesium 24	20	Ca	calcium 40	38	ട്	strontium 88	56	Ва	barium 137	88	Ra	radium
	_			8	<u> </u>	lithium 7	1	Na	sodium 23	19	¥	potassium 39	37	Rb	rubidium 85	55	S	caesium 133	87	ъ.	francium —

		_			
r Lu	lutetium 175	103	۲	lawrencium	ı
°	ytterbium 173	102	%	nobelium	ı
ee Tm	thulium 169	101	Md	mendelevium	ı
88 Fr	erbium 167	100	Fm	ferminm	I
67 H0	holmium 165	66	Es	einsteinium	ı
° ^	dysprosium 163	86	ŭ	californium	1
65 <b>Tb</b>	terbium 159	97	Ř	perkelium	ı
<sup>2</sup> Q	gadolinium 157	96	Cm	curium	ı
в Еп	europium 152	92	Am	americium	ı
Sm	samarium 150	94	Pu	plutonium	ı
Pm	promethium —	93	dN	neptunium	ı
。 P N	neodymium 144	92	$\supset$	uranium	238
59 Pr	praseodymium 141	91	Ра	protactinium	231
C 28	cerium 140	06	H	thorium	232
57 <b>La</b>	lanthanum 139	88	Ac	actinium	ı

lanthanoids

actinoids

The volume of one mole of any gas is 24 dm³ at room temperature and pressure (r.t.p.).