



# Mark Scheme (Results)

January 2023

Pearson Edexcel International GCSE in  
Chemistry (4CH1)  
Paper 1CR

## **Edexcel and BTEC Qualifications**

Edexcel and BTEC qualifications are awarded by Pearson, the UK's largest awarding body. We provide a wide range of qualifications including academic, vocational, occupational and specific programmes for employers. For further information visit our qualifications websites at [www.edexcel.com](http://www.edexcel.com) or [www.btec.co.uk](http://www.btec.co.uk). Alternatively, you can get in touch with us using the details on our contact us page at [www.edexcel.com/contactus](http://www.edexcel.com/contactus).

## **Pearson: helping people progress, everywhere**

Pearson aspires to be the world's leading learning company. Our aim is to help everyone progress in their lives through education. We believe in every kind of learning, for all kinds of people, wherever they are in the world. We've been involved in education for over 150 years, and by working across 70 countries, in 100 languages, we have built an international reputation for our commitment to high standards and raising achievement through innovation in education. Find out more about how we can help you and your students at: [www.pearson.com/uk](http://www.pearson.com/uk)

January 2023

Question Paper Log Number P71893A

Publications Code 4CH1\_1CR\_MS\_2301

All the material in this publication is copyright



© Pearson Education Ltd 2023

## General Marking Guidance

- All candidates must receive the same treatment. Examiners must mark the first candidate in exactly the same way as they mark the last.
- Mark schemes should be applied positively. Candidates must be rewarded for what they have shown they can do rather than penalised for omissions.
- Examiners should mark according to the mark scheme not according to their perception of where the grade boundaries may lie.
- There is no ceiling on achievement. All marks on the mark scheme should be used appropriately.
- All the marks on the mark scheme are designed to be awarded. Examiners should always award full marks if deserved, i.e. if the answer matches the mark scheme. Examiners should also be prepared to award zero marks if the candidate's response is not worthy of credit according to the mark scheme.
- Where some judgement is required, mark schemes will provide the principles by which marks will be awarded and exemplification may be limited.
- When examiners are in doubt regarding the application of the mark scheme to a candidate's response, the team leader must be consulted.
- Crossed out work should be marked UNLESS the candidate has replaced it with an alternative response.

| Question number | Answer  | Notes   | Marks          |
|-----------------|---|---|----------------|
| 1 (a)           | X evaporating<br>Y condensing<br>Z freezing   | ALLOW evaporation<br><br>ALLOW condensation                               | 3              |
| (b)             | M1 solid particles vibrate about a fixed position<br>M2 gas particles move randomly | REJECT do not move<br><br>ALLOW gas particles move rapidly/quickly/freely | 2              |
|                 |   |   | <b>Total 5</b> |

| Question number | Answer   | Notes  | Marks          |
|-----------------|--|--|----------------|
| 2 (a) (i)       | nitrogen   | <b>ALLOW</b> N <sub>2</sub><br><b>IGNORE</b> N   | 1              |
| (ii)            | argon  | <b>ALLOW</b> Ar  | 1              |
| (iii)           | carbon dioxide   | <b>ALLOW</b> CO <sub>2</sub> /H <sub>2</sub> O(g)/water vapour/CH <sub>4</sub> /methane  | 1              |
| (b) (i)         | brown/red-brown/orange-brown   | <b>ALLOW</b> orange<br><b>IGNORE</b> red<br><br><b>ALLOW</b> rusty/rust coloured (looks like)rust/rusted   | 1              |
| (ii)            | <b>M1</b> (change in length of column =) 84 – 69 OR 15 (mm)<br><br><b>M2</b> $\frac{15 \times 100}{84} = 17.86/17.9 (=18)$ | <b>M2</b> subsumes <b>M1</b><br><br>Working must be shown to score <b>M2</b><br><b>Ecf for M2 eg 18/84 x100 =21.4</b><br><b>REJECT</b> 17.85/17.8 as wrongly rounded | 2              |
| (iii)           | not all the oxygen in the sample of air had reacted with the iron wool OWTTE /not enough iron wool                         | <b>ALLOW</b> there is water vapour in the column of air/changes in temperature / pressure / location<br><b>ALLOW</b> Reaction incomplete/reaction too slow           | 1              |
|                 |  |  | <b>Total 7</b> |

| Question number    | Answer  | Notes   | Marks   |      |        |                   |                                |                   |                               |                 |                                  |  |   |
|--------------------|---|---|---|------|--------|-------------------|--------------------------------|-------------------|-------------------------------|-----------------|----------------------------------|--|---|
| 3 (a)              | <table><tr><td>structural formula</td><td>CH<sub>3</sub>CH<sub>2</sub>CH<sub>2</sub>CH<sub>3</sub></td></tr><tr><td>name</td><td>butane</td></tr><tr><td>molecular formula</td><td>C<sub>4</sub>H<sub>10</sub></td></tr><tr><td>empirical formula</td><td>C<sub>2</sub>H<sub>5</sub></td></tr><tr><td>general formula</td><td>C<sub>n</sub>H<sub>2n+2</sub></td></tr></table> <p>1 mark for each correct answer</p> | structural formula  | CH <sub>3</sub> CH <sub>2</sub> CH <sub>2</sub> CH <sub>3</sub> | name | butane | molecular formula | C <sub>4</sub> H <sub>10</sub> | empirical formula | C <sub>2</sub> H <sub>5</sub> | general formula | C <sub>n</sub> H <sub>2n+2</sub> |  | 4 |
| structural formula | CH <sub>3</sub> CH <sub>2</sub> CH <sub>2</sub> CH <sub>3</sub>   |   |   |      |        |                   |                                |                   |                               |                 |                                  |  |   |
| name               | butane  |   |   |      |        |                   |                                |                   |                               |                 |                                  |  |   |
| molecular formula  | C <sub>4</sub> H <sub>10</sub>  |   |   |      |        |                   |                                |                   |                               |                 |                                  |  |   |
| empirical formula  | C <sub>2</sub> H <sub>5</sub>   |   |   |      |        |                   |                                |                   |                               |                 |                                  |  |   |
| general formula    | C <sub>n</sub> H <sub>2n+2</sub>  |   |   |      |        |                   |                                |                   |                               |                 |                                  |  |   |
| (b) (i)            | <b>M1</b> (compounds with the) same molecular formula<br><br><b>M2</b> (but with) different structural/displayed formulae   | <b>ALLOW</b> same numbers of each atom<br><b>ALLOW</b> different arrangement of atoms | 2   |      |        |                   |                                |                   |                               |                 |                                  |  |   |
| (ii)               | <b>M1</b> displayed formula of butane<br><br><b>M2</b> displayed formula of methylpropane<br>  |   | 2   |      |        |                   |                                |                   |                               |                 |                                  |  |   |
| (c) (i)            | HBr   | REJECT incorrect case letters<br><br>Ignore name                                      | 1   |      |        |                   |                                |                   |                               |                 |                                  |  |   |
| (ii)               | <b>D</b> substitution<br><br>A is incorrect as it is not an addition reaction<br>B is incorrect as it is not a decomposition reaction<br>C is incorrect as it is not a neutralisation reaction  |   | 1   |      |        |                   |                                |                   |                               |                 |                                  |  |   |
| (iii)              | ultraviolet (radiation)   | <b>ACCEPT</b> UV (radiation)<br><br><b>ALLOW</b> ultraviolet/UV light/sunlight        | 1   |      |        |                   |                                |                   |                               |                 |                                  |  |   |
| (d) (i)            | 2C <sub>2</sub> H <sub>6</sub> + 7O <sub>2</sub> → 4CO <sub>2</sub> + 6H <sub>2</sub> O<br><b>M1</b> all formulae correct<br><b>M2</b> balancing of correct formulae  | <b>ALLOW</b> multiples and fractions<br><br><b>M2</b> dep on <b>M1</b>                | 2   |      |        |                   |                                |                   |                               |                 |                                  |  |   |
| (ii)               | An explanation that links the following points<br><br><b>M1</b> carbon monoxide/CO (is the gas produced)<br><br><b>M2</b> (carbon monoxide) limits the capacity of the blood/haemoglobin to carry oxygen OWTTE  |   | 2   |      |        |                   |                                |                   |                               |                 |                                  |  |   |
|                    |   |   | <b>Total 15</b>   |      |        |                   |                                |                   |                               |                 |                                  |  |   |

| Question number | Answer   | Notes  | Marks           |
|-----------------|--|--|-----------------|
| 4 (a)           | Any two from<br><b>M1</b> concentration of solution A<br><br><b>M2</b> concentration of solution B<br><br><b>M3</b> volume of solution B   | ALLOW amount of solution B<br>Ignore apparatus   | 2               |
| (b) (i)         | all points plotted correctly to the nearest grid line  |  | 1               |
| (ii)            | anomalous point at 25°C circled  | <b>ALLOW</b> ecf from incorrect plotting   | 1               |
| (iii)           | smooth curve of best fit ignoring the anomalous point  | ALLOW Ecf if 35,130 circled  | 1               |
| (iv)            | Any one from<br><br><b>M1</b> temperature was higher than 25°C<br><br><b>M2</b> started the timer too late /stopped the timer too early/took reading too early   | <b>ALLOW</b> ecf from incorrect anomalous result circled so 35,130 gives slower as temp<35/timer stopped too late  | 1               |
| (v)             | <b>M1</b> vertical line on graph drawn to curve from 55°C<br><br><b>M2</b> value obtained from candidate's graph   | ALLOW extra point at 55°C on curve<br><br>expected value 115 to 117 s  | 2               |
| (c)             | <b>M1</b> $\frac{1}{156}$ OR 0.00641<br><br><b>M2</b> $6.41 \times 10^{-3}$  | <b>ALLOW</b> use of value from graph<br><br>ALLOW $6.4 \times 10^{-3}$   | 2               |
| (d)             | An explanation that links the following three points<br><br><b>M1</b> rate (of reaction) increases<br><br><b>M2</b> (mean) kinetic energy of particles increases<br><br><br><br><b>M3</b> more successful collisions per second/unit time/ more frequent successful collisions | <b>ALLOW</b> reaction is faster/ speeds up<br><b>ALLOW</b> particles move faster<br><br><b>IGNORE</b> vibrate more /faster<br><br><b>ALLOW</b> more frequent collisions having energy $\geq$ activation energy | 3               |
|                 |  |  | <b>Total 13</b> |





| Question number | Answer   | Notes   | Marks                    |
|-----------------|--|---|--------------------------|
| 6 (a) (i)       | <b>B</b> bromine<br><br>A is incorrect as astatine is a solid at room temperature<br>C is incorrect as chlorine is a gas at room temperature<br>D is incorrect as fluorine is a gas at room temperature  |   | 1                        |
| (ii)            | <b>C</b> dark grey<br><br>A is incorrect as solid iodine is not black<br>B is incorrect as solid iodine is not dark brown<br>D is incorrect as solid iodine is not purple  |   | 1                        |
| (iii)           | <b>M1</b> test with (damp blue) litmus paper<br><br><b>M2</b> bleaches/turns white   | ALLOW Universal indicator paper/ pH paper<br><br><b>ACCEPT</b> turns red and then bleaches  | 2                        |
| (b)             | <b>M1</b> $71.2 \times 35 + 28.8 \times 37$ <b>OR</b> 3557.6<br><br><b>M2</b> $\frac{71.2 \times 35 + 28.8 \times 37}{100}$ <b>OR</b> $\frac{3557.6}{100}$ <b>OR</b> 35.576<br><br><b>M3</b> 35.6  | Correct answer without working scores 3<br><br><b>M2</b> subsumes <b>M1</b><br><br>35.5 without working scores 0  | 3                        |
| (c)             | An explanation that links the following four points<br><br><b>M1</b> add chlorine (solution) to sodium iodide (solution)<br><b>M2</b> solution turns brown<br><br><b>M3</b> iodine/I <sub>2</sub> is displaced<br><br><b>M4</b> (so) chlorine is more reactive (than iodine) ORA | <b>ALLOW</b> mix the two solutions<br><br><b>ALLOW</b> iodine/I <sub>2</sub> is formed<br><br><b>REJECT</b> incorrect use of iodide or chloride once only | 4<br><br><b>Total 11</b> |

| Question number | Answer   | Notes   | Marks           |
|-----------------|--|---|-----------------|
| 7 (a)           | <p><b>M1</b> (bright) white flame/light</p> <p><b>M2</b> white powder/solid (formed)</p>   | <p><b>ALLOW</b> white smoke/ash</p> <p><b>ALLOW</b> grey powder</p> <p><b>REJECT</b> white precipitate</p>  | 2               |
| (b) (i)         | gives out/releases heat (energy)/thermal energy  | <b>IGNORE</b> energy alone  | 1               |
| (ii)            | $2\text{Al} + \text{Fe}_2\text{O}_3 \rightarrow 2\text{Fe} + \text{Al}_2\text{O}_3$  | <b>ALLOW</b> multiples and fractions  | 1               |
| (iii)           | <p>An explanation that links two of the following pairs of points</p> <p><b>M1</b> aluminium/Al gains oxygen so is oxidised</p> <p><b>M2</b> iron oxide/<math>\text{Fe}_2\text{O}_3</math> loses oxygen so is reduced</p> <p><b>OR</b></p> <p><b>M1</b> aluminium/Al is oxidised and iron oxide/<math>\text{Fe}_2\text{O}_3</math> is reduced</p> <p><b>M2</b> as aluminium/Al gains oxygen and iron oxide/<math>\text{Fe}_2\text{O}_3</math> loses oxygen</p> | <p><b>ACCEPT</b> aluminium loses electrons so is oxidised</p> <p><b>ACCEPT</b> iron ions/<math>\text{Fe}^{3+}</math> ions gain electrons so are reduced</p> <p><b>ACCEPT</b> aluminium loses electrons and iron ions/<math>\text{Fe}^{3+}</math> ions gain electrons</p> <p><b>ALLOW</b> answers in terms of change in oxidation number</p> | 2               |
| (c) (i)         | <p>An explanation that links the following two points</p> <p><b>M1</b> to allow air/oxygen to enter the crucible OWTTE</p> <p><b>M2</b> so that oxygen can react with the magnesium</p>  |   | 2               |
| (ii)            | <p>A description that refers to the following points</p> <p><b>M1</b> heat the crucible again and reweigh</p> <p><b>M2</b> repeat until constant mass</p>  | Heat and reweigh to constant mass scores 2  | 2               |
|                 |  |   | <b>Total 10</b> |

| Question number | Answer  | Notes  | Marks           |
|-----------------|---|--|-----------------|
| 8 (a) (i)       | (thermal) decomposition   |  | 1               |
| (ii)            | <b>M1</b> amount of $\text{PbCO}_3 = \frac{5.34}{267} = 0.02(00)$ (mol)<br><br><b>M2</b> mass of $\text{PbO} = 0.02(00) \times 223 = 4.46$ (g)  | Correct answer without working scores 2<br><br><b>ACCEPT</b> alternative methods   | 2               |
| (b) (i)         | <b>M1</b> diagram showing delivery tube going into test tube containing liquid  | REJECT if sealed with a bung   | 2               |
| (ii)            | <b>M2</b> limewater labelled (limewater) turns cloudy/milky   | <b>ALLOW</b> white precipitate<br><br>(ii) dep on mention of limewater in either (i) or (ii)   | 1               |
| (c)             | An explanation that links six of the following points<br><br><b>M1</b> silicon dioxide has a giant (covalent) structure<br><br><b>M2</b> covalent bonds are (very) strong<br><br><b>M3</b> (in silicon dioxide) many covalent bonds need to be broken<br><br><b>M4</b> a large amount of energy/more energy is required to break the bonds in silicon dioxide<br><br><b>M5</b> carbon dioxide has a simple molecular structure/is a simple molecule<br><br><b>M6</b> the forces between the molecules/intermolecular forces (in carbon dioxide) are weak<br><br><b>M7</b> very little energy/less energy is needed to overcome the forces between the molecules/intermolecular forces (in carbon dioxide) | No <b>M3</b> or <b>M4</b> if reference to intermolecular forces in silicon dioxide<br><br>No <b>M6</b> or <b>M7</b> if any reference to weak covalent bonds or breaking of covalent bonds in carbon dioxide<br>Accept bonds between molecules weak<br><br>A statement such as 'more energy is needed to break the bonds in silicon dioxide than to overcome the forces between the molecules/intermolecular forces (in carbon dioxide)' scores <b>M4</b> and <b>M7</b> | 6               |
|                 |   |  | <b>Total 12</b> |



| Question number                              | Answer  | Notes   | Marks |                                       |      |                          |     |   |   |
|--|---|---|-------|---------------------------------------|------|--------------------------|-----|---|---|
| 10 (a)                                       | <p><b>M1</b> so that the solid/ammonium nitrate dissolves more quickly</p> <p><b>M2</b> so that the temperature is even throughout the solution OWTTE</p>   | <p>IGNORE speed up reaction</p> <p>ALLOW heat transfers evenly (throughout the solution)</p>  | 2     |                                       |      |                          |     |   |   |
| (b)  | <table><tr><td>initial temperature of distilled water in °C</td><td>23.4</td></tr><tr><td>minimum temperature of solution in °C</td><td>19.4</td></tr><tr><td>temperature change in °C</td><td>4.0</td></tr></table>  | initial temperature of distilled water in °C  | 23.4  | minimum temperature of solution in °C | 19.4 | temperature change in °C | 4.0 | <p>must be to 1 dp</p> <p><b>ALLOW</b> ecf on incorrect minimum temperature</p> | 2 |
| initial temperature of distilled water in °C | 23.4  |   |       |                                       |      |                          |     |   |   |
| minimum temperature of solution in °C        | 19.4  |   |       |                                       |      |                          |     |   |   |
| temperature change in °C                     | 4.0   |   |       |                                       |      |                          |     |   |   |
| (c) (i)                                      | <p><b>M1</b> (<math>Q =</math>) <math>50 \times 4.2 \times 3.9</math> (J)</p> <p><b>M2</b> (<math>Q =</math>) 819/820 (J)</p>   | <p>answer of 819 or 820 without working scores 2</p> <p><b>ALLOW</b> use of 4.0 giving an answer of 840</p>   | 2     |                                       |      |                          |     |   |   |
| (ii)   | <ul style="list-style-type: none"><li>find moles of <math>\text{NH}_4\text{NO}_3</math></li><li>division of <math>Q</math> by moles</li><li>conversion to kJ/mol</li><li>answer with correct sign</li></ul> <p><b>M1</b> (amount of <math>\text{NH}_4\text{NO}_3 =</math>) <math>2.8 \div 80</math> <b>OR</b> 0.035 (mol)</p> <p><b>M2</b> <math>819 \div 0.035</math> <b>OR</b> 23 400 (J/mol)</p> <p><b>M3</b> <math>23\,400 \div 1000</math> <b>OR</b> 23.4 (kJ/mol)</p> <p><b>M4</b> (<math>\Delta H =</math>) +23.4/+23 (kJ/mol)</p> | <p>correct answer without working scores 4</p> <p>use of 820 gives 23 429<br/>use of 800 gives 22 857<br/>use of 840 gives 24 000<br/>use of 820 gives 23.4<br/>use of 800 gives 22.9<br/>use of 840 gives 24.0</p> | 4     |                                       |      |                          |     |   |   |
| (d)  | <p>A description that refers to the following points</p> <p><b>M1</b> add sodium hydroxide (solution to the ammonium nitrate and warm)</p> <p><b>M2</b> test the gas/ammonia evolved with (damp) red litmus paper/(damp) universal indicator paper</p> <p><b>M3</b> (red litmus) turns blue/ (universal indicator) turns blue/purple</p>  | <p><b>M2</b> and <b>M3</b> dep on <b>M1</b></p> <p>No <b>M2</b> or <b>M3</b> if solution tested with litmus/ universal indicator paper</p>  | 3     |                                       |      |                          |     |   |   |

|     |  |  |                 |
|-----|--|--|-----------------|
| (e) | An explanation that links the following points<br><b>M1</b> the temperature increases/rises<br><b>M2</b> so the reaction is exothermic |  | 2               |
|     |  |  | <b>Total 15</b> |

