1 Choose a gas from the following list to answer the questions below. Each gas may be used once, more than once or not at all.


## ammonia carbon dioxide carbon monoxide fluorine hydrogen krypton nitroge sulfur dioxide propene (a) It is a product of respiration. [1] (e) It is a very reactive non-metal. [1] (f) It is used to kill micro-organisms in fruit juice. [1] [Total: 7]

Antimo	ny, Sb, is an element in Group V.				
zin					
De	escribe how each of these changes in the extraction of antimony is carried out.				
(i)	(i) antimony sulfide to antimony oxide				
	[1]				
(ii)	antimony oxide to antimony				
	[1]				
	timony oxide is a white powder which is insoluble in water. scribe how you would find out if it is a basic, an acidic or an amphoteric oxide.				
	[4]				
(a) \/\/h					
	en antimony chloride is added to water, a faint white precipitate forms and the mixture wly goes cloudy.				
	forward				
	$SbCl_3(aq) + H_2O(I) \rightleftharpoons 2HCl(aq) + SbOCl(s)$				
	colourless backward white				
(i)	Explain why after some time the appearance of the mixture remains unchanged.				
	[2]				
(ii)	When a few drops of concentrated hydrochloric acid are added to the mixture, it changes to a colourless solution. Suggest an explanation.				
	[1]				
(iii)	Suggest how you could make the colourless solution go cloudy.				
	[1]				
	[Total: 10]				

2

3	The hydrolysis of complex carbohydrates to simple sugars is catalysed by enzymes carbohydrases and also by dilute acids.		
	(a) (	(i)	They are both catalysts. How do enzymes differ from catalysts such as dilute acids?
			[1]
	<b>(</b> i	ii)	Explain why ethanol, $C_2H_6O$ , is not a carbohydrate but glucose, $C_6H_{12}O_6$ , is a carbohydrate.
			[2]
			w the structure of a complex carbohydrate, such as starch. The formula of a simple ar can be represented by HO — OH.

- (c) Iodine reacts with starch to form a deep blue colour.
  - (i) In the experiment illustrated below, samples are removed at intervals and tested with iodine in potassium iodide solution.



Typical results of this experiment are shown in the table.

time/min	colour of sample tested with iodine in potassium iodide solution
0	deep blue
10	pale blue
30	colourless

	Explain these results.
	[3
(ii)	If the experiment was repeated at a higher temperature, 60 °C, all the samples stayed blue. Suggest an explanation.
	[1
	[Total: 10

4 Complete the following table.

gas	test for gas
ammonia	
	bleaches damp litmus paper
hydrogen	
	relights a glowing splint
	turns limewater milky

[Total: 5]

5	Choose a gas from the following list to answer the questions below. Each gas may be used once, more than once or not at all.					
		ammonia	argon		carbon monoxide	chlorine
		ethene	hydrogen	nitrogen	oxygen	
		Which gas				
	(i)	is a noble g	jas,			
	(ii)	is an acidic	oxide			
	(,	io dir dolaio	OXIGO,			
	(iii)	can be poly	merised,			
	(:)	io the eather		af air		
	(iv)	is the active	e component	ot air,		
	(v)	is used in t	he treatment	of water,		
	(vi)	is a produc	t of respiration	n?		

[1]	
(ii) Explain what is meant by the term fossil fuel.	
[2	2]
(b) The burning of fossil fuels is largely responsible for the formation of acid rain.  Two of the acids in acid rain are sulfuric acid and nitric acid.	
(i) Explain how the combustion of coal can form sulfuric acid.	
[3	3]

(a) (i) Coal is a solid fossil fuel. Name another fossil fuel.

6

(II)	of nitric acid. Explain.
	[3]
ii)	Nitric acid contains nitrate ions.
	Describe a test for nitrate ions.
	[2]
<b>v</b> )	Explain how you could determine which one of two samples of acid rain had the higher concentration of hydrogen ions.
	[2]

[Total: 13]

		chlorine bromine iodine	
(a)		How does their colour change down the Group?	
			[1]
	(ii)	How does their physical state (solid, liquid or gas) change down the Group?	41
			[1]
	(iii)	Predict the colour and physical state of fluorine.	
		colour	•••
		physical state [2	[2]
(b)		scribe how you could distinguish between aqueous potassium bromide and aqueouassium iodide.	us
	test	t	
	res	ult with bromide	
	res	ult with iodide [	3]
(c)		15 moles of iodine react with 0.045 moles of chlorine to form 0.030 moles of a sing duct. Complete the equation.	ıle
	I2	Cl₂ →	[2]
(d)		ices of chlorine can be separated from bromine vapour by diffusion. ich gas would diffuse the faster and why?	
			•••
		[	[2]

Three of the halogens in Group VII are: