

| Question | Answer  | Marks | Guidance   |
|----------|---|-------|--|
| 1(a)     | ( $\text{CuCO}_3 \rightarrow \text{CuO} + \text{CO}_2$ ;<br>$(\text{Cu}(\text{OH})_2 \rightarrow \text{CuO} + \text{H}_2\text{O}$ ;<br>$(2\text{Cu}(\text{NO}_3)_2 \rightarrow 2\text{CuO} + (4\text{NO}_2) + \text{O}_2$<br>species;<br>balancing; | 4     | A multiples<br>I state symbols   |
| (b)(i)   | (black to) pink/brown/orange;   | 1     | red  |
| (b)(ii)  | (hot) copper reacts/is oxidised;<br>with oxygen/air;  | 2     | A forms copper oxide for 2 marks   |
| (iii)    | monoxide/ammonia/methane;   | 1     |  |
| (b)(iv)  | /graphite or any metal more reactive than copper;   | 1     |  |
| 1(c)(i)  | 79.28<br>79.6205853;<br>84.7161572;   | 2     | Minimum 3 sig figs<br>A rounding or truncating<br>All three correct = 2 marks,<br>Two correct = 1 mark |
| (c)(ii)  | the last one <b>OR</b> Cu and O <sub>2</sub> <b>OR</b> the one from copper;<br><br>not all the copper oxidised <b>OR</b> the outside of the pieces of copper oxidised<br>but the inside did not <b>OR</b> (still) contains copper (metal);          | 2     | ecf of biggest for M1  |

2 (a) (i)  $82.76/12$  and  $17.2(4)/(1)$  [1]  
or evaluation:  $6.89 / 6.9(0)$  and  $17.2(4)$

$C_2H_5$  [1]

**OR**

$82.76/100 \times 58 = 48$  and  $17.24/100 \times 58 = 10$  [1]  
or evaluation i.e. 48 and 10

$C_2H_5$  [1]

(ii)  $(C_2H_5 =) 29$  [1]

$(58/29 = 2)$   $C_4H_{10}$  [1]

**OR:**

$82.76/100 \times 58 = 48$  and  $17.24/100 \times 58 = 10$  [1]  
or evaluation i.e. 48 and 10

$48/12 = 4$   $10/1 = 10$  (therefore)  $C_4H_{10}$  [1]

(b) (i)  $C_nH_{2n}$  [1]

(ii)  $CH_2$  [1]

(c) (contains) double bond/triple bond/multiple bond(s)/not all bonds are single [1]

(contains) carbon and hydrogen **only** [1]


(d) bromine/bromine water [1]

no change/stays brown/orange/yellow/red-brown or only changes in UV [1]

(brown/orange/yellow) to colourless/decolourised

(e) (i) circle/brackets around any 2 consecutive carbon atoms in the main chain [1]  
and all attached atoms

e.



(ii)  $CH_3CH_2CH=CH_2$  /  $C_2H_5CH=CH_2$  (double bond must be shown) [1]

butene / but-1-ene

(iii)  $(\text{CH}_3)_2\text{C}=\text{CH}_2 / \text{CH}_3\text{CH}=\text{CHCH}_3 / (\text{CH}_2)_2\text{CHCH}_3 / (\text{CH}_2)_4$  [1]

[Total:15]

3 (a) Any **two** from:  
yeast / 20–40 °C / anaerobic or without oxygen or without air / (aqueous)  
solution or water or aqueous [2]

(b) (i)  $M_r = 180$  (1)  $(30/180) = 0.167$  (1) [2]

(ii)  $2 \times 0.167$  or  $2 \times 46$  or  $0.333$  or  $92$  [1]

$(2 \times 0.167 \times 46) = 15.3(33)$  (g) [1]

(iii)  $(2 \times 0.167 \times 24) = 8$  (dm<sup>3</sup>) [1]

(c) (i) Crude oil / petroleum

(ii)  $\text{C}_2\text{H}_4 + \text{H}_2\text{O} \rightarrow \text{C}_2\text{H}_5\text{OH} / \text{CH}_3\text{CH}_2\text{OH}$  [

[Total:9]

4 (a) (i) (the number of particles which is equal to the number of atoms in) 12 g of carbon 12  
**or**  
the mass in grams which contains the Avogadro's constant number of particles  
**or**  
Avogadro's constant **or** 6 to  $6.023 \times 10^{23}$  of atoms / ions / molecules / electrons / particles  
**or**  
(the amount of substance which has a mass equal to) its relative formula mass / relative atomic mass / relative molecular mass in grams  
**or**  
(the amount of substance which has a volume equal to) 24 dm<sup>3</sup> of a gas at RTP [1]

(ii) (Avogadro's constant is the) number of particles / atoms / ions / molecules in one mole of a substance  
**or**  
the number of carbon atoms in 12 g of C(12).  
**or**  
the number of particles / molecules in 24 dm<sup>3</sup> of a gas at RTP  
**or**  
 $6$  to  $6.023 \times 10^{23}$  (particles / atoms / ions / molecules / electrons) [1]

(b)  $\text{CH}_4$  and  $\text{SO}_2$  [1]

$2/16 = 1/8$  or 0.125 moles of  $\text{CH}_4$  **AND**  $8/64 = 1/8$  or 0.125 moles of  $\text{SO}_2$  [1]

- (c) (i)  $4.8/40 = 0.12$  moles of Ca  
 $3.6/18 = 0.2$  moles of  $H_2O$  **both** correct [1]
- (ii) Ca is in excess (**no mark**) (because 0.12 moles of Ca need) 0.24 moles / 4.32g of  $H_2O$  to react [1]  
 there is not enough / there are 0.2 moles / 3.6g of  $H_2O$  [1]  
**or**  
 Ca is in excess (**no mark**) (because 0.2 moles / 3.6g of water will react with) 0.1moles/4.0g of Ca [1]  
 there is more than that / there are 0.12 moles / 4.8g of Ca [1]  
**or**  
 Ca is in excess (**no mark**) because the mole ratio Ca: $H_2O$  is 3:5 / mass ratio 4:3 [1]  
 which is bigger than the required mole ratio of 1:2 / mass ratio 10:9 [1]  
**or**  
 Ca is in excess (**no mark**) because the mole ratio  $H_2O$ :Ca is 5:3 / mass ratio 3:4 [1]  
 which is smaller than the required mole ratio of 2:1 / mass ratio 9:10 [1]
- (iii)  $0.02 \times 40 = 0.8$  (g) [1]

- 5 (a)  $72/24 = 3$  and  $28/14 = 2$  [1]  
 $Mg_3N_2$  [1]  
**accept** just formula for [2] even with incorrect or no working  
**NOT** ecf

- (b)  $Al_4C_3 + 12H_2O = 4Al(OH)_3 + 3CH_4$  [2]  
 For  $Al_4C_3$  ONLY [1]

- (c) silicon is limiting reagent [1]  
 0.07 moles of Si and  $25/160 = 0.156$  moles of  $Br_2$  [1]  
 because  $0.14 (2 \times 0.07) < 0.156$  [1]  
 If 80 used to find moles of  $Br_2$  the mark 1 and 3 still available  
 arguments based on masses can be used

- (ii) 0.07 [1]  
**NOT** ecf

[Total: 8]

### Question 6

- (a)(i) moles of  $\text{NiCO}_3$  reacted = 0.08 [1]  
mass of nickel carbonate reacted = 9.52 g [1]  
mass of nickel carbonate unreacted = 2.48 g [1]
- (ii) maximum number of moles of hydrated salt = 0.08 [1]  
maximum mass of salt =  $0.08 \times 281 = 22.48$  g [1]  
percentage yield  $10.4/22.48 \times 100 = 46.3\%$  [1]
- (b)(i) sulphuric acid  
**COND** description of titration  
repeat without indicator **or** with carbon  
evaporation  
any **TWO** [3]
- (ii) suitable reactants calcium chloride and sodium fluoride [1]  
**COND** upon correct reagents  
filter [1]  
wash and dry precipitate [1]
- OR** Accept synthesis  
calcium [1]  
fluorine [1]  
burn **or** heat [1]

**TOTAL = 12**