Milankovitch Cycles and Glaciation

The episodic nature of the Earth's glacial and interglacial periods within the present Ice Age (the last couple of million years) have been caused primarily by cyclical changes in the Earth's circumnavigation of the Sun. Variations in the Earth's eccentricity, axial tilt, and precession comprise the three dominant cycles, collectively known as the Milankovitch Cycles for Milutin Milankovitch, the Serbian astronomer who is generally credited with calculating their magnitude. Taken in unison, variations in these three cycles creates alterations in the seasonality of solar radiation reaching the Earth's surface. These times of increased or decreased solar radiation directly influence the Earth's climate system, thus impacting the advance and retreat of Earth's glaciers.

It is of primary importance to explain that climate change, and subsequent periods of glaciation, resulting from the following three variables is not due to the total amount of solar energy reaching Earth. The three Milankovitch Cycles impact the **seasonality and location** of solar energy around the Earth, thus impacting contrasts between the seasons.

The first of the three Milankovitch Cycles is the Earth's *eccentricity*. Eccentricity is, simply, the shape of the Earth's orbit around the Sun. This constantly fluctuating, orbital shape ranges between more and less elliptical (0 to 5% ellipticity) on a cycle of about 100,000 years. These oscillations, from more elliptic to less elliptic, are of prime importance to glaciation in that it alters the distance from the Earth to the Sun, thus changing the distance the Sun's short wave radiation must travel to reach Earth, subsequently reducing or increasing the amount of radiation received at the Earth's surface in different seasons.

LESS ELLIPTICAL

Eccentricity

ECCENTRICITY

MORE ELLIPTICAL



ORBIT

PERIODICITY:

100,000 YEARS

Today a difference of only about 3 percent occurs between aphelion (farthest point) and perihelion (closest point). This 3 percent difference in distance means that Earth experiences a 6 percent increase in received solar energy in January than in July. This 6 percent range of variability is not always the case, however. When the Earth's orbit is most elliptical the amount of solar energy received at the perihelion would be in the range of 20 to 30 percent more than at aphelion. Most certainly these continually altering amounts of received solar energy



Axial Tilt

Axial tilt, the second of the three Milankovitch Cycles, is the inclination of the Earth's axis in relation to its plane of orbit around the Sun. Oscillations in the degree of Earth's axial tilt occur on a periodicity of 41,000 years from 21.5 to 24.5 degrees.



PERIODICITY:

41,000 YEARS

Today the Earth's axial tilt is about 23.5 degrees, which largely accounts for our seasons. Because of the periodic variations of this angle the severity of the Earth's seasons changes. With less axial tilt the Sun's solar radiation is more evenly distributed between winter and summer. However, less tilt also increases the difference in radiation receipts between the equatorial and polar regions.

One hypothesis for Earth's reaction to a smaller degree of axial tilt is that it would promote the growth of ice sheets. This response would be due to a warmer winter, in which warmer air would be able to hold more moisture, and subsequently produce a greater amount of snowfall. In addition, summer temperatures would be cooler, resulting in less melting of the winter's accumulation. At present, axial tilt is in the middle of its range.



The third and final of the Milankovitch Cycles is Earth's precession. Precession is the Earth's slow wobble as it spins on axis. This wobbling of the Earth on its axis can be likened to a top running down, and beginning to wobble back and forth on its axis. The precession of Earth wobbles from pointing at Polaris (North Star) to pointing at the star Vega. When this shift to the axis pointing at Vega occurs, Vega would then be considered the North Star. This top-like wobble, or precession, has a periodicity of 23,000 years.

Milankovitch Cycles



Due to this wobble a climatically significant alteration must take place. When the axis is tilted towards Vega the positions of the Northern Hemisphere winter and summer solstices will coincide with the aphelion and perihelion, respectively. This means that the Northern Hemisphere will experience winter when the Earth is furthest from the Sun and summer when the Earth is closest to the Sun. This coincidence will result in greater seasonal contrasts. At present, the Earth is at perihelion very close to the winter solstice.

Summary

These variables are only important because the Earth has an asymmetric distribution of landmasses, with virtually all (except Antarctica) located in the Northern Hemisphere.









http://www.homepage.montana.edu/~geol445/hyperglac/time1/milankov.htm (3 of 4)10/11/2006 11:49:13 AM



(From NOAA's National Geophysical Data Center) At times when Northern Hemisphere summers are coolest (farthest from the Sun due to precession and greatest orbital eccentricity) and winters are warmest (minimum tilt), snow can accumulate on and cover broad areas of northern America and Europe. At present, only precession is in the glacial mode, with tilt and eccentricity not favorable to glaciation

Even when all of the orbital parameters favor glaciation, the increase in winter snowfall and decrease in summer melt would barely enough to trigger glaciation, not to grow large ice sheets. Ice sheet growth requires the support of positive feedback loops, the most obvious of which is that snow and ice have a much lower albedo than ground and vegetation, thus ice masses tend to reflect more radiation back into space, thus cooling the climate and allowing glaciers to expand.



http://www.homepage.montana.edu/~geol445/hyperglac/time1/milankov.htm (4 of 4)10/11/2006 11:49:13 AM