



**Fig. 1 Adapted news article about an El Niño weather event**

### **El Niño: 75% chance of extreme weather phenomenon in next three months**

Experts have warned there is a 75% chance of an El Niño weather system forming within the next three months. This naturally-occurring weather event changes temperatures across the Pacific Ocean and drives extreme weather across the world.

Droughts and flooding are likely to follow as worldwide climate patterns are thrown into disarray for the first part of 2019. The most recent El Niño event ended in 2016, and was associated with catastrophic coral bleaching on the Great Barrier Reef off the Australian coast, as global temperatures reached the highest levels ever recorded. El Niño weather events can affect rainfall and temperature patterns in many regions, with important consequences for the farming industry and food security in many LIDCs.

Chances of dry conditions and drought will increase in nations from the Caribbean to southern Africa, while heavy rainfall will likely hit parts of the US and Europe. Under the most extreme conditions currently predicted, sea temperatures will rise up to 1.2°C above average.

The Met Office has previously warned that El Niño could impact winter weather in the UK next year. "An El Niño could create wetter and windier conditions at the start of winter" a spokesperson said.

Fig. 2 Effect of El Niño weather event on world climate patterns




Fig. 3 River Flow in the UK, December 2018







Fig. 4 Top twenty export destinations from Heathrow Airport, 2008



**Fig. 5 Climate Graph for Isachen, Canada**

**Fig. 6 The impact of climate change on polar ecosystems**




The impact of climate change on a particular species in a food web can go on to affect a wide range of other organisms. For example, the figure above shows the complex nature of the food web for polar bears. Not only is the decline of sea ice affecting polar bear populations by reducing the size of their primary habitat, it is also negatively impacting them by the effect on food webs. A decline in sea ice in the Arctic leads to declines in the abundance of ice algae. These algae are eaten by zooplankton, which are in turn eaten by Arctic cod, an important food source for many marine mammals, including seals. Seals are eaten by polar bears. Hence, declines in ice algae can contribute to declines in polar bear populations.


Fig. 7 Urban regeneration in a UK town centre





**Fig. 8 Results of coastal fieldwork**

Question 1: Since the new coastal defence measures have been built, has damage to your property changed?



Student photo of new groynes



**BLANK PAGE**

**BLANK PAGE**

**Copyright Information**

OCR is committed to seeking permission to reproduce all third-party content that it uses in its assessment materials. OCR has attempted to identify and contact all copyright holders whose work is used in this paper. To avoid the issue of disclosure of answer-related information to candidates, all copyright acknowledgements are reproduced in the OCR Copyright Acknowledgements Booklet. This is produced for each series of examinations and is freely available to download from our public website ([www.ocr.org.uk](http://www.ocr.org.uk)) after the live examination series.

If OCR has unwittingly failed to correctly acknowledge or clear any third-party content in this assessment material, OCR will be happy to correct its mistake at the earliest possible opportunity.

For queries or further information please contact The OCR Copyright Team, The Triangle Building, Shaftesbury Road, Cambridge CB2 8EA.

OCR is part of the Cambridge Assessment Group; Cambridge Assessment is the brand name of University of Cambridge Local Examinations Syndicate (UCLES), which is itself a department of the University of Cambridge.