

C1 January 2007 (MA)

Q1)  $y = 4x^3 - 1 + 2x^{\frac{1}{2}}, x > 0$

$$y = 4x^3 - 1 + 2x^{\frac{1}{2}}$$

$$\frac{dy}{dx} = 12x^2 + x^{-\frac{1}{2}}$$


$$= \boxed{12x^2 + \frac{1}{\sqrt{x}}}$$

Q2a)  $\sqrt{108} = \sqrt{36 \times 3} = \sqrt{36} \sqrt{3} = \boxed{6\sqrt{3}}$

b)  $(2 - \sqrt{3})^2 = (2 - \sqrt{3})(2 - \sqrt{3})$   
 $= 4 - 2\sqrt{3} - 2\sqrt{3} + 3$   
 $= \boxed{7 - 4\sqrt{3}}$

Q3a)  $f(x) = \frac{1}{x}, x \neq 0$

$y = f(x) + 3$  is a transformation of  $f(x)$   
 up 3 y-coordinates.



**Asymptotes at  $y=3$  and  $x=0$**

$$b) \quad \frac{1}{x} + 3 = 0 \quad \leftarrow \text{Since } f(x) = \frac{1}{x}$$

$$\frac{1}{x} = -3$$

$$1 = -3x$$

$$\underline{x = -\frac{1}{3}}$$

$\therefore$  the curve  $y = f(x) + 3$  crosses the x-axis at  $\boxed{\left(-\frac{1}{3}, 0\right)}$

Q4)

$$y = x - 2 \quad \Rightarrow \textcircled{1}$$

$$y^2 + x^2 = 10 \quad \Rightarrow \textcircled{2}$$

Substitute  $\textcircled{1}$  into  $\textcircled{2}$ :

$$(x-2)^2 + x^2 = 10$$

$$(x-2)(x-2) + x^2 = 10$$

$$x^2 - 4x + 4 + x^2 - 10 = 0$$

$$2x^2 - 4x - 6 = 0$$

$$x^2 - 2x - 3 = 0$$

$$(x-3)(x+1) = 0$$

Either  $x=3$  or  $x=-1$

When  $x=3$ , substitute into ① for  $y$ :

$$y = x - 2$$

$$y = 3 - 2$$

$$\underline{y = 1}$$

When  $x = -1$ , substitute into ① for  $y$ :

$$y = x - 2$$

$$y = -1 - 2$$

$$\underline{y = -3}$$

Solution set:

$x = 3, y = 1$	$x = -1, y = -3$
----------------	------------------

Q5)  $2x^2 - 3x - (k+1) = 0$  has no real roots

\* If a quadratic has no real roots, then the discriminant is less than 0

$$\therefore b^2 - 4ac < 0$$

$$(-3)^2 - (4)(2)(-k-1) < 0$$

$$9 - 8(-k-1) < 0$$

$$9 + 8k + 8 < 0$$

$$17 + 8k < 0$$

$$8k < -17$$

$$\boxed{k < -\frac{17}{8}}$$

$$\text{Q6a) } (4 + 3\sqrt{x})^2 = (4 + 3\sqrt{x})(4 + 3\sqrt{x})$$

$$= 16 + 12\sqrt{x} + 12\sqrt{x} + 9x$$

$$= \boxed{16 + 24\sqrt{x} + 9x}$$

$$\therefore \underline{k = 24}$$

$$\text{b) } \int (4 + 3\sqrt{x})^2 = \int (16 + 24\sqrt{x} + 9x) dx$$

$$= \int (16 + 24x^{\frac{1}{2}} + 9x) dx$$

$$= 16x + \frac{24x^{\frac{3}{2}}}{\frac{3}{2}} + \frac{9x^2}{2} + C$$

$$= 16x + 16x^{\frac{3}{2}} + \frac{9x^2}{2} + C$$

$$= \boxed{16x + 16x\sqrt{x} + \frac{9x^2}{2} + C}$$

$$Q7a) f'(x) = 3x^2 - 6 - \frac{8}{x^2}$$

$$f'(x) = 3x^2 - 6 - 8x^{-2}$$

$$f(x) = \int (3x^2 - 6 - 8x^{-2}) dx$$

$$= \frac{3x^3}{3} - 6x - \frac{8x^{-1}}{-1} + C$$

$$= x^3 - 6x + 8x^{-1} + C$$

$$= x^3 - 6x + \frac{8}{x} + C$$

We know that (2,1) lies on  $f(x)$ , so substitute in  $x=2$  and  $y=1$ :

$$1 = (2)^3 - 6(2) + \frac{8}{2} + C$$

$$1 = 8 - 12 + 4 + C$$

$$1 = C$$

$$\therefore \underline{C = 1}$$

$$\text{So, } \boxed{f(x) = x^3 - 6x + \frac{8}{x} + 1}$$

b) To find gradient of tangent to  $f(x)$  at  $(2,1)$ , substitute in  $x=2$  to  $f'(x)$ :

$$f'(x) = 3x^2 - 6 - \frac{8}{x^2}$$

$$\text{When } x=2, f'(x) = 3(2)^2 - 6 - \frac{8}{(2)^2}$$

$$= 3(4) - 6 - \frac{8}{4}$$

$$= 12 - 6 - 2$$

$$= \underline{4}$$

So the gradient,  $m$ , = 4

Equation of tangent :  $y - y_1 = m(x - x_1)$

Using  $(2,1)$   $\rightarrow y - 1 = 4(x - 2)$

$$y - 1 = 4x - 8$$

$$\boxed{y = 4x - 7}$$

$$Q8) \quad y = 4x + 3x^{\frac{3}{2}} - 2x^2, \quad x > 0$$

$$a) \quad \frac{dy}{dx} = 4 + \frac{9}{2}x^{\frac{1}{2}} - 4x$$

$$= \boxed{4 + \frac{9\sqrt{x}}{2} - 4x}$$

b) Substitute in  $x=4, y=8$  into  $y$ :

$$y = 4x + 3x^{\frac{3}{2}} - 2x^2$$

$$8 = 4(4) + 3(4)^{\frac{3}{2}} - 2(4)^2$$

$$8 = 16 + 3(8) - 2(16)$$

$$8 = 16 + 24 - 32$$

$$\underline{8 = 8}$$

$\therefore$  the point  $P(4, 8)$  lies on  $C$ .

$$c) \quad \text{when } x=4, \quad \frac{dy}{dx} = 4 + \frac{9(\sqrt{4})}{2} - 4(4)$$

$$= 4 + 9 - 16$$

$$= \underline{-3}$$

$\therefore$  the gradient of the normal at  $P$  is  $\underline{\frac{1}{3}}$ .

Equation of normal:  $y - y_1 = m(x - x_1)$

Using  $P(4, 8) \rightarrow y - 8 = \frac{1}{3}(x - 4)$

$$3(y - 8) = 1(x - 4)$$

$$3y - 24 = x - 4$$

$$\boxed{3y = x + 20}$$

d) Normal at P cuts x-axis at Q.

When normal cuts x-axis,  $y = 0$

$$\text{So, } 3(0) = x + 20$$

$$0 = x + 20$$

$$x = -20$$

$$\therefore \underline{Q \text{ is at } (-20, 0)}$$

Also, P is at  $(4, 8)$

$$\text{Length } PQ = \sqrt{(x_1 - x_2)^2 + (y_2 - y_1)^2}$$




$$= \sqrt{(4 - (-20))^2 + (8 - 0)^2}$$

$$= \sqrt{24^2 + 8^2}$$

$$= \sqrt{640} = \sqrt{64 \times 10} = \sqrt{64} \sqrt{10}$$

$$= \boxed{8\sqrt{10} \text{ units}}$$



Q9) Row 1		- 4 sticks
Row 2		- 7 sticks
Row 3		- 10 sticks

a) 4 sticks in Row, so first term,  $a = 4$

Common difference,  $d = 3$

$$U_n = a + (n-1)d$$

$$U_n = 4 + (n-1)3$$

$$U_n = 4 + 3n - 3$$

$$\boxed{U_n = 3n + 1}$$

b)  $S_n = \frac{n}{2} (2a + (n-1)d)$

$$S_{10} = \frac{10}{2} (2(4) + (10-1)3)$$

$$= 5(8 + (9)3)$$

$$= 5(8 + 27)$$

$$= 5(35)$$

$$\boxed{= 175}$$

- c) Ann started with 1750 sticks.  
She has enough to complete  $k$  rows,  
but not  $(k+1)$  rows.

$$\text{So, } S_k < 1750$$

$$\frac{k}{2} (2(4) + (k-1)3) < 1750$$

$$\frac{k}{2} (8 + 3k - 3) < 1750$$

$$\frac{k}{2} (3k + 5) < 1750$$

$$\frac{3k^2}{2} + \frac{5k}{2} < 1750$$

$$\frac{3k^2}{2} + \frac{5k}{2} - 1750 < 0$$

$$3k^2 + 5k - 3500 < 0$$

$$\therefore \boxed{(3k - 100)(k + 35) < 0}$$

- d) Either  $3k = 100$  or  $k = -35$

Since no. of rows,  $k$  has to be positive,

$$3k = 100 \Rightarrow k = \frac{100}{3} = 33.\bar{3}$$

$$\boxed{k = 33}$$

Q10a)

i)  $y = x^2(x-2)$

$$y = x^3 - 2x^2$$

When  $x=0, y=0$ 

When  $y=0, x^3 - 2x^2 = 0$

$$x^2(x-2) = 0$$

$x=0$  or  $x=0$  or  $x=2$

ii)  $y = x(6-x)$


$$y = 6x - x^2$$

When  $x=0, y=0$ 

When  $y=0, 6x - x^2 = 0$

$$x(6-x) = 0$$

$x=0$  or  $x=6$



b) To find the coordinates of the points where the graphs intersect, solve simultaneous eqs

$$y = x^3 - 2x^2 \quad \textcircled{1}$$

$$y = 6x - x^2 \quad \textcircled{2}$$

Substitute  $\textcircled{1}$  into  $\textcircled{2}$

$$x^3 - 2x^2 = 6x - x^2$$

$$x^3 - x^2 - 6x = 0$$

$$x(x^2 - x - 6) = 0$$

Either  $x=0$  or  $x^2 - x - 6 = 0$

For  $x^2 - x - 6 = 0$ ,  $(x-3)(x+2) = 0$

Either  $x=3$  or  $x=-2$

Substitute into  $\textcircled{2}$  for  $y$ :

$$\text{When } x=0, y = 6(0) - 0^2 = \underline{0}$$

$$\text{When } x=3, y = 6(3) - 3^2 = 18 - 9 = \underline{9}$$

$$\text{When } x=-2, y = 6(-2) - (-2)^2 = -12 - 4 = \underline{-16}$$

The coordinates of the points where the graphs intersect are:

$$\boxed{(0,0), (3,9), \text{ and } (-2,-16)}$$