

PhysicsAndMathsTutor.com

# C1 January 2009 (MA)

$$Q1a) 125^{1/3} = \sqrt[3]{125} \quad \boxed{= 5}$$

$$b) 125^{-2/3} = \frac{1}{125^{2/3}} = \frac{1}{(\sqrt[3]{125})^2} = \frac{1}{5^2} \quad \boxed{= \frac{1}{25}}$$

$$Q2) \int (12x^5 - 8x^3 + 3) dx = \frac{12x^6}{6} - \frac{8x^4}{4} + 3x + C$$

$$= \boxed{2x^6 - 2x^4 + 3x + C}$$

$$Q3) (\sqrt{7} + 2)(\sqrt{7} - 2) = 7 - 2\sqrt{7} + 2\sqrt{7} - 4$$

$$\boxed{= 3}$$

$$Q4) f'(x) = 3x^2 - 3x^{1/2} - 7$$

$$f(x) = \int (3x^2 - 3x^{1/2} - 7) dx$$

$$= \frac{3x^3}{3} - \frac{3x^{3/2}}{3/2} - 7x + C$$

$$= x^3 - 2x^{3/2} - 7x + C$$

$$= \underline{x^3 - 2x\sqrt{x} - 7x + C}$$

Since (4, 22) lies on  $f(x)$ , substitute in  $x=4$  and  $f(x)=22$ :

$$22 = x^3 - 2x\sqrt{x} - 7x + C$$

$$22 = (4)^3 - 2(4)(\sqrt{4}) - 7(4) + C$$


$$22 = 64 - 16 - 28 + c$$

$$22 = 20 + c$$


$$\therefore \underline{c = 2}$$

$$f(x) = x^3 - 2x\sqrt{x} - 7x + 2$$

Q5a)  $y = f(x+3)$  - transformation of  $-3$  along the  $x$ -axis:



b)  $y = f(-x)$  - reflection in the  $y$ -axis:



$$\begin{aligned}
 \text{Q6a)} \quad \frac{2x^2 - x^{3/2}}{\sqrt{x}} &= \frac{2x^2 - x^{3/2}}{x^{1/2}} \\
 &= \frac{2x^2}{x^{1/2}} - \frac{x^{3/2}}{x^{1/2}} \\
 &= \boxed{2x^{3/2} - x^1}
 \end{aligned}$$

$$\begin{aligned}
 \text{b)} \quad y &= 5x^4 - 3 + \frac{2x^2 - x^{3/2}}{\sqrt{x}} \\
 y &= 5x^4 - 3 + 2x^{3/2} - x \\
 \frac{dy}{dx} &= 20x^3 + 3x^{1/2} - 1 \\
 &= \boxed{20x^3 + 3\sqrt{x} - 1}
 \end{aligned}$$

Q7)  $kx^2 + 4x + (5-k) = 0$  has 2 distinct real solutions.

a) If a quadratic has 2 distinct real roots then the discriminant is greater than 0.

$$\therefore b^2 - 4ac > 0$$

$$4^2 - (4)(k)(5-k) > 0$$

$$16 - 20k + 4k^2 > 0$$

$$4k^2 - 20k + 16 > 0$$

$$\therefore \boxed{k^2 - 5k + 4 > 0}$$


b) For  $k^2 - 5k + 4 = 0$

$$(k-4)(k-1) = 0$$

Either  $k=4$  or  $k=1$

Set of possible values for  $k$ :

$$\boxed{k < 1 \text{ or } k > 4}$$



Choosing values 'above' the x-axis since  $b^2 - 4ac > 0$

Q8a)  $y = (x+1)^2(2-x)$

When  $x=1$ ,  $y = (1+1)^2(2-1) = (2)^2(1) = 4$

$$\therefore \boxed{a=4}$$

bi)  $y = (x+1)^2(2-x)$

When  $x=0$ ,  $y = (0+1)^2(2-0) = (1)^2(2) = \underline{2}$

When  $y=0$ ,  $(x+1)^2(2-x) = 0$

Either  $x=-1$  or  $x=-1$  or  $x=2$


ii)  $y = \frac{2}{x}$

When  $x=0$ ,  $y \rightarrow \infty$

When  $x=-0$ ,  $y \rightarrow -\infty$

When  $y=0$ ,  $x \rightarrow \infty$

When  $y=-0$ ,  $x \rightarrow -\infty$



c) For the equation  $(x+1)^2(2-x) = \frac{2}{x}$ ,

there are 2 real solutions, since there are 2 intersections between the curves

Q9) First term =  $a$ . common difference =  $d$

$$U_n = a + (n-1)d$$

a)  $U_{18} = a + (18-1)d$

$$\boxed{25 = a + 17d} \quad (1)$$

$$U_{21} = a + (21-1)d$$

$$\boxed{32.5 = a + 20d} \quad (2)$$

$$b) \quad \textcircled{2} - \textcircled{1} : 7.5 = 3d$$

$$\boxed{\therefore d = 2.5}$$

Substitute into  $\textcircled{1}$  for  $a$ :

$$25 = a + 17d$$

$$25 = a + 17(2.5)$$

$$25 = a + 42.5$$

$$\therefore \boxed{a = -17.5}$$

$$c) \quad S_n = 2750$$

$$\text{Since } S_n = \frac{n}{2} (2a + (n-1)d)$$

$$S_n = \frac{n}{2} (2(-17.5) + (n-1)2.5)$$

$$S_n = \frac{n}{2} (-35 + 2.5n - 2.5)$$

$$S_n = \frac{n}{2} (2.5n - 37.5)$$

$$S_n = \frac{2.5n^2}{2} - \frac{37.5n}{2}$$

$$2S_n = 2.5n^2 - 37.5n$$

$$4S_n = 5n^2 - 75n$$

$$4S_n = n(5n - 75)$$

Substitute in  $S_n = 2750$

$$4 \times 2750 = n(5n - 75)$$

$$4 \times 550 = n(n - 15)$$

$$\therefore n^2 - 15n = 55 \times 40$$

d)  $n^2 - 15n = 55 \times 40$

$$n^2 - 15n = 2200$$

$$n^2 - 15n - 2200 = 0$$

$$(n - 55)(n + 40) = 0$$

Since  $n > 0$ ,  $n = 55$

Q10a) The line  $l_1$  passes through  $A(2, 5)$  and has gradient  $-\frac{1}{2}$ .

Equation of  $l_1$  :  $y - y_1 = m(x - x_1)$

Using  $A(2, 5)$   
and  $m = -\frac{1}{2}$

$$y - 5 = -\frac{1}{2}(x - 2)$$

$$2(y - 5) = -1(x - 2)$$

$$2y - 10 = -x + 2$$

$$2y = -x + 12$$

$$y = -\frac{x}{2} + 6$$

b) Using  $x = -2, y = 7$ :

$$y = -\frac{x}{2} + 6$$

$$7 = -\frac{(-2)}{2} + 6$$

$$7 = 1 + 6$$

$$\underline{7 = 7}$$

$\therefore$  point B  $(-2, 7)$  lies on  $l$ ,

c) length  $AB = \sqrt{(x_2 - x_1)^2 + (y_2 - y_1)^2}$

$$= \sqrt{(-2 - 2)^2 + (7 - 5)^2}$$

$$= \sqrt{(-4)^2 + 2^2}$$

$$= \sqrt{20} = \sqrt{4 \times 5} = \sqrt{4} \sqrt{5}$$

$$\boxed{= 2\sqrt{5} \text{ units}}$$

d) point C lies on  $l$ , and has  $x$ -coordinate equal to  $p$ . length AC is 5 units.

$$\therefore \sqrt{(x_2 - x_1)^2 + (y_2 - y_1)^2} = 5$$

$$\sqrt{(2 - p)^2 + (5 - (-\frac{1}{2}p + 6))^2} = 5$$

$$(2 - p)^2 + (5 - (-\frac{1}{2}p + 6))^2 = 25$$

$$(2 - p)(2 - p) + (5 + \frac{1}{2}p - 6)(5 + \frac{1}{2}p - 6) = 25$$



$$(2-p)(2-p) + (-1 + \frac{1}{2}p)(-1 + \frac{1}{2}p) = 25$$

$$(2-p)(2-p) + (\frac{1}{2}p - 1)(\frac{1}{2}p - 1) = 25$$

$$4 - 4p + p^2 + \frac{1}{4}p^2 - p + 1 = 25$$

$$p^2 - 4p + 4 + \frac{1}{4}p^2 - p + 1 = 25$$

$$p^2 - 4p + 4 + \frac{1}{4}p^2 - p + 1 - 25 = 0$$

$$\frac{5}{4}p^2 - 5p - 20 = 0$$

$$5p^2 - 20p - 80 = 0$$

$$\therefore \boxed{p^2 - 4p - 16 = 0}$$

Q11)  $y = 9 - 4x - \frac{8}{x}$ ,  $x > 0$

a)  $y = 9 - 4x - 8x^{-1}$

$$\frac{dy}{dx} = -4 + 8x^{-2}$$

$$= -4 + \frac{8}{x^2} = \frac{8}{x^2} - 4$$

When  $x=2$ ,  $\frac{dy}{dx} = \frac{8}{2^2} - 4 = \underline{-2}$

$\therefore$  the gradient of the tangent at  $x=$   
is  $-2$

Also, when  $x=2, y=9-4(2)-\frac{8}{2}$

$$y=9-8-4$$

$$\underline{y=-3}$$

Equation of normal:  $y-y_1 = m(x-x_1)$

Using  $P(2, -3)$   
and  $m=-2$

$$y - (-3) = -2(x-2)$$

$$y+3 = -2x+4$$

$$y = -2x+1$$

$$\boxed{y=1-2x}$$

b) Gradient of tangent at P is  $-2$ , so gradient of normal at P is  $\frac{1}{2}$ .

Equation of normal:  $y-y_1 = m(x-x_1)$

Using  $P(2, -3)$   
and  $m=\frac{1}{2}$

$$y - (-3) = \frac{1}{2}(x-2)$$

$$2(y+3) = 1(x-2)$$

$$2y+6 = x-2$$

$$2y = x-8$$

$$\boxed{y = \frac{x}{2} - 4}$$

c) For the tangent at P, when this meets the x-axis at A,  $y=0$

$$\therefore y = 1 - 2x$$

$$0 = 1 - 2x$$

$$2x = 1$$

$$x = \frac{1}{2}$$

$\therefore A$  is at  $(\frac{1}{2}, 0)$

For the normal at P, when this meets the x-axis at B,  $y=0$

$$\therefore y = \frac{x}{2} - 4$$


$$0 = \frac{x}{2} - 4$$

$$\frac{x}{2} = 4$$

$$x = 8$$

$\therefore B$  is at  $(8, 0)$

Find the area of the triangle APB



Length  $AP = \sqrt{(x_2 - x_1)^2 + (y_2 - y_1)^2}$

$$= \sqrt{\left(2 - \frac{1}{2}\right)^2 + (-3 - 0)^2}$$

$$= \sqrt{\left(\frac{3}{2}\right)^2 + (-3)^2}$$

$$= \sqrt{\frac{9}{4} + 9}$$

$$= \sqrt{\frac{45}{4}}$$

$$= \frac{\sqrt{45}}{\sqrt{4}} = \frac{\sqrt{9 \times 5}}{2} = \frac{\sqrt{9} \sqrt{5}}{2}$$

$$= \frac{3\sqrt{5}}{2} \text{ units}$$


---

$$\begin{aligned}\text{Length } PB &= \sqrt{(x_2 - x_1)^2 + (y_2 - y_1)^2} \\ &= \sqrt{(2 - 8)^2 + (-3 - 0)^2} \\ &= \sqrt{(-6)^2 + (-3)^2} \\ &= \sqrt{36 + 9} \\ &= \sqrt{45} = \sqrt{9 \times 5} = \sqrt{9} \sqrt{5} \\ &= \underline{3\sqrt{5} \text{ units}}\end{aligned}$$

$$\begin{aligned}\text{Area } \triangle APB &= \frac{1}{2} bh = \frac{1}{2} (PB)(AP) \\ &= \frac{1}{2} (3\sqrt{5}) \left( \frac{3\sqrt{5}}{2} \right) \\ &= \frac{1}{2} \left( \frac{45}{2} \right) \\ &= \frac{45}{4} \\ &= \boxed{11.25 \text{ units}^2}\end{aligned}$$