Write your name here		
Surname	Other na	mes
Pearson Edexcel GCE	Centre Number	Candidate Number
Core Mat Advanced Subsid		s C1
Wednesday 16 May 201	•	Paper Reference 6663/01
Time: 1 hour 30 minute	<b>-5</b>	10000,01

## Calculators may NOT be used in this examination.

## Instructions

- Use **black** ink or ball-point pen.
- If pencil is used for diagrams/sketches/graphs it must be dark (HB or B). Coloured pencils and highlighter pens must not be used.
- **Fill in the boxes** at the top of this page with your name, centre number and candidate number.
- Answer **all** questions and ensure that your answers to parts of questions are clearly labelled.
- Answer the questions in the spaces provided
  - there may be more space than you need.
- You should show sufficient working to make your methods clear. Answers without working may not gain full credit.

## Information

- The total mark for this paper is 75.
- The marks for **each** question are shown in brackets
  - use this as a guide as to how much time to spend on each question.

## **Advice**

- Read each question carefully before you start to answer it.
- Try to answer every question.
- Check your answers if you have time at the end.

Turn over ▶



P51518A
©2018 Pearson Education Ltd.
1/1/1/1/1/1/1/



1. (i) Simplify

$$\sqrt{48}-\frac{6}{\sqrt{3}}$$

Write your answer in the form  $a\sqrt{3}$ , where a is an integer to be found.

**(2)** 

(ii) Solve the equation

$$3^{6x-3} = 81$$

Write your answer as a rational number	r.
--	----

	(1)
	•
١.	•




P 5 1 5 1 8 A 0 2 3 2

	Leave blank
Question 1 continued	
	Q1
/FD 4.15 1 N	
(Total 5 marks)	



Given

$$y = 3\sqrt{x} - 6x + 4, \qquad x > 0$$

(a) find  $\int y dx$ , simplifying each term.

**(3)** 

- (b) (i) Find  $\frac{dy}{dx}$ 
  - (ii) Hence find the value of x such that  $\frac{dy}{dx} = 0$

(4)









10	Ρ
-	lij.
11	АÜ
÷	Ħ.
۵	ø
ш	
÷	i
5	ь
U	PA.
	ч
=	=
1	×
ы	÷
	7
ŀ	=
	₽
4	-
÷	÷
Ц	1
ы	н
片	_
ъ.	
-	=
Ώ	⊏
Ξ	_
2	30
-	Si
200	-
10.1	
ŧ	=
	6
E.	ı
_	_
72	⇗
1	_
٠	. 1
r	٦
•	J
-	ĸ.
1	_1
_	=

			ı
Ä			
		7.1	
Š			
X			
<u>/ i</u>			
V			
		N	
М	И	Ы	k
K		Ņ	k
K		N	k
K		V	k
K		N	į
K		N 2	į
K		N 2	
K		N 2	į
K		N Z	l de
K		N 2	
K		N 2	
		N 2 2	
		N 2 2 X	
K		N 2 2	
K			
K			
K			
K			
K			
K			k ive v
K			
K			k it is a contract of
K			k i kekk Krkink
K			k identification
K			k iverview
K			
K			
K			L idea or cardia
K			k items created
K			
K			
K			k identify the control of the contro
K			k iver crystylly er k
K			k idea or did to the
K			
K			
K			
K			

$\times$			
Χ.,	×.		
ᄼ	/		
×			
Κ.	×	-2	Ś
ğ	d		ľ
U	۰,		ŀ
	×		2
Þ	D	п	
×Ι			
×	Z	$\leq$	
M	Ď		۰
ΚĪ	м		i
×	. 7		
Χ.,			,
Š			ŀ
X	7	7	۰
	×		
ğ		и	
v	ы	r	
~	7		
×			
Κ.	×	2	Ĺ
~	X D		7
×	△	к	
Κŝ	≂	7	ť
. Ж	u	S.	
V	в	7	۰
ø			
X			
K.	₹		
-/3	2	ĸ	
V		×	
ĸũ	ĸ	è	i
×			
۲.	Χ	2	
Ø	Ŵ	ь	
K			
ĸ.	х		
Ø	k	6	
M	12	7	٦
< £	4	4	è
×	7	₹	
×.	m		ļ
4	Ċ		
$\times$	-	7	
×Ψ	ю,	ù	í
.4	ò		,
v	-		ŀ
KJ	ø	щ	e
×			
N.	ĸ		
И			
K			
O	ы		
N.	r		
×J	6	ú	
1	-2	≾	
M	-		
ĸ.	×	2	
А	ø	۰	ļ
V			
ĸ^	ú	è	ś
- 24	۳	75	٩

Question 2 continued		Lea bla
		Q2
	(Total 7 marks)	



3.  $f(x) = x^2 - 10x + 23$ 

(a) Express f(x) in the form  $(x + a)^2 + b$ , where a and b are constants to be found.

**(2)** 

(b) Hence, or otherwise, find the exact solutions to the equation

$$x^2 - 10x + 23 = 0$$

**(2)** 

(c) Use your answer to part (b) to find the larger solution to the equation

$$y - 10y^{0.5} + 23 = 0$$

Write your solution in the form  $p + q\sqrt{r}$ , where p, q and r are integers.

Leave

10	Ρ
-	lij.
11	АÜ
÷	Ħ.
۵	ø
ш	
÷	i
5	ь
U	PA.
	ч
=	=
1	×
ы	÷
	7
ŀ	=
	₽
4	-
÷	÷
Ц	1
ы	н
片	_
ъ.	
-	=
Ώ	⊏
Ξ	_
2	30
-	Si
200	-
10.1	
ŧ	=
	6
E.	ı
_	_
72	~
1	_
٠	. 1
r	٦
•	J
-	ĸ.
1	_1
_	=

Question 3 continued		blank
		Q3
	(Total 6 marks)	



4.	Each year, Andy pays into a savings scheme. In year one he pays in £600. His payments increase by £120 each year so that he pays £720 in year two, £840 in year three and so on, so that his payments form an arithmetic sequence.
	(a) Find out how much Andy pays into the savings scheme in year ten. (2)
	Kim starts paying money into a different savings scheme at the same time as Andy. In year one she pays in £130. Her payments increase each year so that she pays £210 in year two, £290 in year three and so on, so that her payments form a different arithmetic sequence.
	At the end of year N, Andy has paid, in total, twice as much money into his savings scheme as Kim has paid, in total, into her savings scheme.
	(b) Find the value of $N$ . (5)



	Leave
	blank
Overtion 4 continued	June
Question 4 continued	



tion 4 continued	

		ú	á	ø
1		3	ľ	3
1	i	ú	ï	ì
4	Ŀ	ě	è	ı
4	Ė	b	ė	ø
1	Ė	ă	Ė	
ì	è	ø	è	ø
	٦	d	ķ	ù
1	Ĺ	I	ī	)
'n	i	ė		ė
ì	e	ė	ŧ	ú
ì				ė
Ì	Ł	_		i
1	г	7	7	-
i	6	ė	ė	i
ì	é	ĕ	2	_
j	Ξ	Ξ	Ξ	3
Ē	7		ī	7
į	١	ı	۰	ı
ŧ		7		7
J	Þ	e		ij
å	ì		i	
4	þ	b	è	ø
1	è	ä	è	
Þ	9	4	ė	
1	Ę		S	2
J	ø	P	9	7
9				
4	r			Ŧ
ď	þ	P	٩	h
ļ	Ġ		à	p
9	i	Ę	ġ	ø
1	ė		_	ì
÷			Ī	7
ì	è		ġ	ĸ
4	L	٠	i	)
٠	ă	ă	í	ŀ
1	ľ	ï	1	١

Question 4 continued	Leave blank
	Q4
(Total 7 marks)	



5.




Figure 1

Figure 1 shows the sketch of a curve with equation  $y = f(x), x \in \mathbb{R}$ .

The curve crosses the y-axis at (0, 4) and crosses the x-axis at (5, 0).

The curve has a single turning point, a maximum, at (2, 7).

The line with equation y = 1 is the only asymptote to the curve.

- (a) State the coordinates of the turning point on the curve with equation y = f(x 2). (1)
- (b) State the solution of the equation f(2x) = 0 (1)
- (c) State the equation of the asymptote to the curve with equation y = f(-x). (1)

Given that the line with equation y = k, where k is a constant, meets the curve y = f(x) at only one point,

(d) state the set of possible values for k.

	Leave
	blank
Overtion 5 continued	) June
Question 5 continued	



estion 5 continued	

2	φ
-	ы
1.1	ιũ
ы	ш
÷	i
₽	-
٠.	ès
×	L
	7
BF 4	m.
U	9
=	-i
100	-
ш	Ŀ
	_
ŀ	Ė
э.	
1	90
4	_
Ξ	☶
=	=
	٠ů
Ш	ч
1L	_
ŀ	_
=	ė
~	iα
m	_
1	~
1	30
_	
ы	÷
Œ	7
r	٦
ъ.	J
=	9
2	_
	=
-	i.
г	٦
-	ø
P	ď
£	4

Question 5 continued	Leave blank
	Q5
(Total 5 marks)	



blank

**6.** A sequence  $a_1, a_2, a_3, \dots$  is defined by

$$a_{_{1}}=4$$
 
$$a_{_{n+1}}=\frac{a_{_{n}}}{a_{_{n}}+1}, \qquad n\geqslant 1,\, n\in\mathbb{N}$$

(a) Find the values of  $a_2$ ,  $a_3$  and  $a_4$ 

Write your answers as simplified fractions.

(3)

Given that

$$a_n = \frac{4}{pn+q}$$
, where p and q are constants

(b) state the value of p and the value of q.

- (c) Hence calculate the value of N such that  $a_N = \frac{4}{321}$  (2)

ì	ø	ij	۴	,
٩	7	힉	b	i
1	L	Ħ		i
1	ė	ě	Ħ	ı
1	À	ù	ä	ø
ł	Ė	ă	Ė	ı
	4	d	ò	ø
J	ξ	a	L	ı
				٩
'n	ŕ	à	n	i
1	ы	y		9
'n	i	ė		i
ì	×	b	ø	â
ì			_	i
	7	7	7	7
ł	þ		Ė	ı
	٢			
ž	٠	ŧ	á	ß
á	e		Ξ	
c	Ξ	Ξ	Ξ	3
3	-	7		۰
ì	r	i		i
ļ	b	ě		ı
1	L	_	_	1
1	Г	7		7
4	ŧ		ŧ	ı
1	ρ	٧	ř	ø
	Ė	ä	Ė	ı
þ	۹	6	ė	
1	Ġ		2	7
s	ρ	ø	3	•
ì				
1	h	Ė		i
1	L	i	Ĺ.	
1	r	"	7	١
4	þ	ú	ø	,
9	÷	5	3	ø
ï			_	i
÷	1	•	1	7
٠,	à		ò	c
4	ľ		1	0
	9	Ħ	p	
Ì	Р	7	٩	١
1	h	Ė	ď	ı

	Š	
¥į		

DO NOT WRITE IN THIS AREA

Question 6 continued	Leave blank
Question o continueu	
	<b>Q6</b>
(Total 7 marks)	



7. The equation $20x^2 = 4kx - 13kx^2 + 2$ , where k is a constant, has n	no real roots.
(a) Show that $k$ satisfies the inequality	
$2k^2 + 13k + 20 < 0$	(4
(b) Find the set of possible values for $k$ .	(4



3		
1		
J		
1		
'n		
3		
Ì		
4		
1		
1		
þ		
1		
1		
1		
1		
1		
4		
1		

Question 7 continued	Leave blank
	<b>Q7</b>
(Total 8 marks)	



blank

8.




Figure 2 shows the straight line  $l_1$  with equation 4y = 5x + 12

(a) State the gradient of  $l_1$ 

**(1)** 

The line  $l_2$  is parallel to  $l_1$  and passes through the point E (12, 5), as shown in Figure 2.

(b) Find the equation of  $l_2$ . Write your answer in the form y = mx + c, where m and c are constants to be determined.

(3)

The line  $l_2$  cuts the x-axis at the point C and the y-axis at the point B.

- (c) Find the coordinates of
  - (i) the point B,
  - (ii) the point C.

**(2)** 

The line  $l_1$  cuts the y-axis at the point A.

The point D lies on  $l_1$  such that ABCD is a parallelogram, as shown in Figure 2.

(d) Find the area of ABCD.

	]
uestion 8 continued	



stion 8 continued	

1	
∾.	
ш	
-	
_	
ď	
II A	
ഗ	
_	
_	
- 1	
$\vdash$	
_	
==	
Sec. 2	
ш	
-	
<del>-</del>	
<u> </u>	
-	
œ	
1	
$\rightarrow$	
-	
1	
$\vdash$	
-	
<b>(</b> )	
-	
=	
~	
-	
$\Box$	
$\overline{}$	
-	

Question 8 continued	Leave blank
	<b>Q8</b>
(Total 8 marks)	



**9.** The curve C has equation y = f(x), where

$$f'(x) = (x-3)(3x+5)$$

Given that the point P(1, 20) lies on C,

(a) find f(x), simplifying each term.

(5)

(b) Show that

$$f(x) = (x - 3)^2(x + A)$$

where *A* is a constant to be found.

**(3)** 

(c) Sketch the graph of C. Show clearly the coordinates of the points where C cuts or meets the x-axis and where C cuts the y-axis.

**(4)** 


	L
Question 9 continued	b:
guestion 5 continued	



estion 9 continued	

Leave

Question 9 continued	blank
	Q9
(Total 12 marks)	
	27



**10.** 




Figure 3

Figure 3 shows a sketch of part of the curve C with equation

$$y = \frac{1}{2}x + \frac{27}{x} - 12, \qquad x > 0$$

The point *A* lies on *C* and has coordinates  $\left(3, -\frac{3}{2}\right)$ .

(a) Show that the equation of the normal to C at A can be written as 10y = 4x - 27 (5)

The normal to C at A meets C again at the point B, as shown in Figure 3.

(b) Use algebra to find the coordinates of B.



	Leave
	blank
Question 10 continued	
	1



	Lea
Question 10 continued	blaı
Question to continued	
	_
	_
	_
	-
	_
	_
	_
	_
	-
	_
	_
	_
	-
	_
	-
	_
	-
	_
	_
	_
	_
	-
	_
	_
	_
	-
	_
	-
	_
	_
	_

	Le bl
Question 10 continued	



uestion 10 continued	Leave blank
uconon 10 continucu	
	_
	_
	_
	-
	_
	_
	-
	-
	-
	-
	-
	-
	_
	_
	-
	-
	_
	_
	_
	_
	_
	_
	_
	_
	_
	_
	Q10
(Total 10 marks	)
TOTAL FOR PAPER: 75 MARK	S
END	