

GCE Examinations
Advanced Subsidiary

Core Mathematics C1

Paper F

MARKING GUIDE

This guide is intended to be as helpful as possible to teachers by providing concise solutions and indicating how marks could be awarded. There are obviously alternative methods that would also gain full marks.

Method marks (M) are awarded for knowing and using a method.

Accuracy marks (A) can only be awarded when a correct method has been used.

(B) marks are independent of method marks.





Written by Shaun Armstrong

© *Solomon Press*

These sheets may be copied for use solely by the purchaser's institute.

C1 Paper F – Marking Guide

1. $x^4 - 5x^2 - 14 = 0$, $(x^2 + 2)(x^2 - 7) = 0$ M1
 $x^2 = -2$ (no solutions) or 7 A1
 $x = \pm\sqrt{7}$ A1 (3)
-
2. $= \frac{2}{3\sqrt{5}+7} \times \frac{3\sqrt{5}-7}{3\sqrt{5}-7} = \frac{6\sqrt{5}-14}{45-49} = \frac{7}{2} - \frac{3}{2}\sqrt{5}$ M2 A1 (3)
-
3. (a) $x = (\sqrt[3]{27})^2 = 3^2 = 9$ M1 A1
 (b) $= (\frac{9}{4})^{-\frac{1}{2}} = \sqrt{\frac{4}{9}} = \frac{2}{3}$ M1 A1 (4)
-
4. cubic, coeff of $x^3 = 1$, crosses x -axis at $(-1, 0)$, touches at $(3, 0)$
 $\therefore y = (x + 1)(x - 3)^2$ M1 A1
 $= (x + 1)(x^2 - 6x + 9)$
 $= x^3 - 6x^2 + 9x + x^2 - 6x + 9$ M1
 $= x^3 - 5x^2 + 3x + 9$
 $\therefore a = -5, b = 3, c = 9$ A2 (5)
-
5. (a) $y = \frac{1}{2}x^2 - \frac{3}{2}x^{-2}$ M1 A1
 $\frac{dy}{dx} = x + 3x^{-3}$ M1 A1
 (b) $\frac{d^2y}{dx^2} = 1 - 9x^{-4} = \frac{x^4 - 9}{x^4}$ M1 A1 (6)
-
6. (a)  B2
 B3
 (b) $x^2 - 4x + 4 > 2x - 1$
 $x^2 - 6x + 5 > 0$
 $(x - 1)(x - 5) > 0$ 
 $x < 1$ or $x > 5$ M1
 M1
 A1 (8)
-
7. (a) $\frac{dy}{dx} = \frac{1}{2} + x^{-2}$ M1 A1
 $\text{grad} = \frac{1}{2} + 2^{-2} = \frac{3}{4}$ M1 A1
 (b) $x = 2 \therefore y = \frac{7}{2}$ B1
 $y - \frac{7}{2} = \frac{3}{4}(x - 2)$ M1
 $4y - 14 = 3x - 6$
 $3x - 4y + 8 = 0$ A1
 (c) at B, $\text{grad} = \frac{3}{4}$
 $\therefore \frac{1}{2} + x^{-2} = \frac{3}{4}$ M1
 $x^2 = 4, x = 2$ (at A), -2 A1
 $\therefore B(-2, \frac{5}{2})$ A1 (10)

8.	(a)	$y - 3 = \frac{3}{2}(x - 5)$	M1	
		$y = \frac{3}{2}x - \frac{9}{2}$	A1	
	(b)	$3x - 4(\frac{3}{2}x - \frac{9}{2}) + 3 = 0$	M1	
		$x = 7$	A1	
		$\therefore B(7, 6)$	A1	
	(c)	$= (\frac{5+7}{2}, \frac{3+6}{2}) = (6, \frac{9}{2})$	M1 A1	
	(d)	$l_2: y = \frac{3}{4}x + \frac{3}{4} \therefore \text{grad} = \frac{3}{4}$	B1	
		$\therefore y - \frac{9}{2} = \frac{3}{4}(x - 6)$	M1	
		$y = \frac{3}{4}x$	A1	
		when $x = 0, y = 0 \therefore$ passes through origin	A1	(11)

9.	(a)	$a + 2d = 5\frac{1}{2}$ (1)	B1	
		$\frac{4}{2}(2a + 3d) = 22\frac{3}{4}$ (2)	M1 A1	
		(2) $\Rightarrow 4a + 6d = 22\frac{3}{4}$		
		(1) $\Rightarrow 3a + 6d = 16\frac{1}{2}$		
		subtracting, $a = 22\frac{3}{4} - 16\frac{1}{2} = 6\frac{1}{4}$	M1 A1	
		$d = \frac{1}{2}(5\frac{1}{2} - 6\frac{1}{4}) = -\frac{3}{8}$	M1 A1	
	(b)	$6\frac{1}{4} - \frac{3}{8}(n - 1) > 0$	M1	
		$50 - 3(n - 1) > 0$		
		$n < 17\frac{2}{3} \therefore 17$ positive terms	M1 A1	
	(c)	$= S_{17} = \frac{17}{2} [12\frac{1}{2} + (16 \times -\frac{3}{8})]$	M1	
		$= \frac{17}{2} (12\frac{1}{2} - 6) = \frac{17}{2} \times \frac{13}{2} = \frac{221}{4} = 55\frac{1}{4}$	A1	(12)

10.	(a)	grad = $8 - 2 = 6$	B1	
		$\therefore y - 1 = 6(x - 1)$	M1	
		$y = 6x - 5$	A1	
	(b)	$y = \int (8x - \frac{2}{x^3}) dx$		
		$y = 4x^2 + x^{-2} + c$	M1 A2	
		(1, 1) $\therefore 1 = 4 + 1 + c$		
		$c = -4$	M1	
		$y = 4x^2 + x^{-2} - 4$	A1	
	(c)	$4x^2 + x^{-2} - 4 = 0$		
		$4x^4 - 4x^2 + 1 = 0$	M1	
		$(2x^2 - 1)^2 = 0$	M1	
		$x^2 = \frac{1}{2}$		
		$x = \pm \frac{1}{\sqrt{2}}$	A1	
		$x = \pm \frac{1}{\sqrt{2}} \times \frac{\sqrt{2}}{\sqrt{2}} = \pm \frac{1}{2}\sqrt{2}$	M1 A1	(13)

Total **(75)**

Performance Record – C1 Paper F

Question no.	1	2	3	4	5	6	7	8	9	10	Total
Topic(s)	quad.	surds	indices	cubic	diff.	curve sketch, unequal.	diff., tangents	straight lines	AP	integr., tangent	
Marks	3	3	4	5	6	8	10	11	12	13	75
Student											