

Solutionbank C1

Edexcel Modular Mathematics for AS and A-Level

Algebraic fractions

Exercise A, Question 1

Question:

Factorise completely

(a) $2x^3 - 13x^2 - 7x$

(b) $9x^2 - 16$

(c) $x^4 + 7x^2 - 8$

Solution:

(a)

$$\begin{aligned} 2x^3 - 13x^2 - 7x \\ &= x(2x^2 - 13x - 7) \\ &= x(2x^2 + x - 14x - 7) \\ &= x[x(2x + 1) - 7(2x + 1)] \\ &= x(2x + 1)(x - 7) \end{aligned}$$

x is a common factor

So take x outside the bracket.

For the quadratic, $ac = -14$ and

$$1 - 14 = -13 = b$$

Factorise

(b)

$$\begin{aligned} 9x^2 - 16 \\ &= (3x)^2 - 4^2 \\ &= (3x + 4)(3x - 4) \end{aligned}$$

This is a difference of two

squares, $(3x)^2$ and 4^2

$$\text{Use } x^2 - y^2 = (x + y)(x - y)$$

(c)

$$\begin{aligned} x^4 + 7x^2 - 8 \\ &= y^2 + 7y - 8 \\ &= y^2 - y + 8y - 8 \\ &= y(y - 1) + 8(y - 1) \\ &= (y - 1)(y + 8) \\ &= (x^2 - 1)(x^2 + 8) \\ &= (x + 1)(x - 1) \\ &\quad (x^2 + 8) \end{aligned}$$

squares,

$$\text{Let } y = x^2$$

$$ac = -8 \text{ and } -1 + 8 = +7 = b$$

Factorise

Replace y by x^2

$x^2 - 1$ is a difference of two

$$\text{so use } x^2 - y^2 = (x + y)(x - y)$$

Solutionbank C1

Edexcel Modular Mathematics for AS and A-Level

Algebraic fractions

Exercise A, Question 2

Question:

Find the value of

(a) $81^{\frac{1}{2}}$

(b) $81^{\frac{3}{4}}$

(c) $81^{-\frac{3}{4}}$

Solution:

(a)

$$\begin{aligned} 81^{1/2} \\ &= \sqrt{81} \\ &= 9 \end{aligned}$$

$$\text{Use } a^{\frac{1}{m}} = m\sqrt[m]{a}, \text{ so } a^{\frac{1}{2}} = \sqrt{a}$$

(b)

$$\begin{aligned} 81^{\frac{3}{4}} \\ &= (\sqrt[4]{81})^3 \quad \text{then cube this} \\ &= 3^3 \quad \sqrt[4]{81} = 3 \text{ because } 3 \times 3 \times 3 \times 3 = 81 \\ &= 27 \end{aligned}$$

$$a^{\frac{n}{m}} = m\sqrt[m]{(a^n)} \text{ or } (m\sqrt[m]{a})^n$$

It is easier to find the fourth root,

4

(c)

$$\begin{aligned} 81^{-\frac{3}{4}} &= \frac{1}{81^{3/4}} \\ &= \frac{1}{27} \end{aligned}$$

$$\text{Use } a^{-m} = \frac{1}{a^m}$$

Use the answer from part (b)

Solutionbank C1

Edexcel Modular Mathematics for AS and A-Level

Algebraic fractions

Exercise A, Question 3

Question:

(a) Write down the value of $8^{\frac{1}{3}}$.

(b) Find the value of $8^{-\frac{2}{3}}$.

Solution:

(a)

$$\begin{aligned} 8^{\frac{1}{3}} &= \sqrt[3]{8} \\ &= 2 \end{aligned}$$

$$\text{Use } a^{\frac{1}{m}} = \sqrt[m]{a}, \text{ so } a^{\frac{1}{3}} = \sqrt[3]{a}$$

$$\sqrt[3]{8} = 2 \text{ because } 2 \times 2 \times 2 = 8$$

(b)

$$8^{-\frac{2}{3}}$$

$$\begin{aligned} 8^{\frac{2}{3}} &= (\sqrt[3]{8})^2 \\ &= 2^2 = 4 \end{aligned} \quad (\sqrt[m]{a})^n$$

$$\text{First find } 8^{\frac{2}{3}} \quad a^{\frac{n}{m}} = \sqrt[m]{(a^n)} \text{ or}$$

$$\begin{aligned} 8^{-\frac{2}{3}} &= \frac{1}{8^{\frac{2}{3}}} \\ &= \frac{1}{4} \end{aligned}$$

$$\text{Use } a^{-m} = \frac{1}{a^m}$$

Solutionbank C1

Edexcel Modular Mathematics for AS and A-Level

Algebraic fractions

Exercise A, Question 4

Question:

(a) Find the value of $125^{\frac{4}{3}}$.

(b) Simplify $24x^2 \div 18x^{\frac{4}{3}}$.

Solution:

(a)

$$\begin{aligned} 125^{\frac{4}{3}} &= (\sqrt[3]{125})^4 \\ &= 5^4 \\ &= 625 \end{aligned}$$

$$a^{\frac{n}{m}} = m\sqrt{(a^n)} \quad \text{or} \quad (m\sqrt{a})^n$$

It is easier to find the cube root,
then the fourth power

$$\sqrt[3]{125} = 5 \quad \text{because} \quad 5 \times 5 \times 5 = 125$$

(b)

$$24x^2 \div 18x^{\frac{4}{3}}$$

$$= \frac{24x^2}{18x^{\frac{4}{3}}} = \frac{4x^2}{3x^{\frac{4}{3}}}$$

by 6

Divide

$$= \frac{4x^{\frac{2}{3}}}{3}$$

$$\text{Use } a^m \div a^n = a^{m-n}$$

$$\left(\text{or } \frac{4}{3}x^{\frac{2}{3}} \right)$$

$$\frac{2}{3}$$

Solutionbank C1

Edexcel Modular Mathematics for AS and A-Level

Algebraic fractions

Exercise A, Question 5

Question:

- (a) Express $\sqrt{80}$ in the form $a\sqrt{5}$, where a is an integer.
- (b) Express $(4 - \sqrt{5})^2$ in the form $b + c\sqrt{5}$, where b and c are integers.

Solution:

(a)

$$\begin{aligned}\sqrt{80} &= \sqrt{16} \times \sqrt{5} \\ &= 4\sqrt{5} \quad (a = 4)\end{aligned}$$

Use $\sqrt{(ab)} = \sqrt{a}\sqrt{b}$

(b)

$$\begin{aligned}(4 - \sqrt{5})^2 &= (4 - \sqrt{5})(4 - \sqrt{5}) \\ &= 4(4 - \sqrt{5}) - \sqrt{5}(4 - \sqrt{5}) \\ &= 16 - 4\sqrt{5} - 4\sqrt{5} + 5 \\ &= 21 - 8\sqrt{5}\end{aligned}$$

Multiply the brackets.
 $\sqrt{5} \times \sqrt{5} = 5$

$(b = 21 \text{ and } c = -8)$

Solutionbank C1

Edexcel Modular Mathematics for AS and A-Level

Algebraic fractions

Exercise A, Question 6

Question:

(a) Expand and simplify $(4 + \sqrt{3})(4 - \sqrt{3})$.

(b) Express $\frac{26}{4 + \sqrt{3}}$ in the form $a + b\sqrt{3}$, where a and b are integers.

Solution:

(a)

$$\begin{aligned} & (4 + \sqrt{3})(4 - \sqrt{3}) \\ &= 4(4 - \sqrt{3}) + \sqrt{3}(4 - \sqrt{3}) \\ &= 16 - 4\sqrt{3} + 4\sqrt{3} - 3 \\ &= 13 \end{aligned}$$

Multiply the brackets.
 $\sqrt{3} \times \sqrt{3} = 3$

(b)

$$\frac{26}{4 + \sqrt{3}} \times \frac{4 - \sqrt{3}}{4 - \sqrt{3}}$$

To rationalise the denominator, multiply top and

$$\text{bottom by } 4 - \sqrt{3}$$

$$= \frac{26(4 - \sqrt{3})}{(4 + \sqrt{3})(4 - \sqrt{3})}$$

$$= \frac{26(4 - \sqrt{3})}{13}$$

answer from part (a)

Use the

$$= 2(4 - \sqrt{3})$$

Divide by 13

$$= 8 - 2\sqrt{3}$$

$$(a = 8 \text{ and } b = -2)$$

Solutionbank C1

Edexcel Modular Mathematics for AS and A-Level

Algebraic fractions

Exercise A, Question 7

Question:

- (a) Express $\sqrt{108}$ in the form $a\sqrt{3}$, where a is an integer.
- (b) Express $(2 - \sqrt{3})^2$ in the form $b + c\sqrt{3}$, where b and c are integers to be found.

Solution:

$$\begin{aligned} \text{(a)} \quad \sqrt{108} &= \sqrt{36} \times \sqrt{3} && \text{Use } \sqrt{(ab)} = \sqrt{a}\sqrt{b} \\ &= 6\sqrt{3} \quad (a = 6) \end{aligned}$$

$$\begin{aligned} \text{(b)} \quad (2 - \sqrt{3})^2 &= (2 - \sqrt{3})(2 - \sqrt{3}) \\ &= 2(2 - \sqrt{3}) - \sqrt{3}(2 - \sqrt{3}) && \text{Multiply the brackets} \\ &= 4 - 2\sqrt{3} - 2\sqrt{3} + 3 && \sqrt{3} \times \sqrt{3} = 3 \\ &= 7 - 4\sqrt{3} \end{aligned}$$

$(b = 7 \text{ and } c = -4)$

Solutionbank C1

Edexcel Modular Mathematics for AS and A-Level

Algebraic fractions

Exercise A, Question 8

Question:

- (a) Express $(2\sqrt{7})^3$ in the form $a\sqrt{7}$, where a is an integer.
- (b) Express $(8 + \sqrt{7})(3 - 2\sqrt{7})$ in the form $b + c\sqrt{7}$, where b and c are integers.
- (c) Express $\frac{6+2\sqrt{7}}{3-\sqrt{7}}$ in the form $d + e\sqrt{7}$, where d and e are integers.

Solution:

(a)

$$\begin{aligned} (2\sqrt{7})^3 &= 2\sqrt{7} \times 2\sqrt{7} \times 2\sqrt{7} && \text{Multiply the 2s.} \\ &= 8(\sqrt{7} \times \sqrt{7} \times \sqrt{7}) \\ &= 8(7\sqrt{7}) && \sqrt{7} \times \sqrt{7} = 7 \\ &= 56\sqrt{7} \quad (a = 56) \end{aligned}$$

(b)

$$\begin{aligned} (8 + \sqrt{7})(3 - 2\sqrt{7}) &= 8(3 - 2\sqrt{7}) + \sqrt{7}(3 - 2\sqrt{7}) && \sqrt{7} \times 2\sqrt{7} = 2 \times 7 \\ &= 24 - 16\sqrt{7} + 3\sqrt{7} - 14 \\ &= 10 - 13\sqrt{7} \\ (b = 10 \text{ and } c = -13) \end{aligned}$$

(c)

$$\frac{6+2\sqrt{7}}{3-\sqrt{7}} \times \frac{3+\sqrt{7}}{3+\sqrt{7}}$$

To rationalise the denominator, multiply top and bottom

$$\begin{aligned} &\text{by } 3 + \sqrt{7} \\ &= \frac{(6+2\sqrt{7})(3+\sqrt{7})}{(3-\sqrt{7})(3+\sqrt{7})} \\ &= \frac{6(3+\sqrt{7}) + 2\sqrt{7}(3+\sqrt{7})}{3(3+\sqrt{7}) - \sqrt{7}(3+\sqrt{7})} \\ &= \frac{18 + 6\sqrt{7} + 6\sqrt{7} + 14}{9 + 3\sqrt{7} - 3\sqrt{7} - 7} \\ &= \frac{32 + 12\sqrt{7}}{2} = 16 + 6\sqrt{7} && \text{Divide by 2} \\ (d = 16 \text{ and } e = 6) \end{aligned}$$

Solutionbank C1

Edexcel Modular Mathematics for AS and A-Level

Algebraic fractions

Exercise A, Question 9

Question:

Solve the equations

(a) $x^2 - x - 72 = 0$

(b) $2x^2 + 7x = 0$

(c) $10x^2 + 9x - 9 = 0$

Solution:

(a)

$$x^2 - x - 72 = 0$$

$$(x + 8)(x - 9) = 0$$

Factorise

$x + 8 = 0, x - 9 = 0$ the equation could be solved using the

Although

$x = -8, x = 9$ formula or 'completing

quadratic

the square', factorisation is quicker.

(b)

$$2x^2 + 7x = 0$$

$$x(2x + 7) = 0$$

$x = 0, 2x + 7 = 0$ forget the $x = 0$ solution.

Don't

$$x = 0, x = -\frac{7}{2}$$

Use

the factor x.

(b)

$$2x^2 + 7x = 0$$

$$x(2x + 7) = 0$$

$x = 0, 2x + 7 = 0$ forget the $x = 0$ solution.

Don't

$$x = 0, x = -\frac{7}{2}$$

Use

the factor x.

(c)

$$10x^2 + 9x - 9 = 0$$

$$(2x + 3)(5x - 3) = 0$$

$$2x + 3 = 0, 5x - 3 = 0$$

$$x = -\frac{3}{2}, x = \frac{3}{5}$$

Factorise

Solutionbank C1

Edexcel Modular Mathematics for AS and A-Level

Algebraic fractions

Exercise A, Question 10

Question:

Solve the equations, giving your answers to 3 significant figures

(a) $x^2 + 10x + 17 = 0$

(b) $2x^2 - 5x - 1 = 0$

(c) $(2x - 3)^2 = 7$

Solution:

(a)

$$x^2 + 10x + 17 = 0$$

$$x = \frac{-b \pm \sqrt{(b^2 - 4ac)}}{2a}$$

quadratic will not factorise.

$$a = 1, b = 10, c = 17$$

$$x = \frac{-10 \pm \sqrt{(100 - 68)}}{2}$$

the formula first.

Since the question requires answers to

3 significant figures, you know that the

$$= \frac{-10 \pm \sqrt{(100 - 68)}}{2}$$

$$= \frac{-10 \pm \sqrt{32}}{2}$$

$$= \frac{-10 \pm 5.656 \dots}{2}$$

at least 4 sig. figs.

$$= \frac{-10 + 5.656 \dots}{2},$$

$$\frac{-10 - 5.656 \dots}{2}$$

$$x = -2.17, x = -7.83$$

Alternative method:

$$x^2 + 10x + 17 = 0$$

$$x^2 + 10x = -17$$

$$(x + 5)^2 - 25 = -17$$

$$(x + 5)^2 = -17 + 25$$

$$(x + 5)^2 = 8$$

$$x + 5 = \pm \sqrt{8}$$

$$x = -5 \pm \sqrt{8}$$

$$x = -5 + \sqrt{8}, x = -5 - \sqrt{8}$$

$$x = -2.17, x = -7.83$$

Use the quadratic formula, quoting

Intermediate working should be to

Divide by 2, and round to 3 sig. figs.

Subtract 17 to get LHS in the required form.

Complete the square for $x^2 + 10x$

Add 25 to both sides

Square root both sides.

Subtract 5 from both sides.

(b)

$$2x^2 - 5x - 1 = 0$$

$$x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$$

$$a = 2, b = -5, c = -1$$

$$x = \frac{5 \pm \sqrt{(-5)^2 - (4 \times 2 \times -1)}}{4}$$

Use the quadratic formula, quoting

the formula first.

$$= \frac{5 \pm \sqrt{(25 + 8)}}{4} = \frac{5 \pm \sqrt{33}}{4}$$

$$= \frac{5 + 5.744 \dots}{4}, \frac{5 - 5.744 \dots}{4}$$

$$x = 2.69, x = -0.186$$

Divide by 4, and round to 3 sig. figs.

(c)

$$(2x - 3)^2 = 7$$

$$2x - 3 = \pm \sqrt{7}$$

The quickest method is to take the square root

of both sides.

$$2x = 3 \pm \sqrt{7}$$

Add 3 to both sides.

$$x = \frac{3 + \sqrt{7}}{2}, x =$$

Divide both

$$\frac{3 - \sqrt{7}}{2}$$

sides by 2

$$x = 2.82, x = 0.177$$

Solutionbank C1

Edexcel Modular Mathematics for AS and A-Level

Algebraic fractions

Exercise A, Question 11

Question:

$$x^2 - 8x - 29 \equiv (x + a)^2 + b,$$

where a and b are constants.

(a) Find the value of a and the value of b .

(b) Hence, or otherwise, show that the roots of

$$x^2 - 8x - 29 = 0$$

are $c \pm d\sqrt{5}$, where c and d are integers to be found.

Solution:

(a)

$$\begin{aligned} x^2 - 8x &= (x - 4)^2 - 16 && \text{Complete the square} \\ x^2 - 8x - 29 &= (x - 4)^2 - 16 - 29 && \text{for } x^2 - 8x \\ &= (x - 4)^2 - 45 \end{aligned}$$

$$(a = -4 \text{ and } b = -45)$$

(b)

$$\begin{aligned} x^2 - 8x - 29 &= 0 \\ (x - 4)^2 - 45 &= 0 && \text{Use} \\ (x - 4)^2 &= 45 && \text{the result from part (a)} \\ x - 4 &= \pm\sqrt{45} && \text{Take} \\ x &= 4 \pm\sqrt{45} && \text{the square root of both sides.} \\ \frac{\sqrt{45}}{\sqrt{5}} = \sqrt{9} \times \sqrt{5} = 3 & && \text{Use } \sqrt{(ab)} \\ \frac{\sqrt{45}}{\sqrt{5}} &= \sqrt{a}\sqrt{b} \end{aligned}$$

$$\begin{aligned} \text{Roots are } &4 \pm 3\sqrt{5} \\ (c = 4 \text{ and } d = 3) \end{aligned}$$

Solutionbank C1

Edexcel Modular Mathematics for AS and A-Level

Algebraic fractions

Exercise A, Question 12

Question:

Given that

$$f(x) = x^2 - 6x + 18, \quad x \geq 0,$$

(a) express $f(x)$ in the form $(x - a)^2 + b$, where a and b are integers.

The curve C with equation $y = f(x)$, $x \geq 0$, meets the y -axis at P and has a minimum point at Q .

(b) Sketch the graph of C , showing the coordinates of P and Q .

The line $y = 41$ meets C at the point R .

(c) Find the x -coordinate of R , giving your answer in the form $p + q\sqrt{2}$, where p and q are integers.

Solution:

(a)

$$f(x) = x^2 - 6x + 18$$

$$x^2 - 6x = (x - 3)^2 - 9 \quad \text{for } x^2 - 6x \quad \text{Complete the square}$$

$$\begin{aligned} x^2 - 6x + 18 &= (x - 3)^2 - 9 + 18 \\ &= (x - 3)^2 + 9 \end{aligned}$$

$$(a = 3 \text{ and } b = 9)$$

(b)

$$y = x^2 - 6x + 18$$

$$y = (x - 3)^2 + 9$$

$$(x - 3)^2 \geq 0$$

Squaring a number cannot give a negative result

The minimum value of $(x - 3)^2$ is zero, when $x = 3$.


So the minimum value of y is $0 + 9 = 9$, when $x = 3$.


Q is the point $(3, 9)$


The curve crosses the y -axis where $x = 0$.

For $x = 0$, $y = 18$

P is the point $(0, 18)$

The graph of $y = x^2 - 6x + 18$ is a  shape.

For $y = ax^2 + bx + c$, if $a > 0$, the shape is 



Use the information about P and Q to sketch the curve $x \geq 0$, so the part where $x < 0$ is not needed.

(c)

$$y = (x - 3)^2 + 9$$

$$41 = (x - 3)^2 + 9$$

$$32 = (x - 3)^2$$

$$(x - 3)^2 = 32$$

$$x - 3 = \pm \sqrt{32}$$

$$x = 3 \pm \sqrt{32}$$

$$\sqrt{32} = \sqrt{16} \times \sqrt{2} = 4\sqrt{2}$$

$$x = 3 \pm 4\sqrt{2}$$

x-coordinate of R is $3 + 4\sqrt{2}$

The other value $3 - 4\sqrt{2}$ is less than 0,

so not

needed

Put $y = 41$ into the equation of C.

Subtract 9 from both sides.

Take the square root of both sides.

$$= \sqrt{a}\sqrt{b}$$

Use $\sqrt{(ab)}$

Solutionbank C1

Edexcel Modular Mathematics for AS and A-Level

Algebraic fractions

Exercise A, Question 13

Question:

Given that the equation $kx^2 + 12x + k = 0$, where k is a positive constant, has equal roots, find the value of k .

Solution:

$$Kx^2 + 12x + K = 0$$

$$a = K, b = 12, c = K$$

For equal roots, $b^2 = 4ac$
(or $b^2 - 4ac = 0$)

$$12^2$$

$$4K^2$$

$$K^2$$

$$K$$

$$\text{So } K$$

Write down the
values of a , b and c
for the quadratic
equation.

$$= 4 \times K \times K$$

$$= 144$$

$$= 36$$

$$= \pm 6$$

$$= 6$$

The question says
that K is a positive constant.

Solutionbank C1

Edexcel Modular Mathematics for AS and A-Level

Algebraic fractions

Exercise A, Question 14

Question:

Given that

$$x^2 + 10x + 36 \equiv (x + a)^2 + b,$$

where a and b are constants,

- (a) find the value of a and the value of b .
- (b) Hence show that the equation $x^2 + 10x + 36 = 0$ has no real roots.

The equation $x^2 + 10x + k = 0$ has equal roots.

- (c) Find the value of k .
- (d) For this value of k , sketch the graph of $y = x^2 + 10x + k$, showing the coordinates of any points at which the graph meets the coordinate axes.

Solution:

(a)

$$x^2 + 10x + 36$$

$$x^2 + 10x = (x + 5)^2 - 25$$

Complete the square for $x^2 + 10x$

$$x^2 + 10x + 36 = (x + 5)^2 - 25 + 36$$

$$= (x + 5)^2 + 11$$

$$a = 5 \text{ and } b = 11$$

(b)

$$x^2 + 10x + 36 = 0$$

$$(x + 5)^2 + 11 = 0 \quad \text{used}$$

'Hence' implies that part (a) must be

$$(x + 5)^2 = -11$$

A real number squared cannot

be negative, \therefore no real roots

(c)

$$x^2 + 10x + K = 0$$

$$a = 1, b = 10, c = K$$



For equal roots, $b^2 = 4ac$

$$10^2 = 4 \times 1 \times K$$

$$4K = 100$$

$$K = 25$$

(d)

The graph of $y = x^2 + 10x + 25$ is a  shape. For $y = ax^2 + bx + c$, if $a > 0$, the shape is .

$x = 0 : y = 0 + 0 + 25 = 25$

Meets y -axis at $(0, 25)$

Put $x = 0$ to find intersections with the

$y = 0 : x^2 + 10x + 25 = 0$


axis, and $y = 0$ to find intersections

$(x + 5)(x + 5) = 0$

with the x -axis.

$x = -5$

Meets x -axis at $(-5, 0)$



The graph meets the x -axis at just one point, so it 'touches' the x -axis

Solutionbank C1

Edexcel Modular Mathematics for AS and A-Level

Algebraic fractions

Exercise A, Question 15

Question:

$$x^2 + 2x + 3 \equiv (x + a)^2 + b.$$

- (a) Find the values of the constants a and b .
- (b) Sketch the graph of $y = x^2 + 2x + 3$, indicating clearly the coordinates of any intersections with the coordinate axes.
- (c) Find the value of the discriminant of $x^2 + 2x + 3$.
Explain how the sign of the discriminant relates to your sketch in part (b).

The equation $x^2 + kx + 3 = 0$, where k is a constant, has no real roots.

- (d) Find the set of possible values of k , giving your answer in surd form.

Solution:

(a)

$$x^2 + 2x + 3$$

$$x^2 + 2x = (x + 1)^2 - 1 \quad \text{for } x^2 + 2x$$


$$x^2 + 2x + 3 = (x + 1)^2 - 1 + 3$$

$$= (x + 1)^2 + 2$$


$$a = 1 \text{ and } b = 2$$

Complete the square

(b)

The graph of $y = x^2 + 2x + 3$ is a  shape

For $y = ax^2 + bx + c$,

if $a > 0$, the shape is 

$$x = 0 : y = 0 + 0 + 3$$

Meets y -axis at $(0, 3)$

Put $x = 0$ to find intersections with the y -axis,

$$y = 0 : x^2 + 2x + 3 = 0$$


$$(x + 1)^2 + 2 = 0$$

$$(x + 1)^2 = -2$$

A real number squared cannot be negative, \therefore

no real roots, so no intersection with x -axis.

and $y = 0$ to find intersections with the x -axis.



The minimum value of $(x + 1)^2$ is zero, when $x = -1$, so the minimum point on the graph is at $x = -1$

(c)

$$x^2 + 2x + 3$$

$$a = 1, b = 2, c = 3$$

$$b^2 - 4ac = 2^2 - 4 \times 1 \times 3 = -8$$

Since the discriminant is negative

$$(b^2 - 4ac < 0), x^2 + 2x + 3 = 0$$

has no real roots, so the graph

does not cross the x -axis.

$$\text{real roots: } b^2 < 4ac$$

The discriminant is $b^2 - 4ac$

No

(d)

$$x^2 + kx + 3 = 0$$


$$a = 1, b = k, c = 3$$

For no real roots, $b^2 < 4ac$

$$k^2 < 12$$

$$k^2 - 12 < 0$$

This is a quadratic inequality with critical values $-\sqrt{12}$ and $\sqrt{12}$



$$(k + \sqrt{12})(k - \sqrt{12}) < 0$$

Critical values:

$$K = -\sqrt{12}, K = \sqrt{12}$$

$$-\sqrt{12} < K < \sqrt{12}$$

$$\left(\begin{array}{l} \sqrt{12} = \sqrt{4} \times \sqrt{3} = 2\sqrt{3} \\ -2\sqrt{3} < K < 2\sqrt{3} \end{array} \right)$$

$$\sqrt{(ab)} = \sqrt{a}\sqrt{b}$$

The surds can be simplified using

Solutionbank C1

Edexcel Modular Mathematics for AS and A-Level

Algebraic fractions

Exercise A, Question 16

Question:

Solve the simultaneous equations

$$\begin{aligned}x + y &= 2 \\x^2 + 2y &= 12\end{aligned}$$

Solution:

$y = 2 - x$		Rearrange the linear equation to get $y = \dots$
$x^2 + 2(2 - x)$	$= 12$	Substitute into the quadratic equation
$x^2 + 4 - 2x$	$= 12$	
$x^2 - 2x + 4 - 12$	$= 0$	
$x^2 - 2x - 8$	$= 0$	
$(x + 2)(x - 4)$	$= 0$	Solve for x using factorisation
$x = -2$ or $x = 4$		
$x = -2 : y = 2 -$ $(-2) = 4$		Substitute the x values back into $y = 2 - x$
$x = 4 : y = 2 - 4 = -2$		
Solution: $x = -2, y = 4$ and $x = 4, y = -2$		

Solutionbank C1

Edexcel Modular Mathematics for AS and A-Level

Algebraic fractions

Exercise A, Question 17

Question:

(a) By eliminating y from the equations

$$\begin{aligned}y &= x - 4, \\ 2x^2 - xy &= 8,\end{aligned}$$

show that

$$x^2 + 4x - 8 = 0.$$

(b) Hence, or otherwise, solve the simultaneous equations

$$\begin{aligned}y &= x - 4, \\ 2x^2 - xy &= 8,\end{aligned}$$

giving your answers in the form $a \pm b\sqrt{3}$, where a and b are integers.

Solution:

(a)

$$\frac{2x^2 - x}{(x - 4)} = 8 \text{ equation.}$$

$$2x^2 - x^2 + 4x = 8$$

$$x^2 + 4x - 8 = 0$$

Substitute $y = x - 4$ into the quadratic

(b)

(a). The $\sqrt{3}$
factorisation
quadratic

Solve the equation found in part
in the given answer suggests that
will not be possible, so use the
formula, or complete the square.

$$\begin{aligned}
 x^2 + 4x - 8 &= 0 \\
 x^2 + 4x &= (x + 2)^2 - 4 \\
 (x + 2)^2 - 4 - 8 &= 0 \\
 (x + 2)^2 &= 12 \\
 x + 2 &= \pm \sqrt{12} \\
 x &= -2 \pm \sqrt{12} \\
 \sqrt{12} &= \sqrt{4 \times 3} = 2\sqrt{3} \\
 x &= -2 \pm 2\sqrt{3} \\
 (a = -2 \text{ and } b = 2) \\
 \text{Using } y &= x - 4, \\
 y &= (-2 \pm 2\sqrt{3}) - 4 \\
 &= -6 \pm 2\sqrt{3} \\
 \text{Solution: } x &= -2 \pm 2\sqrt{3} \\
 y &= -6 \pm 2\sqrt{3}
 \end{aligned}$$

Complete the square for $x^2 + 4x$

$$\text{Use } \sqrt{(ab)} = \sqrt{a}\sqrt{b}$$

Solutionbank C1

Edexcel Modular Mathematics for AS and A-Level

Algebraic fractions

Exercise A, Question 18

Question:

Solve the simultaneous equations

$$\begin{aligned} 2x - y - 5 &= 0 \\ x^2 + xy - 2 &= 0 \end{aligned}$$

Solution:

$$y = 2x - 5$$

the linear

Rearrange

equation to

get $y = \dots$

$$x^2 + x(2x - 5) - 2 = 0$$

Substitute
into the quadratic equation.

$$x^2 + 2x^2 - 5x - 2 = 0$$

$$3x^2 - 5x - 2 = 0$$

$$(3x + 1)(x - 2) = 0$$

Solve
for x using factorisation

$$x = -\frac{1}{3} \text{ or } x = 2$$

$$x = -\frac{1}{3} : y = -$$

Substitute

$$\frac{2}{3} - 5 = -\frac{17}{3} \quad \text{the } x \text{ values}$$

$$x = 2 : y = 4 - 5 = -1 \quad \text{into } y = 2x - 5$$

back

Solution $x = -$

$$\frac{1}{3}, y = -\frac{17}{3}$$

$$\text{and } x = 2, y = -1$$

Solutionbank C1

Edexcel Modular Mathematics for AS and A-Level

Algebraic fractions

Exercise A, Question 19

Question:

Find the set of values of x for which

(a) $3(2x + 1) > 5 - 2x$,

(b) $2x^2 - 7x + 3 > 0$,

(c) both $3(2x + 1) > 5 - 2x$ and $2x^2 - 7x + 3 > 0$.

Solution:

(a)

$$3(2x + 1) > 5 - 2x$$

$$6x + 3 > 5 - 2x$$

$$6x + 2x + 3 > 5$$

$$8x > 2$$

$$x > \frac{1}{4}$$

Multiply out

Add $2x$ to both sides.

Subtract 3 from both sides

Divide both sides by 8


(b)

$$2x^2 - 7x + 3 = 0$$

$$\begin{aligned} (2x - 1) \\ (x - 3) \end{aligned} = 0 \text{ quadratic equation.}$$

$$x = \frac{1}{2}, x = 3$$

Factorise to solve the



Sketch the graph of $y = 2x^2 - 7x + 3$. The

shape is  The sketch does not need to be accurate.


$2x^2 - 7x + 3 > 0$ where the part

$$x < \frac{1}{2} \text{ or } x > 3$$

$$2x^2 - 7x + 3 > 0 \quad (y > 0) \text{ for}$$

of the graph above the x -axis

(c)



$$\frac{1}{4} < x <$$

$$\frac{1}{2}, x > 3$$

(a)

Use a number line. The

two sets of values (from part

and part (b)) overlap for

$$\frac{1}{4} < x < \frac{1}{2} \text{ and } x > 3$$

Solutionbank C1

Edexcel Modular Mathematics for AS and A-Level

Algebraic fractions

Exercise A, Question 20

Question:

Find the set of values of x for which


(a) $x(x - 5) < 7x - x^2$

(b) $x(3x + 7) > 20$

Solution:

(a)
 $x(x - 5) < 7x - x^2$
 $x^2 - 5x < 7x - x^2$
 $2x^2 - 12x < 0$
 $2x(x - 6) < 0$

$2x(x - 6) = 0$
 $x = 0, x = 6$



$2x^2 - 12x < 0$ where
 $0 < x < 6$

(b)
 $x(3x + 7) > 20$
 $3x^2 + 7x > 20$
 $3x^2 + 7x - 20 > 0$
 $(3x - 5)(x + 4) > 0$
 $(3x - 5)(x + 4) = 0$
 $x = \frac{5}{3}, x = -4$

Multiply out

Factorise using the common factor $2x$

Solve the quadratic equation to find the critical values


Sketch the graph of
 $y = 2x^2 - 12x$

$2x^2 - 12x < 0$ ($y < 0$)
 for the part of the graph below the x -axis

Multiply out

Factorise

Solve the quadratic equation to find the critical values



$$3x^2 + 7x - 20 > 0 \text{ where}$$
$$x < -4 \text{ or } x > \frac{5}{3}$$

Sketch the graph of
 $y = 3x^2 + 7x - 20$

$$3x^2 + 7x - 20 > 0 \quad (y > 0)$$

for the part of the graph

above the x -axis.

Solutionbank C1

Edexcel Modular Mathematics for AS and A-Level

Algebraic fractions

Exercise A, Question 21

Question:

(a) Solve the simultaneous equations

$$\begin{aligned}y + 2x &= 5, \\ 2x^2 - 3x - y &= 16.\end{aligned}$$

(b) Hence, or otherwise, find the set of values of x for which

$$2x^2 - 3x - 16 > 5 - 2x.$$

Solution:

(a)

$$y = 5 - 2x$$

the linear equation

Rearrange

to

$$\text{get } y = \dots$$

$$2x^2 - 3x - (5 - 2x) = 16$$

Substitute
into the quadratic equation.

$$2x^2 - 3x - 5 + 2x = 16$$

$$2x^2 - x - 21 = 0$$

$$(2x - 7)(x + 3) = 0$$

Solve
for x using factorisation.

$$x = 3\frac{1}{2} \text{ or } x = -3$$

$$x = 3$$

Substitute

$$\frac{1}{2} : y = 5 - 7 = -2 \quad \text{the } x\text{-values back into}$$

$$x = -3 : y = 5 + 6 = 11$$

$$y = 5 - 2x$$

Solution $x = 3$

$$\frac{1}{2}, y = -2$$

$$\text{and } x = -3, y = 11$$

(b)

The equations in (a) could be written as

$$y = 5 - 2x \text{ and } y = 2x^2 - 3x - 16.$$

The solutions to $2x^2 - 3x - 16 = 5 - 2x$ are the x solutions from (a). These are the critical values for $2x^2 - 3x - 16 > 5 - 2x$.


Critical values

$$x = 3\frac{1}{2} \text{ and } x = -3.$$

$$2x^2 - 3x - 16 > 5 - 2x$$

$$(2x^2 - 3x - 16 - 5 + 2x > 0)$$

$$2x^2 - x - 21 > 0$$



$$x < -3 \text{ or } x > 3\frac{1}{2}$$

Sketch the graph of
 $y = 2x^2 - x - 21$

$2x^2 - x - 21 > 0$ ($y > 0$) for the
part of the graph above the x -axis.

Solutionbank C1

Edexcel Modular Mathematics for AS and A-Level

Algebraic fractions

Exercise A, Question 22

Question:

The equation $x^2 + kx + (k + 3) = 0$, where k is a constant, has different real roots.

(a) Show that $k^2 - 4k - 12 > 0$.

(b) Find the set of possible values of k .

Solution:

(a)

$$x^2 + kx + (k + 3) = 0$$

$$a = 1, b = k, c = k + 3$$

$$b^2 > 4ac$$

$$k^2 > 4(k + 3)$$

$$k^2 > 4k + 12$$

$$k^2 - 4k - 12 > 0$$

Write down a , b and c for the equation

For different real roots, $b^2 > 4ac$


(b)

$$k^2 - 4k - 12 = 0 \text{ equation.}$$


$$\begin{aligned} (k + 2) \\ (k - 6) \end{aligned} = 0$$

$$k = -2, k = 6$$

Factorise to solve the quadratic



Sketch the graph

The shape is  The sketch does not need to be accurate

$$\begin{aligned} k^2 - 4k - 12 > 0 \text{ where} \\ k < -2 \text{ or } k > 6 \end{aligned}$$

$k^2 - 4k - 12 > 0$ ($y > 0$) for the part of the graph above the k -axis.

Solutionbank C1

Edexcel Modular Mathematics for AS and A-Level

Algebraic fractions

Exercise A, Question 23

Question:

Given that the equation $kx^2 + 3kx + 2 = 0$, where k is a constant, has no real roots, find the set of possible values of k .

Solution:

$$kx^2 + 3kx + 2 = 0$$

$a = k$, $b = 3k$, $c = 2$ Write down a , b and c for the equation.

$$b^2 < 4ac$$

$$(3k)^2 < 4 \times k \times 2 \quad \text{no real roots, } b^2 < 4ac.$$


$$9k^2 < 8k$$

$$9k^2 - 8k < 0$$

$$9k^2 - 8k = 0$$

$$k(9k - 8) = 0$$

$$k = 0, k = \frac{8}{9}$$



$$9k^2 - 8k < 0 \quad \text{where}$$


$$0 < k < \frac{8}{9}$$

For

to solve the quadratic equation

Factorise

Sketch the graph of $y = 9k^2 - 8k$. The shape is



. The sketch does not need to be accurate.

$9k^2 - 8k < 0$ ($y < 0$) for the part of the graph below the k -axis.

Solutionbank C1

Edexcel Modular Mathematics for AS and A-Level

Algebraic fractions

Exercise A, Question 24

Question:

The equation $(2p + 5)x^2 + px + 1 = 0$, where p is a constant, has different real roots.

- (a) Show that $p^2 - 8p - 20 > 0$
- (b) Find the set of possible values of p .

Given that $p = -3$,

- (c) find the exact roots of $(2p + 5)x^2 + px + 1 = 0$.

Solution:

(a)

$$(2p + 5)x^2 + px + 1 = 0$$

$$a = 2p + 5, b = p, c = 1$$

$$b^2 > 4ac$$

$$p^2 > 4(2p + 5)$$

$$p^2 > 8p + 20$$

$$p^2 - 8p - 20 > 0$$

Write down a , b and c for the equation.

For different real roots, $b^2 > 4ac$


(b)

$$p^2 - 8p - 20 = 0$$

$$\begin{aligned} (p + 2) \\ (p - 10) \end{aligned} = 0 \text{ equation.}$$


$$p = -2, p = 10$$

Factorise to solve the quadratic



Sketch the graph of

$$y = p^2 - 8p - 20$$

The shape is . The sketch does not need to be accurate

$$p^2 - 8p - 20 > 0 \text{ where } p < -2 \text{ or } p > 10$$

$p^2 - 8p - 20 > 0$ ($y > 0$) for the part of the graph above the p -axis

(c)

For $p = -3$

$$(-6 + 5)x^2 - 3x + 1 = 0$$

$$-x^2 - 3x + 1 = 0$$

$$x^2 + 3x - 1 = 0$$

$$a = 1, b = 3, c = -1$$

$$x = \frac{-b \pm \sqrt{(b^2 - 4ac)}}{2a}$$

$$x = \frac{-3 \pm \sqrt{9 + 4}}{2}$$

$$x = \frac{1}{2} (-3 \pm \sqrt{13})$$

$\sqrt{13}$ cannot be simplified.

$$x = \frac{1}{2} (-3 + \sqrt{13}) \quad \text{or} \quad x =$$

$$\frac{1}{2} (-3 - \sqrt{13})$$

Substitute $p = -3$ into the equation.

Multiply by -1

The equation does not factorise,

so use the quadratics formula.

Quote the formula.

Exact roots are required.

Solutionbank C1

Edexcel Modular Mathematics for AS and A-Level

Algebraic fractions

Exercise A, Question 25

Question:

(a) Factorise completely $x^3 - 4x$

(b) Sketch the curve with equation $y = x^3 - 4x$, showing the coordinates of the points where the curve crosses the x -axis.

(c) On a separate diagram, sketch the curve with equation
 $y = (x - 1)^3 - 4(x - 1)$
 showing the coordinates of the points where the curve crosses the x -axis.

Solution:

(a)

$$x^3 - 4x$$

$$= x(x^2 - 4)$$

squares

$$= x(x + 2)$$

$$(x - 2)$$

x is a common factor

$(x^2 - 4)$ is a difference of

(b)

Curve crosses x -axis where $y = 0$

$$x(x + 2)(x - 2) = 0$$

$$x = 0, x = -2, x = 2$$

When $x = 0$, $y = 0$

curve crosses

the y -axis.

When $x \rightarrow \infty$, $y \rightarrow \infty$

large

When $x \rightarrow -\infty$, $y \rightarrow -\infty$


of x

Put $y = 0$ and solve for x

Put $x = 0$ to find where the

Check what happens to y for

positive and negative values



Crosses at $(0, 0)$

Crosses x -axis at $(-2, 0)$, $(2, 0)$.

(c)

$$y = x^3 - 4x$$

$$y = (x - 1)^3 - 4(x - 1)$$

This is a translation of $+1$ in the x -direction.


(b).

Compare with the equation from part

x has been replaced by $x - 1$.

$f(x + a)$ is a translation of

$-a$ in the x -direction.



Crosses x -axis at $(-1, 0)$, $(1, 0)$, $(3, 0)$

The shape is the same as in part (b).


Solutionbank C1

Edexcel Modular Mathematics for AS and A-Level

Algebraic fractions

Exercise A, Question 26

Question:



The figure shows a sketch of the curve with equation $y = f(x)$. The curve crosses the x -axis at the points $(2, 0)$ and $(4, 0)$. The minimum point on the curve is $P(3, -2)$.

In separate diagrams, sketch the curve with equation


(a) $y = -f(x)$

(b) $y = f(2x)$

On each diagram, give the coordinates of the points at which the curve crosses the x -axis, and the coordinates of the image of P under the given transformation.

Solution:

(a)




The transformation $-f(x)$ multiplies the y -coordinates by -1 . This turns the graph upside-down.

Crosses the x -axis at $(2, 0)$, $(4, 0)$

Image of P is $(3, 2)$

(b)



Crosses the x -axis at $(1, 0)$,
 $(2, 0)$

Image of P is $(1\frac{1}{2}, -2)$

unchanged.

$f(2x)$ is a stretch of $\frac{1}{2}$
in the x -direction. (Multiply

x -coordinates by $\frac{1}{2}$.)


y -coordinates are

Solutionbank C1

Edexcel Modular Mathematics for AS and A-Level

Algebraic fractions
Exercise A, Question 27

Question:



The figure shows a sketch of the curve with equation $y = f(x)$. The curve passes through the points $(0, 3)$ and $(4, 0)$ and touches the x -axis at the point $(1, 0)$.

On separate diagrams, sketch the curve with equation

(a) $y = f(x + 1)$

(b) $y = 2f(x)$


(c) $y = f\left(\frac{1}{2}x\right)$

On each diagram, show clearly the coordinates of all the points where the curve meets the axes.

Solution:

(a)

$f(x + 1)$ is a translation of




Meets the x -axis at $(0, 0)$, $(3, 0)$

Meets the y -axis at $(0, 0)$

-1 in the x -direction.

(b)



Meets the x -axis at $(1, 0)$,
 $(4, 0)$


Meets the y -axis at $(0, 6)$

unchanged.

$2f(x)$ is a stretch of scale
factor 2 in the y -direction
(Multiply y -coordinates by 2)

x -coordinates are

(c)



$f\left(\frac{1}{2}x\right)$ is a stretch of scale

factor $\frac{1}{\left(\frac{1}{2}\right)} = 2$ in the

x -direction. (Multiply
 x -coordinates by 2)

Meets the x -axis at $(2, 0)$,
 $(8, 0)$

Meets the y -axis at $(0, 3)$ unchanged.

y -coordinates are

Solutionbank C1

Edexcel Modular Mathematics for AS and A-Level

Algebraic fractions

Exercise A, Question 28


Question:

Given that $f(x) = \frac{1}{x}$, $x \neq 0$,

- (a) sketch the graph of $y = f(x) + 3$ and state the equations of the asymptotes.
 (b) Find the coordinates of the point where $y = f(x) + 3$ crosses a coordinate axis.

Solution:

(a)



You should know the shape of this curve.

$f(x) + 3$ is a translation of $+3$ in the y -direction.

$y = 3$ is an asymptote
 $x = 0$ is an asymptote

is $x = 0$

The equation of the y -axis

(b)

The graph does not cross

get

the y -axis (see sketch in
(a)) .

undefined ,

Crosses the x -axis where $y = 0$:

$$\frac{1}{x} + 3 = 0$$

$$\frac{1}{x} = -3$$

$$x = -\frac{1}{3} \quad \left(-\frac{1}{3}, 0 \right)$$

If you used $x = 0$ you would

$$y = \frac{1}{0} + 3 \text{ but } \frac{1}{0} \text{ is}$$

or infinite.

Solutionbank C1

Edexcel Modular Mathematics for AS and A-Level

Algebraic fractions

Exercise A, Question 29

Question:

Given that $f(x) = (x^2 - 6x)(x - 2) + 3x$,

(a) express $f(x)$ in the form $x(ax^2 + bx + c)$, where a , b and c are constants

(b) hence factorise $f(x)$ completely

(c) sketch the graph of $y = f(x)$, showing the coordinates of each point at which the graph meets the axes

Solution:

(a)

$$\begin{aligned}
 f(x) &= (x^2 - 6x)(x - 2) + 3x && \text{Multiply} \\
 &= x^2(x - 2) - 6x(x - 2) && \text{out the bracket} \\
 &\quad + 3x \\
 &= x^3 - 2x^2 - 6x^2 + 12x + 3x \\
 &= x^3 - 8x^2 + 15x && \text{common factor} \\
 &= x(x^2 - 8x + 15) && x \text{ is a} \\
 (a = 1, b = -8, c = 15)
 \end{aligned}$$

(b)

$$\begin{aligned}
 &x(x^2 - 8x + 15) && \text{Factorise the quadratic} \\
 f(x) &= x(x - 3)(x - 5)
 \end{aligned}$$

(c)

Curve meets x -axis
where $y = 0$.

$$x(x - 3)(x - 5) = 0$$

$$x = 0, x = 3, x = 5$$

When $x = 0$, $y = 0$

Put $y = 0$ and solve for x


Put $x = 0$ to find where the curve crosses the y -axis

Check what happens to y for

When $x \rightarrow \infty$, $y \rightarrow \infty$ large

When $x \rightarrow -\infty$, $y \rightarrow -\infty$ of x .

positive and negative values



Meets x -axis at $(0, 0)$, $(3, 0)$, $(5, 0)$

Meets y -axis at $(0, 0)$

© Pearson Education Ltd 2008

Solutionbank C1

Edexcel Modular Mathematics for AS and A-Level

Algebraic fractions Exercise A, Question 30

Question:

- (a) Sketch on the same diagram the graph of $y = x(x + 2)(x - 4)$ and the graph of $y = 3x - x^2$, showing the coordinates of the points at which each graph meets the x -axis.
 (b) Find the exact coordinates of each of the intersection points of $y = x(x + 2)(x - 4)$ and $y = 3x - x^2$.

Solution:

(a)
 $y = x(x + 2)(x - 4)$
 Curve meets x -axis where $y = 0$.
 $x(x + 2)(x - 4) = 0$
 $x = 0, x = -2, x = 4$
 When $x = 0, y = 0$


Put $y = 0$ and solve for x .


Put $x = 0$ to find where the curve crosses the y -axis

When $x \rightarrow \infty, y \rightarrow \infty$
 When $x \rightarrow -\infty, y \rightarrow -\infty$

Check what happens to y for large positive and negative values of x .

$$y = 3x - x^2$$

The graph of $y = 3x - x^2$ is a  shape

For $y = ax^2 + bx + c$,
 if $a < 0$, the shape is 

$$3x - x^2 = 0$$

$$x(3 - x) = 0$$


$$x = 0, x = 3$$

Put $y = 0$ and solve for x .

When $x = 0, y = 0$

Put $x = 0$ to find where the curve crosses the y -axis.

the y -axis.



$y = x(x + 2)(x - 4)$ meets the x -axis at $(-2, 0), (0, 0), (4, 0)$
 $y = 3x - x^2$ meets the x -axis at $(0, 0), (3, 0)$

(b)

$$x(x + 2)(x - 4) = 3x - x^2$$

$$x(x + 2)(x - 4) = x(3 - x)$$

$$(x + 2)(x - 4) = 3 - x$$

One solution is $x = 0$

$$x^2 - 2x - 8 = 3 - x$$

$$x^2 - 2x + x - 8 - 3 = 0$$

$$x^2 - x - 11 = 0$$

to give an equation in x .
 $x = 0$ is a solution.

To find where the graphs intersect, equate the two expressions for y

If you divide by x , remember that

$$a = 1, b = -1, c = -11$$

use the quadratic formula.

The equation does not factorise, so

$$x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$$

Quote the formula

$$x = \frac{1 \pm \sqrt{(-1)^2 - (4 \times 1 \times -11)}}{2}$$

$$= \frac{1 \pm \sqrt{45}}{2}$$

Exact values are required, not rounded decimals, so leave the answers in surd form.

$$\sqrt{45} = \sqrt{9 \times 5} = 3\sqrt{5}$$

$$x = \frac{1}{2} (1 \pm 3\sqrt{5})$$

$$x = \frac{1}{2} (1 + 3\sqrt{5}) \text{ or } x = \frac{1}{2} (1 - 3\sqrt{5})$$

$x = 0 : y = 0$ The y-coordinates for the intersection

$$x = \frac{1}{2} (1 + 3\sqrt{5})$$

points are also needed.

$$y = \frac{3(1 + 3\sqrt{5})}{2} - \frac{(1 + 3\sqrt{5})^2}{4}$$

Use $y = 3x - x^2$, the simpler equation

$$\begin{aligned} (1 + 3\sqrt{5})^2 &= (1 + 3\sqrt{5})(1 + 3\sqrt{5}) \\ &= 1(1 + 3\sqrt{5}) + 3\sqrt{5}(1 + 3\sqrt{5}) \\ &= 1 + 3\sqrt{5} + 3\sqrt{5} + 45 \\ &= 46 + 6\sqrt{5} \end{aligned}$$

$$\sqrt{5} \times \sqrt{5} = 5$$

$$y = \frac{6(1 + 3\sqrt{5})}{4} - \frac{46 + 6\sqrt{5}}{4}$$

Use a common denominator 4.

$$= \frac{6 + 18\sqrt{5} - 46 - 6\sqrt{5}}{4}$$

$$= \frac{-40 + 12\sqrt{5}}{4} = -10 + 3\sqrt{5}$$

$$x = \frac{1}{2} (1 - 3\sqrt{5})$$

$$y = \frac{3(1 - 3\sqrt{5})}{2} - \frac{(1 - 3\sqrt{5})^2}{4}$$

$$y = \frac{6(1 - 3\sqrt{5})}{4} - \frac{46 - 6\sqrt{5}}{4}$$

that for The working will be similar to

repeated. $1 + 3\sqrt{5}$, so need not be fully

$$= \frac{6 - 18\sqrt{5} - 46 + 6\sqrt{5}}{4}$$

$$= \frac{-40 - 12\sqrt{5}}{4} = -10 - 3\sqrt{5}$$

$\sqrt{5}$

Intersection points are :

Finally, write down the coordinates of all the points you have found. You can compare

$$(0, 0), \left(\frac{1}{2}(1 + 3\sqrt{5}), -10 + 3\sqrt{5}\right)$$

these with your sketch, as a rough check.

$$\text{and } \left(\frac{1}{2}(1 - 3\sqrt{5}), -10 - 3\sqrt{5}\right)$$