

## C1

## DIFFERENTIATION

## Answers - Worksheet A

1 a

$x$ -coordinate of $B$	$y$ -coordinate of $B$	gradient of $AB$
2	4	$\frac{4-1}{2-1} = 3$
1.1	1.21	$\frac{1.21-1}{1.1-1} = 2.1$
1.01	1.0201	$\frac{1.0201-1}{1.01-1} = 2.01$
1.001	1.002001	$\frac{1.002001-1}{1.001-1} = 2.001$

b gradient = 2

c

$x$ -coordinate of $B$	$y$ -coordinate of $B$	gradient of $AB$
0	0	$\frac{1-0}{1-0} = 1$
0.9	0.81	$\frac{1-0.81}{1-0.9} = 1.9$
0.99	0.9801	$\frac{1-0.9801}{1-0.99} = 1.99$
0.999	0.998001	$\frac{1-0.998001}{1-0.999} = 1.999$

this table supports the answer to part **b** as the gradient of the chord  $AB$  again gets closer to 2 as  $B$  gets closer to  $A$

2 possible tables of values are:

**a**

$x$ -coordinate of $B$	$y$ -coordinate of $B$	gradient of $AB$
3	9	$\frac{9-4}{3-2} = 5$
2.1	4.41	$\frac{4.41-4}{2.1-2} = 4.1$
2.01	4.0401	$\frac{4.0401-4}{2.01-2} = 4.01$
2.001	4.004001	$\frac{4.004001-4}{2.001-2} = 4.001$

$\therefore$  gradient = 4

**b**

$x$ -coordinate of $B$	$y$ -coordinate of $B$	gradient of $AB$
5	25	$\frac{25-16}{5-4} = 9$
4.1	16.81	$\frac{16.81-16}{4.1-4} = 8.1$
4.01	16.0801	$\frac{16.0801-16}{4.01-4} = 8.01$
4.001	16.008001	$\frac{16.008001-16}{4.001-4} = 8.001$

$\therefore$  gradient = 8

**c**

$x$ -coordinate of $B$	$y$ -coordinate of $B$	gradient of $AB$
2.5	6.25	$\frac{6.25-2.25}{2.5-1.5} = 4$
1.6	2.56	$\frac{2.56-2.25}{1.6-1.5} = 3.1$
1.51	2.2801	$\frac{2.2801-2.25}{1.51-1.5} = 3.01$
1.501	2.253001	$\frac{2.253001-2.25}{1.501-1.5} = 3.001$

$\therefore$  gradient = 3

**d**

$x$ -coordinate of $B$	$y$ -coordinate of $B$	gradient of $AB$
-2	4	$\frac{4-9}{-2-(-3)} = -5$
-2.9	8.41	$\frac{8.41-9}{-2.9-(-3)} = -5.9$
-2.99	8.9401	$\frac{8.9401-9}{-2.99-(-3)} = -5.99$
-2.999	8.994001	$\frac{8.994001-9}{-2.999-(-3)} = -5.999$

$\therefore$  gradient = -6

3 **a** gradient =  $2x$

**b** **i** 12      **ii** 4.8      **iii** -6.4

4 possible answers are:

a let  $A$  be  $(1, 1)$

$x$ -coordinate of $B$	$y$ -coordinate of $B$	gradient of $AB$
2	16	$\frac{16-1}{2-1} = 15$
1.1	1.4641	$\frac{1.4641-1}{1.1-1} = 4.641$
1.01	1.04060401	$\frac{1.04060401-1}{1.01-1} = 4.060401$
1.001	1.004006004	$\frac{1.004006004-1}{1.001-1} = 4.006004$

$\therefore$  gradient = 4

b let  $A$  be  $(2, -3)$

$x$ -coordinate of $B$	$y$ -coordinate of $B$	gradient of $AB$
3	-3	$\frac{-3-(-3)}{3-2} = 0$
2.1	-3.09	$\frac{-3.09-(-3)}{2.1-2} = -0.9$
2.01	-3.0099	$\frac{-3.0099-(-3)}{2.01-2} = -0.99$
2.001	-3.000999	$\frac{-3.000999-(-3)}{2.001-2} = -0.999$

$\therefore$  gradient = -1

c let  $A$  be  $(4, 2)$

$x$ -coordinate of $B$	$y$ -coordinate of $B$	gradient of $AB$
5	2.236067977	$\frac{2.236067977-2}{5-4} = 0.236068$
4.1	2.024845673	$\frac{2.024845673-2}{4.1-4} = 0.248457$
4.01	2.002498439	$\frac{2.002498439-2}{4.01-4} = 0.249844$
4.001	2.000249984	$\frac{2.000249984-2}{4.001-4} = 0.249984$

$\therefore$  gradient = 0.25

d let  $A$  be  $(2, 1)$

$x$ -coordinate of $B$	$y$ -coordinate of $B$	gradient of $AB$
3	0.666666667	$\frac{0.666666667-1}{3-2} = -0.333333$
2.1	0.952380952	$\frac{0.952380952-1}{2.1-2} = -0.476190$
2.01	0.995024876	$\frac{0.995024876-1}{2.01-2} = -0.497512$
2.001	0.999500250	$\frac{0.999500250-1}{2.001-2} = -0.499750$

$\therefore$  gradient = -0.5

5 a possible answers are:

i let  $A$  be  $(1, 1)$

$x$ -coordinate of $B$	$y$ -coordinate of $B$	gradient of $AB$
2	8	$\frac{8-1}{2-1} = 7$
1.1	1.331	$\frac{1.331-1}{1.1-1} = 3.31$
1.01	1.030301	$\frac{1.030301-1}{1.01-1} = 3.0301$
1.001	1.003003001	$\frac{1.003003001-1}{1.001-1} = 3.003001$

$\therefore$  gradient = 3

ii let  $A$  be  $(2, 8)$

$x$ -coordinate of $B$	$y$ -coordinate of $B$	gradient of $AB$
3	27	$\frac{27-8}{3-2} = 19$
2.1	9.261	$\frac{9.261-8}{2.1-2} = 12.61$
2.01	8.120601	$\frac{8.120601-8}{2.01-2} = 12.0601$
2.001	8.012006001	$\frac{8.012006001-8}{2.001-2} = 12.006001$

$\therefore$  gradient = 12

iii let  $A$  be  $(3, 27)$

$x$ -coordinate of $B$	$y$ -coordinate of $B$	gradient of $AB$
4	64	$\frac{64-27}{4-3} = 37$
3.1	29.791	$\frac{29.791-27}{3.1-3} = 27.91$
3.01	27.270901	$\frac{27.270901-27}{3.01-3} = 27.0901$
3.001	27.027009	$\frac{27.027009-27}{3.001-3} = 27.009$

$\therefore$  gradient = 27

b gradient =  $3x^2$

c i 48      ii 12      iii 6.75

**C1 DIFFERENTIATION**
**Answers - Worksheet B**

- 1**    **a**  $2x$             **b**  $4x^3$             **c**  $1$             **d**  $9x^8$             **e**  $-3x^{-4}$             **f**  $-x^{-2}$   
          **g**  $8x$             **h**  $7$             **i**  $10x^4$             **j**  $0$             **k**  $-16x^{-3}$             **l**  $-44x^{-5}$
- 2**    **a**  $5x^4 + 2x$             **b**  $1 + 3x^2$             **c**  $4x^3$             **d**  $6x^5 - 2$   
          **e**  $18x^2 - 10x^{-3}$             **f**  $2x - 4$             **g**  $-x^{-2} + 5x^{-6}$             **h**  $12x^2 - 12x^{-5}$
- 3**    **a**  $6t^5$             **b**  $-15t^{-4}$             **c**  $\frac{1}{2}t^{-\frac{1}{2}}$             **d**  $\frac{2}{3}t^{-\frac{1}{3}}$             **e**  $\frac{3}{2}t$             **f**  $2t^{-\frac{3}{4}}$   
          **g**  $7t^{\frac{5}{2}}$             **h**  $-\frac{1}{5}t^{-\frac{6}{5}}$             **i**  $\frac{3}{5}t^{\frac{1}{5}}$             **j**  $-\frac{3}{2}t^{-\frac{5}{2}}$             **k**  $-15t^{-\frac{9}{4}}$             **l**  $\frac{2}{9}t^{\frac{1}{3}}$
- 4**    **a**  $2 + 2x^5$             **b**  $\frac{3}{2}x^{\frac{1}{2}}$             **c**  $1 + 2x^{-\frac{1}{2}}$             **d**  $10x^{\frac{2}{3}} + 4x^{-5}$   
          **e**  $-\frac{4}{5}x^{-\frac{9}{5}}$             **f**  $\frac{1}{3}x^{-\frac{5}{6}} + \frac{3}{4}x^{-\frac{1}{4}}$             **g**  $-3x^{-2} + \frac{15}{2}x^{-\frac{5}{2}}$             **h**  $7x^{-2} - \frac{8}{3}x^{-\frac{11}{3}}$
- 5**    **a**  $y = x^{\frac{1}{2}}$             **b**  $y = 4 - x^{-1}$             **c**  $y = 3x^2 + x^{\frac{1}{3}}$             **d**  $y = 9x + 3x^{-1}$   
           $\frac{dy}{dx} = \frac{1}{2}x^{-\frac{1}{2}}$              $\frac{dy}{dx} = x^{-2}$              $\frac{dy}{dx} = 6x + \frac{1}{3}x^{-\frac{2}{3}}$              $\frac{dy}{dx} = 9 - 3x^{-2}$   
          **e**  $y = \frac{1}{4}x^{-1} - x^{-2}$             **f**  $y = 6x^{-\frac{1}{4}}$             **g**  $y = x^{\frac{5}{2}}$             **h**  $y = 8x^{\frac{1}{2}} + \frac{4}{3}x^{-2}$   
           $\frac{dy}{dx} = -\frac{1}{4}x^{-2} + 2x^{-3}$              $\frac{dy}{dx} = -\frac{3}{2}x^{-\frac{5}{4}}$              $\frac{dy}{dx} = \frac{5}{2}x^{\frac{3}{2}}$              $\frac{dy}{dx} = 4x^{-\frac{1}{2}} - \frac{8}{3}x^{-3}$
- 6**    **a**  $s = t^2 + 3t$             **b**  $s = t^2 - 4t + 4$             **c**  $s = 5t^4 + 20t^2$             **d**  $s = 7t^3 - t$   
           $\frac{ds}{dt} = 2t + 3$              $\frac{ds}{dt} = 2t - 4$              $\frac{ds}{dt} = 20t^3 + 40t$              $\frac{ds}{dt} = 21t^2 - 1$   
          **e**  $s = t^2 + 7t + 6$             **f**  $s = t^2 - 2t - 8$             **g**  $s = t^5 + 3t^3 + 9t$             **h**  $s = 2t^3 - 5t^2 + 3t$   
           $\frac{ds}{dt} = 2t + 7$              $\frac{ds}{dt} = 2t - 2$              $\frac{ds}{dt} = 5t^4 + 9t^2 + 9$              $\frac{ds}{dt} = 6t^2 - 10t + 3$
- 7**    **a**  $y = x^{\frac{3}{2}} - 4x^{\frac{1}{2}}$             **b**  $y = x^2 - 2$             **c**  $y = 4x + x^{-1}$             **d**  $y = x^{\frac{1}{2}} + 3x^{-\frac{1}{2}}$   
           $\frac{dy}{dx} = \frac{3}{2}x^{\frac{1}{2}} - 2x^{-\frac{1}{2}}$              $\frac{dy}{dx} = 2x$              $\frac{dy}{dx} = 4 - x^{-2}$              $\frac{dy}{dx} = \frac{1}{2}x^{-\frac{1}{2}} - \frac{3}{2}x^{-\frac{3}{2}}$   
          **e**  $y = 2x^{-1} - \frac{1}{2}x^2$             **f**  $y = 5x^{-2} + x^{-\frac{3}{2}}$             **g**  $y = 3 - \frac{2}{3}x^{-1}$             **h**  $y = 2x^{\frac{1}{2}} + \frac{1}{4}x^{\frac{5}{2}}$   
           $\frac{dy}{dx} = -2x^{-2} - x$              $\frac{dy}{dx} = -10x^{-3} - \frac{3}{2}x^{-\frac{5}{2}}$              $\frac{dy}{dx} = \frac{2}{3}x^{-2}$              $\frac{dy}{dx} = x^{-\frac{1}{2}} + \frac{5}{8}x^{\frac{3}{2}}$
- 8**    **a**  $\frac{dy}{dx} = 8x - 1$             **b**  $\frac{dy}{dx} = 3x^2 + 10x + 2$             **c**  $\frac{dy}{dx} = 2x^{-2}$   
           $\frac{d^2y}{dx^2} = 8$              $\frac{d^2y}{dx^2} = 6x + 10$              $\frac{d^2y}{dx^2} = -4x^{-3}$   
          **d**  $\frac{dy}{dx} = 8x^3 + 6x$             **e**  $y = 3x^4 - 4x^{-2}$             **f**  $\frac{dy}{dx} = 3x^{-\frac{1}{2}} + \frac{1}{2}x^{-\frac{3}{2}}$   
           $\frac{d^2y}{dx^2} = 24x^2 + 6$              $\frac{dy}{dx} = 12x^3 + 8x^{-3}$              $\frac{d^2y}{dx^2} = -\frac{3}{2}x^{-\frac{3}{2}} - \frac{3}{4}x^{-\frac{5}{2}}$   
           $\frac{d^2y}{dx^2} = 36x^2 - 24x^{-4}$

# C1 DIFFERENTIATION

# Answers - Worksheet C

- 1    **a**  $\frac{dy}{dx} = 3x^2$                       **b**  $\frac{dy}{dx} = 4 - 2x$                       **c**  $\frac{dy}{dx} = 4x - 8$                       **d**  $\frac{dy}{dx} = -3x^{-2}$   
          grad = 27                              grad = -2                              grad = 4                              grad =  $-\frac{1}{3}$
- 2    **a**  $\frac{dy}{dx} = 6x + 1$   
          at (1, -1) grad = 7                      **b**  $\frac{dy}{dx} = 4x^3 + 6x^2$   
          at (-2, 0) grad = -8
- c**  $y = 2x^2 - 3x$ ,  $\frac{dy}{dx} = 4x - 3$   
          at (2, 2) grad = 5                      **d**  $\frac{dy}{dx} = 2x + 2x^{-2}$   
          at (2, 3) grad =  $\frac{9}{2}$
- e**  $\frac{dy}{dx} = 2x + 6$                       **f**  $\frac{dy}{dx} = 4 - 2x^{-3}$   
          at (-3, -1) grad = 0                      at ( $\frac{1}{2}$ , 6) grad = -12
- 3    **a**  $f(x) = x^2 + 2x + 1$     **b**  $f'(x) = \frac{1}{2}x^{-\frac{1}{2}}$                       **c**  $f'(x) = 1 + 8x^{-3}$                       **d**  $f'(x) = -9x^{\frac{1}{2}}$   
           $f'(x) = 2x + 2$                        $f'(4) = \frac{1}{4}$                        $f'(4) = \frac{9}{8}$                        $f'(4) = -18$   
           $f'(4) = 10$
- 4    **a**  $x(x-1)(x-3) = 0$ ,  $x = 0, 1, 3$   
           $\therefore (0, 0), (1, 0)$  and  $(3, 0)$                       **5** **a**  $\frac{dy}{dx} = 4x - 5$   
          **b**  $\frac{dy}{dx} = 3x^2 - 8x + 3$                       **b**  $4x - 5 = 7$   
          at (0, 0) grad = 3                       $x = 3$   
          at (1, 0) grad = -2  
          at (3, 0) grad = 6
- 6     $\frac{dy}{dx} = 3x^2 - 8$                       **7** **a**  $\frac{dy}{dx} = 3x^2 + 2x - 4$   
           $\therefore 3x^2 - 8 = 4$   
           $x^2 = 4$   
           $x = \pm 2$   
           $\therefore (-2, 8)$  and  $(2, -8)$                       **b** grad at  $P = -3$   
          grad at  $Q = -3$   
           $\therefore 3x^2 + 2x - 4 = -3$   
           $3x^2 + 2x - 1 = 0$   
           $(3x-1)(x+1) = 0$   
           $x = -1$  (at  $P$ ) or  $\frac{1}{3}$   
           $\therefore Q(\frac{1}{3}, -\frac{5}{27})$
- 8    **a**  $\frac{dy}{dx} = 2x$ , grad = 4  
           $\therefore y - 4 = 4(x - 2)$     [ $y = 4x - 4$ ]                      **b**  $\frac{dy}{dx} = 2x + 3$ , grad = 1  
           $\therefore y - 2 = x + 1$     [ $y = x + 3$ ]
- c**  $\frac{dy}{dx} = 4x - 6$ , grad = -2                      **d**  $\frac{dy}{dx} = 3x^2 - 8x$ , grad = 3  
           $\therefore y - 4 = -2(x - 1)$     [ $y = -2x + 6$ ]                       $\therefore y + 7 = 3(x - 3)$     [ $y = 3x - 16$ ]

9 a  $\frac{dy}{dx} = -2x$ , grad = 6

$$\therefore y + 6 = 6(x + 3)$$

$$y + 6 = 6x + 18$$

$$6x - y + 12 = 0$$

c  $\frac{dy}{dx} = 4x + 5$ , grad = 7

$$\therefore y - 2 = 7(x - \frac{1}{2})$$

$$2y - 4 = 14x - 7$$

$$14x - 2y - 3 = 0$$

b  $\frac{dy}{dx} = -2x^{-2}$ , grad =  $-\frac{1}{2}$

$$\therefore y - 1 = -\frac{1}{2}(x - 2)$$

$$2y - 2 = -x + 2$$

$$x + 2y - 4 = 0$$

d  $\frac{dy}{dx} = 1 - \frac{3}{2}x^{-\frac{1}{2}}$ , grad =  $\frac{1}{4}$

$$\therefore y + 2 = \frac{1}{4}(x - 4)$$

$$4y + 8 = x - 4$$

$$x - 4y - 12 = 0$$

10 a  $\frac{dy}{dx} = 2x$ , grad = 2

$$\therefore \text{grad of normal} = -\frac{1}{2}$$

$$\therefore y + 3 = -\frac{1}{2}(x - 1)$$

$$2y + 6 = -x + 1$$

$$x + 2y + 5 = 0$$

b  $\frac{dy}{dx} = 6x + 7$ , grad = -5

$$\therefore \text{grad of normal} = \frac{1}{5}$$

$$\therefore y - 5 = \frac{1}{5}(x + 2)$$

$$5y - 25 = x + 2$$

$$x - 5y + 27 = 0$$

c  $\frac{dy}{dx} = 3x^2 - 8$ , grad = 4

$$\therefore \text{grad of normal} = -\frac{1}{4}$$

$$\therefore y + 4 = -\frac{1}{4}(x - 2)$$

$$4y + 16 = -x + 2$$

$$x + 4y + 14 = 0$$

d  $\frac{dy}{dx} = 1 + 6x^{-2}$ , grad =  $\frac{5}{3}$

$$\therefore \text{grad of normal} = -\frac{3}{5}$$

$$\therefore y - 1 = -\frac{3}{5}(x - 3)$$

$$5y - 5 = -3x + 9$$

$$3x + 5y - 14 = 0$$

11 a  $x = 2 \therefore y = 4$

$$\frac{dy}{dx} = 6x - 5, \text{ grad} = 7$$

$$\therefore y - 4 = 7(x - 2)$$

$$y = 7x - 10$$

b  $x = -3 \therefore y = 6$

$$\frac{dy}{dx} = 3x^2 + 10x, \text{ grad} = -3$$

$$\therefore \text{grad of normal} = \frac{1}{3}$$

$$\therefore y - 6 = \frac{1}{3}(x + 3)$$

$$y = \frac{1}{3}x + 7$$

12 a  $\frac{dy}{dx} = 3x^2 + 6x - 16$ , grad = 8

$$\therefore y + 10 = 8(x - 2) \quad [y = 8x - 26]$$

b  $3x^2 + 6x - 16 = 8$

$$x^2 + 2x - 8 = 0$$

$$(x + 4)(x - 2) = 0$$

$$x = 2 \text{ (at } P) \text{ or } -4$$

$$\therefore Q(-4, 50)$$

13 a  $\frac{dy}{dx} = 2x - 3$ , grad = 1

$$\therefore \text{grad of normal} = -1$$

$$\therefore y - 2 = -(x - 2) \quad [y = 4 - x]$$

b  $x^2 - 3x + 4 = 4 - x$

$$x^2 - 2x = 0$$

$$x(x - 2) = 0$$

$$x = 2 \text{ (at } A) \text{ or } 0$$

$$\therefore B(0, 4)$$

14 a  $f'(x) = 3x^2 + 8x$

b  $x = -3 \therefore y = -9$

$$\text{grad} = 3$$

$$\therefore y + 9 = 3(x + 3)$$

$$y = 3x \text{ which passes through } (0, 0)$$

$$15 \quad \text{a} \quad y = 0 \Rightarrow 6 + x - x^2 = 0$$

$$(2 + x)(3 - x) = 0$$

$$x = -2, 3$$

+ve  $x$ -axis  $\therefore P(3, 0)$

$$x = 0 \Rightarrow y = 6 \therefore Q(0, 6)$$

$$\text{b} \quad \frac{dy}{dx} = 1 - 2x$$

grad at  $P = -5$

$$y = -5(x - 3) \quad [y = 15 - 5x]$$

c grad at  $Q = 1$

tangent at  $Q$ :  $y = x + 6$

$$\therefore 15 - 5x = x + 6$$

$$x = \frac{3}{2}$$

$$\therefore \left(\frac{3}{2}, \frac{15}{2}\right)$$

$$16 \quad \text{a} \quad \text{grad of } l = -3$$

for curve,  $\frac{dy}{dx} = 2x - 5$

$$\therefore \text{at } A, \quad 2x - 5 = -3$$

$$x = 1$$

$$\therefore A(1, -1)$$

$$\text{b} \quad y + 1 = -3(x - 1)$$

$$y = -3x + 2$$

$$17 \quad \text{grad of normal} = 2$$

$$\therefore \text{grad of curve} = -\frac{1}{2}$$

for curve,  $\frac{dy}{dx} = -32x^{-3}$

$$\therefore -\frac{32}{x^3} = -\frac{1}{2}$$

$$x^3 = 64$$

$$x = 4 \therefore (4, 1)$$

$$\text{sub. } 1 = 8 + k$$

$$k = -7$$

$$18 \quad \text{a} \quad \frac{ds}{dt} = 3 + 10t$$

$$t = 0.6 \Rightarrow \frac{ds}{dt} = 9 \text{ metres per second}$$

$$\text{b} \quad 54 = 3t + 5t^2$$

$$5t^2 + 3t - 54 = 0$$

$$(5t + 18)(t - 3) = 0$$

$$t > 0 \therefore t = 3$$

$$\therefore \frac{ds}{dt} = 33 \text{ metres per second}$$

$$19 \quad \text{a} \quad \frac{dh}{dt} = \frac{1}{3}kt^{-\frac{2}{3}}$$

when  $t = 1$ ,  $\frac{dh}{dt} = 3$

$$\therefore \frac{1}{3}k = 3$$

$$k = 9$$


$$\text{b} \quad \frac{dh}{dt} = 3 \times 8^{-\frac{2}{3}} = 0.75 \text{ cm per second}$$



# C1 DIFFERENTIATION

## Answers - Worksheet D

1 a



$$\begin{aligned} \text{b } f(x) &= (x+1)(x^2 - 4x + 4) \\ &= x^3 - 4x^2 + 4x + x^2 - 4x + 4 \\ &= x^3 - 3x^2 + 4 \end{aligned}$$

$$f'(x) = 3x^2 - 6x$$

$$\text{c } x = 1 \quad \therefore y = 2 \times (-1)^2 = 2$$

$$\text{grad} = 3 - 6 = -3$$

$$\therefore y - 2 = -3(x - 1)$$

$$y - 2 = -3x + 3$$

$$y = 5 - 3x$$

3

$$\text{a } x^2 + x - 2 = 0$$

$$(x+2)(x-1) = 0$$

$$x = -2, 1 \quad a < b \quad \therefore a = -2, b = 1$$

$$\text{b } \frac{dy}{dx} = 2x + 1$$

$$\text{grad at A} = -3$$

$$\therefore \text{grad of normal} = \frac{1}{3}$$

$$\therefore y - 0 = \frac{1}{3}(x + 2)$$

$$3y = x + 2$$

$$x - 3y + 2 = 0$$

$$\text{c } \text{grad at B} = 3$$

$$\text{tangent at B: } y - 0 = 3(x - 1)$$

$$y = 3x - 3$$

$$\text{at C, } x - 3(3x - 3) + 2 = 0$$

$$x = \frac{11}{8}$$

$$\therefore C \left( \frac{11}{8}, \frac{9}{8} \right)$$

5

$$\text{a } \frac{dy}{dx} = -24x^{-3}$$

$$\text{at A, } y = 3, \text{ grad} = -3$$

$$\therefore y - 3 = -3(x - 2)$$

$$3x + y - 9 = 0$$

$$\text{b } \text{tangent:}$$

$$x = -1 \Rightarrow -3 + y - 9 = 0 \Rightarrow y = 12$$

curve:

$$x = -1 \Rightarrow y = \frac{12}{1} \Rightarrow y = 12$$

$$\therefore \text{tangent intersects curve at } (-1, 12)$$

2

$$\text{a } \frac{dy}{dx} = 1 - \frac{3}{2}x^{-\frac{1}{2}}$$

$$\text{grad at P} = \frac{1}{4}$$

$$\therefore y - 1 = \frac{1}{4}(x - 4)$$


$$y = \frac{1}{4}x \text{ which passes through } (0, 0)$$

$$\text{b } \text{grad of normal} = -4$$

$$\therefore y - 1 = -4(x - 4) \quad [y = 17 - 4x]$$

$$\text{at Q, } x = 0 \Rightarrow y = 17$$

$$\therefore \text{area} = \frac{1}{2} \times 17 \times 4 = 34$$



4

$$y = \frac{1}{3}x^{\frac{3}{2}} - 2x^{\frac{1}{2}} - x^{-\frac{1}{2}}$$

$$\frac{dy}{dx} = \frac{1}{2}x^{\frac{1}{2}} - x^{-\frac{1}{2}} + \frac{1}{2}x^{-\frac{3}{2}}$$

$$= \frac{x^2 - 2x + 1}{2x^{\frac{3}{2}}}$$

$$= \frac{(x-1)^2}{2x^{\frac{3}{2}}} \quad [a = -1, b = 2]$$

6

$$\text{a } \frac{dy}{dx} = 3 + 2kx - 3x^2$$

$$\text{at P, } 3 - 2k - 3 = -6$$

$$k = 3$$

$$\text{b } y = 2 + 3x + 3x^2 - x^3 \quad \therefore P(-1, 3)$$

$$\text{at Q, } 3 + 6x - 3x^2 = -6$$

$$x^2 - 2x - 3 = 0$$

$$(x+1)(x-3) = 0$$

$$x = -1 \text{ (at P) or } 3 \quad \therefore Q(3, 11)$$

$$PQ = \sqrt{16 + 64} = \sqrt{80} = 4\sqrt{5}$$

$$7 \quad = \frac{d}{dx}(x^2 + \frac{1}{2}x^{-1})$$

$$= 2x - \frac{1}{2}x^{-2}$$

$$8 \quad \text{a} \quad \frac{dy}{dx} = 4x - 7$$

$$\text{at } A, y = -5, \text{ grad} = 1$$

$$\therefore y + 5 = 1(x - 2)$$

$$[y = x - 7]$$

**b** grad of normal at  $B = 1$

$$\therefore \text{grad of curve at } B = -1$$

$$\therefore 4x - 7 = -1$$

$$x = \frac{3}{2}, y = 2(\frac{3}{2}) - 7(\frac{3}{2}) + 1 = -5$$

$$\therefore B(\frac{3}{2}, -5)$$

$$9 \quad \text{a} \quad \frac{dy}{dx} = 2x + \frac{3}{2}x^{-\frac{1}{2}}$$

$$\text{b} \quad \frac{d^2y}{dx^2} = 2 - \frac{3}{4}x^{-\frac{3}{2}}$$

$$\therefore 2x \frac{d^2y}{dx^2} + \frac{dy}{dx} - 6x$$

$$= 2x(2 - \frac{3}{4}x^{-\frac{3}{2}}) + 2x + \frac{3}{2}x^{-\frac{1}{2}} - 6x$$

$$= 4x - \frac{3}{2}x^{-\frac{1}{2}} + 2x + \frac{3}{2}x^{-\frac{1}{2}} - 6x$$

$$= 0$$

$$10 \quad \text{a} \quad \frac{dy}{dx} = -4x^{-2}$$

$$\text{grad at } M = -\frac{1}{4}$$

$$\therefore \text{grad of normal} = 4$$

$$\therefore y - 3 = 4(x - 4) \quad [y = 4x - 13]$$

$$\text{b} \quad 4x - 13 = 2 + \frac{4}{x}$$

$$4x^2 - 15x - 4 = 0$$

$$(4x + 1)(x - 4) = 0$$

$$x = 4 \text{ (at } M) \text{ or } -\frac{1}{4}$$

$$\therefore N(-\frac{1}{4}, -14)$$

$$11 \quad \text{a} \quad \frac{dy}{dx} = 3x^2 - 6x - 8$$

$$\text{grad at } P = 1$$

$$\therefore y - 8 = 1(x + 1) \quad [y = x + 9]$$

$$\text{b} \quad \text{at } Q, \quad 3x^2 - 6x - 8 = 1$$

$$x^2 - 2x - 3 = 0$$

$$(x + 1)(x - 3) = 0$$

$$x = -1 \text{ at } P \therefore Q(3, -20)$$

$$\therefore y + 20 = 1(x - 3) \quad [y = x - 23]$$

**c** grad normal =  $-1$

$$\therefore y - 8 = -(x + 1) \quad [y = 7 - x]$$

**d** normal at  $P$  meets  $m$  when

$$7 - x = x - 23$$

$$x = 15 \therefore (15, -8)$$

$$\text{dist between lines} = \text{dist } P \text{ to } (15, -8)$$

$$= \sqrt{16^2 + 16^2} = \sqrt{16^2 \times 2} = 16\sqrt{2}$$

$$12 \quad \text{a} \quad y = kx^{\frac{1}{2}} - x^{\frac{3}{2}}$$

$$\frac{dy}{dx} = \frac{1}{2}kx^{-\frac{1}{2}} - \frac{3}{2}x^{\frac{1}{2}}$$

$$\text{at } P, \quad \frac{1}{2}k(\frac{1}{\sqrt{2}}) - \frac{3}{2}(\sqrt{2}) = \sqrt{2}$$

$$k - 6 = 4$$

$$k = 10$$

$$\text{b} \quad y = \sqrt{x}(10 - x)$$

$$\text{at } P, y = \sqrt{2}(10 - 2) = 8\sqrt{2}$$

$$\text{grad of normal} = -\frac{1}{\sqrt{2}}$$

$$\therefore y - 8\sqrt{2} = -\frac{1}{\sqrt{2}}(x - 2)$$

$$\sqrt{2}y - 16 = -x + 2$$

$$x + \sqrt{2}y = 18 \quad [c = 18]$$