

Exercise 5C

$$1 \text{ a } 1 \rightarrow 2 \rightarrow 4 \rightarrow 8 \rightarrow 16 \rightarrow 32$$

$$\quad \quad \times 2 \quad \times 2 \quad \times 2 \quad \times 2 \quad \times 2$$

Geometric, $r = 2$

$$2 \rightarrow 5 \rightarrow 8 \rightarrow 11 \rightarrow 14$$

$$\quad \quad +3 \quad +3 \quad +3 \quad +3$$

Not geometric

(this is an arithmetic sequence)

$$40 \rightarrow 36 \rightarrow 32 \rightarrow 28$$

$$\quad \quad -4 \quad -4 \quad -4$$

Not geometric (arithmetic)

$$2 \rightarrow 6 \rightarrow 18 \rightarrow 54$$

$$\quad \quad \times 3 \quad \times 3 \quad \times 3$$

Geometric, $r = 3$

$$10 \rightarrow 5 \rightarrow 2.5 \rightarrow 1.25$$

$$\quad \quad \times \frac{1}{2} \quad \times \frac{1}{2} \quad \times \frac{1}{2}$$

Geometric, $r = \frac{1}{2}$

$$5 \rightarrow -5 \rightarrow 5 \rightarrow -5$$

$$\quad \quad \times (-1) \quad \times (-1) \quad \times (-1)$$

Geometric, $r = -1$

$$3 \rightarrow 3 \rightarrow 3 \rightarrow 3 \rightarrow 3$$

$$\quad \quad \times 1 \quad \times 1 \quad \times 1 \quad \times 1$$

Geometric, $r = 1$

$$4 \rightarrow -1 \rightarrow 0.25 \rightarrow -0.0625$$

$$\quad \quad \times \left(-\frac{1}{4}\right) \quad \times \left(-\frac{1}{4}\right) \quad \times \left(-\frac{1}{4}\right)$$

Geometric, $r = -\frac{1}{4}$

$$2 \text{ a } 5 \rightarrow 15 \rightarrow 45 \rightarrow 135 \rightarrow 405 \rightarrow 1215$$

$$\quad \quad \times 3 \quad \times 3 \quad \times 3 \quad \times 3 \quad \times 3$$

$$4 \rightarrow -8 \rightarrow 16 \rightarrow -32 \rightarrow 64 \rightarrow -128$$

$$\quad \quad \times (-2) \quad \times (-2) \quad \times (-2) \quad \times (-2) \quad \times (-2)$$

$$60 \rightarrow 30 \rightarrow 15 \rightarrow 7.5 \rightarrow 3.75 \rightarrow 1.875$$

$$\quad \quad \times \frac{1}{2} \quad \times \frac{1}{2} \quad \times \frac{1}{2} \quad \times \frac{1}{2} \quad \times \frac{1}{2}$$

$$1 \rightarrow \frac{1}{4} \rightarrow \frac{1}{16} \rightarrow \frac{1}{64} \rightarrow \frac{1}{256} \rightarrow \frac{1}{1024}$$

$$\quad \quad \times \frac{1}{4} \quad \times \frac{1}{4} \quad \times \frac{1}{4} \quad \times \frac{1}{4} \quad \times \frac{1}{4}$$

$$1 \rightarrow p \rightarrow p^2 \rightarrow p^3 \rightarrow p^4 \rightarrow p^5$$

$$\quad \quad \times p \quad \times p \quad \times p \quad \times p \quad \times p$$

$$2 \text{ f } x \rightarrow -2x^2 \rightarrow 4x^3 \rightarrow -8x^4$$

$$\quad \quad \times (-2x) \quad \times (-2x) \quad \times (-2x)$$

$$\quad \quad \rightarrow 16x^5 \rightarrow -32x^6$$

$$\quad \quad \times (-2x) \quad \times (-2x)$$

$$3 \text{ a } 3 \quad x \quad 9$$

$$\text{Common ratio} = \frac{\text{term 2}}{\text{term 1}} \text{ or } \frac{\text{term 3}}{\text{term 2}} \frac{x}{3} \text{ or } \frac{9}{x}$$

Therefore,

$$\frac{x}{3} = \frac{9}{x} \quad (\text{cross multiply})$$

$$x^2 = 27$$

$$x = \sqrt{27}$$

$$x = \sqrt{9 \times 3}$$

$$x = 3\sqrt{3}$$

$$b \text{ Term 4} = \text{term 3} \times r$$

$$\text{Term 3} = 9 \text{ and}$$

$$r = \frac{\text{term 2}}{\text{term 1}} = \frac{3\sqrt{3}}{3} = \sqrt{3}$$

$$\text{So term 4} = 9\sqrt{3}$$

$$4 \text{ a } 2, 6, 18, 54, \dots$$

$$6\text{th term} = 2 \times 3^5$$

$$= 2 \times 243$$

$$= 486$$

$$n\text{th term} = 2 \times 3^{n-1}$$

$$b \text{ } 100, 50, 25, 12.5, \dots$$

$$6\text{th term} = 100 \times \left(\frac{1}{2}\right)^5$$

$$= 100 \times \frac{1}{32}$$

$$= \frac{25}{8}$$

$$n\text{th term} = 100 \times \left(\frac{1}{2}\right)^{n-1}$$

$$c \text{ } 1, -2, 4, -8, \dots$$

$$6\text{th term} = 1 \times (-2)^5$$

$$= 1 \times -32$$

$$= -32$$

$$n\text{th term} = (-2)^{n-1}$$

4 d 1, 1.1, 1.21, 1.331, ...
 6th term = $1 \times (1.1)^5$
 $= 1 \times 1.61051$
 $= 1.61051$
 nth term = $(1.1)^{n-1}$

5 nth term = 2×5^n
 1st term = $2 \times 5^1 = 10$
 5th term = $2 \times 5^5 = 6250$

6 Let the first term be a and the common ratio = r

6th term is 32
 $\Rightarrow ar^{6-1} = 32$
 $\Rightarrow ar^5 = 32$ (1)

3rd term is 4
 $\Rightarrow ar^{3-1} = 4$
 $\Rightarrow ar^2 = 4$ (2)

(1) \div (2):

$$\frac{ar^5}{ar^2} = \frac{32}{4}$$

$$r^3 = 8$$

$$r = 2$$

Common ratio is 2.

Substitute $r = 2$ into equation (2)

$$a \times 2^2 = 4$$

$$a \times 4 = 4$$

$$a = 1$$

First term is 1.

7 First term is 4.
 $\Rightarrow a = 4$ (1)

Third term is 1 $\Rightarrow ar^{3-1} = 1$
 $\Rightarrow ar^2 = 1$ (2)

Substitute $a = 4$ into (2)

$$4r^2 = 1$$

$$r^2 = \frac{1}{4}$$

$$r = \pm \frac{1}{2}$$

The sixth term = $ar^{6-1} = ar^5$

If $r = \frac{1}{2}$ then sixth term = $4 \times \left(\frac{1}{2}\right)^5 = \frac{1}{8}$

If $r = -\frac{1}{2}$ then sixth term = $4 \times \left(-\frac{1}{2}\right)^5$
 $= -\frac{1}{8}$

Possible values for sixth term: $\frac{1}{8}, -\frac{1}{8}$.

8 a $\frac{u_2}{u_1} = \frac{u_3}{u_2}$

$$\frac{2x}{8-x} = \frac{x^2}{2x}$$

$$4x^2 = 8x^2 - x^3$$

$$x^3 - 4x^2 = 0$$

b $x^2(x - 4) = 0$

$$x = 0 \text{ or } 4$$

$$\text{As } x > 0, x = 4$$

$$a = 4, r = 2$$

$$\text{20th term} = ar^{19}$$

$$= 4 \times 2^{19}$$

$$= 4 \times 524\,288$$

$$= 2\,097\,152$$

c If 4096 in the sequence then,
 for some n , $ar^{n-1} = 4096$

$$4 \times 2^{n-1} = 4096$$

$$2^{n-1} = 1024$$

$$n - 1 = 10$$

$$n = 11$$

Yes, 4096 is in the sequence as n is an integer.

$$9 \text{ a } a = 200, r = p$$

$$u_6 = 200p^5 = 40$$

$$p^5 = \frac{1}{5}$$

$$\log p^5 = \log \frac{1}{5}$$

$$5 \log p = \log 1 - \log 5$$

$$5 \log p + \log 5 = 0$$

$$b \quad \log p = \frac{-\log 5}{5}$$

$$p = 10^{\frac{-\log 5}{5}}$$

$$p = 0.725$$

$$10 \text{ a } a = 4, u_4 = 108 = 4r^3$$

$$r^3 = 27$$

$$r = 3$$

We want k th term $> 500\,000$

$$\text{So } 4 \times 3^{k-1} > 500\,000$$

$$3^{k-1} > 125\,000$$

$$\log 3^{k-1} > \log 125\,000$$

$$(k-1) \log 3 > \log 125\,000$$

$$k-1 > \frac{\log 125\,000}{\log 3}$$

$$k-1 > 10.68$$

$$k > 11.68$$

$$\text{So } k = 12$$

$$11 \text{ a } a = 9, r = 4$$

$$u_n = 9 \times 4^{n-1} = 383\,616$$

$$4^{n-1} = 42\,624$$

$$\log 4^{n-1} = \log 42\,624$$

$$(n-1) \log 4 = \log 42\,624$$

$$n-1 = \frac{\log 42\,624}{\log 4}$$

$$n-1 = 7.69$$

$$n = 8.69$$

n is not an integer so 383 616 is not in the sequence.

$$12 \text{ a } a = 3, r = -4$$

$$3, -12, 48, -192, 768, -3072, 12\,288, -49\,152$$

So 49 152 is not in the sequence, but -49 152 is.

$$13 \quad 3 \xrightarrow{\times 4} 12 \xrightarrow{\times 4} 48 \dots$$

This is a geometric series with $a = 3$ and $r = 4$.

If a term exceeds 1 000 000 then

$$ar^{n-1} > 1\,000\,000$$

Substitute $a = 3, r = 4$:

$$3 \times 4^{n-1} > 1\,000\,000$$

$$4^{n-1} > \frac{1\,000\,000}{3}$$

$$\log 4^{n-1} > \log \left(\frac{1\,000\,000}{3} \right)$$

$$(n-1) \log 4 > \log \left(\frac{1\,000\,000}{3} \right)$$

$$n-1 > \frac{\log \left(\frac{1\,000\,000}{3} \right)}{\log 4}$$

$$n-1 > 9.173 \dots$$

$$n > 10.173 \dots$$

$$\text{So } n = 11$$

$$\text{Term is } 3 \times 4^{10} = 3\,145\,728$$