

# Solutionbank C2

## Edexcel Modular Mathematics for AS and A-Level


### Exponentials and logarithms

#### Exercise A, Question 1

#### Question:

- (a) Draw an accurate graph of  $y = (1.7)^x$ , for  $-4 \leq x \leq 4$ .
- (b) Use your graph to solve the equation  $(1.7)^x = 4$ .

#### Solution:



(a)

(b) Where  $y = 4$ ,  $x \approx 2.6$

© Pearson Education Ltd 2008

# Solutionbank C2

## Edexcel Modular Mathematics for AS and A-Level


### Exponentials and logarithms

#### Exercise A, Question 2

#### Question:

- (a) Draw an accurate graph of  $y = (0.6)^x$ , for  $-4 \leq x \leq 4$ .
- (b) Use your graph to solve the equation  $(0.6)^x = 2$ .

#### Solution:



(a)

- (b) Where  $y = 2$ ,  $x \approx -1.4$

© Pearson Education Ltd 2008

# Solutionbank C2

## Edexcel Modular Mathematics for AS and A-Level


### Exponentials and logarithms

#### Exercise A, Question 3

#### Question:

Sketch the graph of  $y = 1^x$ .

#### Solution:



# Solutionbank C2

## Edexcel Modular Mathematics for AS and A-Level

### Exponentials and logarithms

#### Exercise B, Question 1

**Question:**

Rewrite as a logarithm:

(a)  $4^4 = 256$

(b)  $3^{-2} = \frac{1}{9}$

(c)  $10^6 = 1\,000\,000$

(d)  $11^1 = 11$

(e)  $(0.2)^3 = 0.008$

**Solution:**

(a)  $\log_4 256 = 4$

(b)  $\log_3 \left( \frac{1}{9} \right) = -2$

(c)  $\log_{10} 1\,000\,000 = 6$

(d)  $\log_{11} 11 = 1$

(e)  $\log_{0.2} 0.008 = 3$

© Pearson Education Ltd 2008

# Solutionbank C2

## Edexcel Modular Mathematics for AS and A-Level

### Exponentials and logarithms

#### Exercise B, Question 2

**Question:**

Rewrite using a power:

(a)  $\log_2 16 = 4$

(b)  $\log_5 25 = 2$

(c)  $\log_9 3 = \frac{1}{2}$

(d)  $\log_5 0.2 = -1$

(e)  $\log_{10} 100\,000 = 5$

**Solution:**

(a)  $2^4 = 16$

(b)  $5^2 = 25$

(c)  $9^{\frac{1}{2}} = 3$

(d)  $5^{-1} = 0.2$

(e)  $10^5 = 100\,000$

© Pearson Education Ltd 2008

# Solutionbank C2

## Edexcel Modular Mathematics for AS and A-Level

### Exponentials and logarithms

#### Exercise B, Question 3

#### Question:

Find the value of:

(a)  $\log_2 8$

(b)  $\log_5 25$

(c)  $\log_{10} 10\,000\,000$

(d)  $\log_{12} 12$

(e)  $\log_3 729$

(f)  $\log_{10} \sqrt{10}$

(g)  $\log_4 (0.25)$

(h)  $\log_{0.25} 16$

(i)  $\log_a (a^{10})$

(j)  $\log \left( \frac{2}{3} \right) \left( \frac{9}{4} \right)$

#### Solution:

(a) If  $\log_2 8 = x$  then  $2^x = 8$ , so  $x = 3$

(b) If  $\log_5 25 = x$  then  $5^x = 25$ , so  $x = 2$

(c) If  $\log_{10} 10\,000\,000 = x$  then  $10^x = 10\,000\,000$ , so  $x = 7$

(d) If  $\log_{12} 12 = x$  then  $12^x = 12$ , so  $x = 1$

(e) If  $\log_3 729 = x$  then  $3^x = 729$ , so  $x = 6$

(f) If  $\log_{10} \sqrt{10} = x$  then  $10^x = \sqrt{10}$ , so  $x = \frac{1}{2}$

(Power  $\frac{1}{2}$  means 'square root'.)

(g) If  $\log_4 (0.25) = x$  then  $4^x = 0.25 = \frac{1}{4}$ , so  $x = -1$

(Negative power means 'reciprocal'.)

$$(h) \log_{0.25} 16 = x$$

$$\Rightarrow 0.25^x = 16$$

$$\Rightarrow \left(\frac{1}{4}\right)^x = 16, \text{ so } x = -2$$

$$\left[ \left(\frac{1}{4}\right)^{-2} = \frac{1}{\left(\frac{1}{4}\right)^2} = \frac{1}{\left(\frac{1}{16}\right)} = 16 \right]$$

$$(i) \log_a (a^{10}) = x$$

$$\Rightarrow a^x = a^{10}, \text{ so } x = 10$$

$$(j) \log \left(\frac{2}{3}\right) \left(\frac{9}{4}\right) = x$$

$$\Rightarrow \left(\frac{2}{3}\right)^x = \frac{9}{4}, \text{ so } x = -2$$

$$\left[ \left(\frac{2}{3}\right)^{-2} = \frac{1}{\left(\frac{2}{3}\right)^2} = \frac{1}{\left(\frac{4}{9}\right)} = \frac{9}{4} \right]$$

# Solutionbank C2

## Edexcel Modular Mathematics for AS and A-Level

### Exponentials and logarithms

#### Exercise B, Question 4

**Question:**

Find the value of  $x$  for which:

(a)  $\log_5 x = 4$

(b)  $\log_x 81 = 2$

(c)  $\log_7 x = 1$

(d)  $\log_x (2x) = 2$

**Solution:**

(a) Using a power,  $5^4 = x$   
So  $x = 625$

(b) Using a power,  $x^2 = 81$   
So  $x = 9$   
(The base of a logarithm cannot be negative, so  $x = -9$  is not possible.)

(c) Using a power,  $7^1 = x$   
So  $x = 7$

(d) Using a power,  
 $x^2 = 2x$   
 $x^2 - 2x = 0$   
 $x(x - 2) = 0$   
 $x = 2$   
(The base of a logarithm cannot be zero, so  $x = 0$  is not possible.)



# Solutionbank C2

## Edexcel Modular Mathematics for AS and A-Level

### Exponentials and logarithms

#### Exercise C, Question 1

**Question:**

Find from your calculator the value to 3 s.f. of:

$$\log_{10} 20$$

**Solution:**

$$\log_{10} 20 = 1.3010 \dots = 1.30 \text{ (3 s.f.)}$$

© Pearson Education Ltd 2008

# Solutionbank C2

## Edexcel Modular Mathematics for AS and A-Level

### Exponentials and logarithms

#### Exercise C, Question 2

**Question:**

Find from your calculator the value to 3 s.f. of:  
 $\log_{10} 4$

**Solution:**

$$\log_{10} 4 = 0.6020 \dots = 0.602 \text{ (3 s.f.)}$$

© Pearson Education Ltd 2008

# Solutionbank C2

## Edexcel Modular Mathematics for AS and A-Level

### Exponentials and logarithms

#### Exercise C, Question 3

**Question:**

Find from your calculator the value to 3 s.f. of:  
 $\log_{10} 7000$

**Solution:**

$$\log_{10} 7000 = 3.8450 \dots = 3.85 \text{ (3 s.f.)}$$

© Pearson Education Ltd 2008

# Solutionbank C2

## Edexcel Modular Mathematics for AS and A-Level

### Exponentials and logarithms

#### Exercise C, Question 4

**Question:**

Find from your calculator the value to 3 s.f. of:  
 $\log_{10} 0.786$

**Solution:**

$$\log_{10} 0.786 = -0.1045 \dots = -0.105 \text{ (3 s.f.)}$$

© Pearson Education Ltd 2008

# Solutionbank C2

## Edexcel Modular Mathematics for AS and A-Level

### Exponentials and logarithms

#### Exercise C, Question 5

**Question:**

Find from your calculator the value to 3 s.f. of:

$$\log_{10} 11$$

**Solution:**

$$\log_{10} 11 = 1.0413 \dots = 1.04 \text{ (3 s.f.)}$$

© Pearson Education Ltd 2008

# Solutionbank C2

## Edexcel Modular Mathematics for AS and A-Level

### Exponentials and logarithms

#### Exercise C, Question 6

**Question:**

Find from your calculator the value to 3 s.f. of:  
 $\log_{10} 35.3$

**Solution:**

$$\log_{10} 35.3 = 1.5477 \dots = 1.55 \text{ (3 s.f.)}$$

© Pearson Education Ltd 2008

# Solutionbank C2

## Edexcel Modular Mathematics for AS and A-Level

### Exponentials and logarithms

#### Exercise C, Question 7

**Question:**

Find from your calculator the value to 3 s.f. of:  
 $\log_{10} 0.3$

**Solution:**

$$\log_{10} 0.3 = -0.5228 \dots = -0.523 \text{ (3 s.f.)}$$

© Pearson Education Ltd 2008

# Solutionbank C2

## Edexcel Modular Mathematics for AS and A-Level

### Exponentials and logarithms

#### Exercise C, Question 8

**Question:**

Find from your calculator the value to 3 s.f. of:  
 $\log_{10} 999$

**Solution:**

$$\log_{10} 999 = 2.9995 \dots = 3.00 \text{ (3 s.f.)}$$

© Pearson Education Ltd 2008



# Solutionbank C2

## Edexcel Modular Mathematics for AS and A-Level

### Exponentials and logarithms

#### Exercise D, Question 1

#### Question:

Write as a single logarithm:

(a)  $\log_2 7 + \log_2 3$

(b)  $\log_2 36 - \log_2 4$

(c)  $3 \log_5 2 + \log_5 10$

(d)  $2 \log_6 8 - 4 \log_6 3$

(e)  $\log_{10} 5 + \log_{10} 6 - \log_{10} \left( \frac{1}{4} \right)$

#### Solution:

(a)  $\log_2 (7 \times 3) = \log_2 21$

(b)  $\log_2 \left( \frac{36}{4} \right) = \log_2 9$

(c)  $3 \log_5 2 = \log_5 2^3 = \log_5 8$   
 $\log_5 8 + \log_5 10 = \log_5 (8 \times 10) = \log_5 80$

(d)  $2 \log_6 8 = \log_6 8^2 = \log_6 64$   
 $4 \log_6 3 = \log_6 3^4 = \log_6 81$

$\log_6 64 - \log_6 81 = \log_6 \left( \frac{64}{81} \right)$

(e)  $\log_{10} 5 + \log_{10} 6 = \log_{10} (5 \times 6) = \log_{10} 30$

$\log_{10} 30 - \log_{10} \left( \frac{1}{4} \right) = \log_{10} \left[ \frac{30}{\left( \frac{1}{4} \right)} \right] = \log_{10} 120$

# Solutionbank C2

## Edexcel Modular Mathematics for AS and A-Level

### Exponentials and logarithms

#### Exercise D, Question 2

#### Question:

Write as a single logarithm, then simplify your answer:

(a)  $\log_2 40 - \log_2 5$

(b)  $\log_6 4 + \log_6 9$

(c)  $2 \log_{12} 3 + 4 \log_{12} 2$

(d)  $\log_8 25 + \log_8 10 - 3 \log_8 5$

(e)  $2 \log_{10} 20 - (\log_{10} 5 + \log_{10} 8)$

#### Solution:

(a)  $\log_2 \left( \frac{40}{5} \right) = \log_2 8 = 3 \quad \left( 2^3 = 8 \right)$

(b)  $\log_6 (4 \times 9) = \log_6 36 = 2 \quad (6^2 = 36)$

(c)  $\log_{12} (3^2) + \log_{12} (2^4)$   
 $= \log_{12} 9 + \log_{12} 16$   
 $= \log_{12} (9 \times 16)$   
 $= \log_{12} 144$   
 $= 2 \quad (12^2 = 144)$

(d)  $\log_8 (25 \times 10) - \log_8 (5^3)$   
 $= \log_8 250 - \log_8 125$   
 $= \log_8 \left( \frac{250}{125} \right)$   
 $= \log_8 2$   
 $= \frac{1}{3} \quad \left( 8^{\frac{1}{3}} = 2 \right)$

(e)  $\log_{10} (20^2) - \log_{10} (5 \times 8)$   
 $= \log_{10} 400 - \log_{10} 40$   
 $= \log_{10} \left( \frac{400}{40} \right)$   
 $= \log_{10} 10$   
 $= 1 \quad (10^1 = 10)$

# Solutionbank C2

## Edexcel Modular Mathematics for AS and A-Level

### Exponentials and logarithms

#### Exercise D, Question 3

#### Question:

Write in terms of  $\log_a x$ ,  $\log_a y$  and  $\log_a z$ :

(a)  $\log_a (x^3y^4z)$

(b)  $\log_a \left( \frac{x^5}{y^2} \right)$

(c)  $\log_a (a^2x^2)$

(d)  $\log_a \left( \frac{x\sqrt{y}}{z} \right)$

(e)  $\log_a \sqrt{ax}$

#### Solution:

(a)  $\log_a x^3 + \log_a y^4 + \log_a z$   
 $= 3 \log_a x + 4 \log_a y + \log_a z$

(b)  $\log_a x^5 - \log_a y^2$   
 $= 5 \log_a x - 2 \log_a y$

(c)  $\log_a a^2 + \log_a x^2$   
 $= 2 \log_a a + 2 \log_a x$   
 $= 2 + 2 \log_a x \quad (\log_a a = 1)$

(d)  $\log_a x + \log_a y^{\frac{1}{2}} - \log_a z$   
 $= \log_a x + \frac{1}{2} \log_a y - \log_a z$

(e)  $\log_a (ax)^{\frac{1}{2}}$   
 $= \frac{1}{2} \log_a (ax)$   
 $= \frac{1}{2} \log_a a + \frac{1}{2} \log_a x$   
 $= \frac{1}{2} + \frac{1}{2} \log_a x$

# Solutionbank C2

## Edexcel Modular Mathematics for AS and A-Level

### Exponentials and logarithms

#### Exercise E, Question 1

#### Question:

Solve, giving your answer to 3 significant figures:

(a)  $2^x = 75$

(b)  $3^x = 10$

(c)  $5^x = 2$

(d)  $4^{2x} = 100$

(e)  $9^{x+5} = 50$

(f)  $7^{2x-1} = 23$

(g)  $3^{x-1} = 8^{x+1}$

(h)  $2^{2x+3} = 3^{3x+2}$

(i)  $8^{3-x} = 10^x$

(j)  $3^{4-3x} = 4^{x+5}$

#### Solution:

(a)  $2^x = 75$

$$\log 2^x = \log 75$$

$$x \log 2 = \log 75$$

$$x = \frac{\log 75}{\log 2}$$

$$x = 6.23 \text{ (3 s.f.)}$$

(b)  $3^x = 10$

$$\log 3^x = \log 10$$

$$x \log 3 = \log 10$$

$$x = \frac{\log 10}{\log 3}$$

$$x = 2.10 \text{ (3 s.f.)}$$

(c)  $5^x = 2$

$$\log 5^x = \log 2$$

$$x \log 5 = \log 2$$

$$x = \frac{\log 2}{\log 5}$$

$$x = 0.431 \text{ (3 s.f.)}$$

(d)  $4^{2x} = 100$

$$\log 4^{2x} = \log 100$$

$$2x \log 4 = \log 100$$

$$x = \frac{\log 100}{2 \log 4}$$

$$x = 1.66 \text{ (3 s.f.)}$$

$$(e) 9^{x+5} = 50$$

$$\log 9^{x+5} = \log 50$$

$$(x+5) \log 9 = \log 50$$

$$x \log 9 + 5 \log 9 = \log 50$$

$$x \log 9 = \log 50 - 5 \log 9$$

$$x = \frac{\log 50 - 5 \log 9}{\log 9}$$

$$x = -3.22 \text{ (3 s.f.)}$$

$$(f) 7^{2x-1} = 23$$

$$\log 7^{2x-1} = \log 23$$

$$(2x-1) \log 7 = \log 23$$

$$2x \log 7 - \log 7 = \log 23$$

$$2x \log 7 = \log 23 + \log 7$$

$$x = \frac{\log 23 + \log 7}{2 \log 7}$$

$$x = 1.31 \text{ (3 s.f.)}$$

$$(g) 3^{x-1} = 8^{x+1}$$

$$\log 3^{x-1} = \log 8^{x+1}$$

$$(x-1) \log 3 = (x+1) \log 8$$

$$x \log 3 - \log 3 = x \log 8 + \log 8$$

$$x (\log 3 - \log 8) = \log 3 + \log 8$$

$$x = \frac{\log 3 + \log 8}{\log 3 - \log 8}$$

$$x = -3.24 \text{ (3 s.f.)}$$

$$(h) 2^{2x+3} = 3^{3x+2}$$

$$\log 2^{2x+3} = \log 3^{3x+2}$$

$$(2x+3) \log 2 = (3x+2) \log 3$$

$$2x \log 2 + 3 \log 2 = 3x \log 3 + 2 \log 3$$

$$2x \log 2 - 3x \log 3 = 2 \log 3 - 3 \log 2$$

$$x (2 \log 2 - 3 \log 3) = 2 \log 3 - 3 \log 2$$

$$x = \frac{2 \log 3 - 3 \log 2}{2 \log 2 - 3 \log 3}$$

$$x = -0.0617 \text{ (3 s.f.)}$$

$$(i) 8^{3-x} = 10^x$$

$$\log 8^{3-x} = \log 10^x$$

$$(3-x) \log 8 = x \log 10$$

$$3 \log 8 - x \log 8 = x \log 10$$

$$3 \log 8 = x (\log 10 + \log 8)$$

$$x = \frac{3 \log 8}{\log 10 + \log 8}$$

$$x = 1.42 \text{ (3 s.f.)}$$

$$(j) 3^{4-3x} = 4^{x+5}$$

$$\log 3^{4-3x} = \log 4^{x+5}$$

$$(4-3x) \log 3 = (x+5) \log 4$$

$$4 \log 3 - 3x \log 3 = x \log 4 + 5 \log 4$$

$$4 \log 3 - 5 \log 4 = x \log 4 + 3x \log 3$$

$$4 \log 3 - 5 \log 4 = x (\log 4 + 3 \log 3)$$

$$x = \frac{4 \log 3 - 5 \log 4}{\log 4 + 3 \log 3}$$

$$x = -0.542 \text{ (3 s.f.)}$$

© Pearson Education Ltd 2008

# Solutionbank C2

## Edexcel Modular Mathematics for AS and A-Level

### Exponentials and logarithms

#### Exercise E, Question 2

#### Question:

Solve, giving your answer to 3 significant figures:

(a)  $2^{2x} - 6(2^x) + 5 = 0$

(b)  $3^{2x} - 15(3^x) + 44 = 0$

(c)  $5^{2x} - 6(5^x) - 7 = 0$

(d)  $3^{2x} + 3^{x+1} - 10 = 0$

(e)  $7^{2x} + 12 = 7^{x+1}$

(f)  $2^{2x} + 3(2^x) - 4 = 0$

(g)  $3^{2x+1} - 26(3^x) - 9 = 0$

(h)  $4(3^{2x+1}) + 17(3^x) - 7 = 0$

#### Solution:

(a) Let  $y = 2^x$

$$y^2 - 6y + 5 = 0$$

$$(y - 1)(y - 5) = 0$$

So  $y = 1$  or  $y = 5$

If  $y = 1$ ,  $2^x = 1$ ,  $x = 0$

If  $y = 5$ ,  $2^x = 5$

$$\log 2^x = \log 5$$

$$x \log 2 = \log 5$$

$$x = \frac{\log 5}{\log 2}$$

$$x = 2.32 \text{ (3 s.f.)}$$

So  $x = 0$  or  $x = 2.32$

(b) Let  $y = 3^x$

$$y^2 - 15y + 44 = 0$$

$$(y - 4)(y - 11) = 0$$

So  $y = 4$  or  $y = 11$

If  $y = 4$ ,  $3^x = 4$

$$\log 3^x = \log 4$$

$$x \log 3 = \log 4$$

$$x = \frac{\log 4}{\log 3}$$

$$x = 1.26 \text{ (3 s.f.)}$$

If  $y = 11$ ,  $3^x = 11$

$$\log 3^x = \log 11$$

$$x \log 3 = \log 11$$

$$x = \frac{\log 11}{\log 3}$$

$$x = 2.18 \text{ (3 s.f.)}$$

So  $x = 1.26$  or  $x = 2.18$

(c) Let  $y = 5^x$

$$y^2 - 6y - 7 = 0$$

$$(y + 1)(y - 7) = 0$$

$$\text{So } y = -1 \text{ or } y = 7$$

If  $y = -1$ ,  $5^x = -1$ . No solution.

$$\text{If } y = 7, 5^x = 7$$

$$\log 5^x = \log 7$$

$$x \log 5 = \log 7$$

$$x = \frac{\log 7}{\log 5}$$

$$x = 1.21 \text{ (3 s.f.)}$$

(d) Let  $y = 3^x$

$$(3^x)^2 + (3^x \times 3) - 10 = 0$$

$$y^2 + 3y - 10 = 0$$

$$(y + 5)(y - 2) = 0$$

$$\text{So } y = -5 \text{ or } y = 2$$

If  $y = -5$ ,  $3^x = -5$ . No solution.

$$\text{If } y = 2, 3^x = 2$$

$$\log 3^x = \log 2$$

$$x \log 3 = \log 2$$

$$x = \frac{\log 2}{\log 3}$$

$$x = 0.631 \text{ (3 s.f.)}$$

(e) Let  $y = 7^x$

$$(7^x)^2 + 12 = 7^x \times 7$$

$$y^2 + 12 = 7y$$

$$y^2 - 7y + 12 = 0$$

$$(y - 3)(y - 4) = 0$$

$$\text{So } y = 3 \text{ or } y = 4$$

$$\text{If } y = 3, 7^x = 3$$

$$x \log 7 = \log 3$$

$$x = \frac{\log 3}{\log 7}$$

$$x = 0.565 \text{ (3 s.f.)}$$

$$\text{If } y = 4, 7^x = 4$$

$$x \log 7 = \log 4$$

$$x = \frac{\log 4}{\log 7}$$

$$x = 0.712 \text{ (3 s.f.)}$$

$$\text{So } x = 0.565 \text{ or } x = 0.712$$

$$(f) 2^{2x} + 3(2^x) - 4 = 0$$

$$\text{Let } y = 2^x$$

$$\text{Then } y^2 + 3y - 4 = 0$$

$$\text{So } (y + 4)(y - 1) = 0$$

$$\text{So } y = -4 \text{ or } y = 1$$

$$2^x = -4 \text{ has no solution}$$

$$\text{Therefore } 2^x = 1$$

$$\text{So } x = 0 \text{ is the only solution}$$

$$(g) 3^{2x+1} - 26(3^x) - 9 = 0$$

$$\text{Let } y = 3^x$$

$$\text{Then } 3y^2 - 26y - 9 = 0$$

$$\text{So } (3y + 1)(y - 9) = 0$$



$$\text{So } y = -\frac{1}{3} \text{ or } y = 9$$

$$3^x = -\frac{1}{3} \text{ has no solution}$$

$$\text{Therefore } 3^x = 9$$

So  $x = 2$  is the only solution

$$\text{(h) } 4(3^{2x+1}) + 17(3^x) - 7 = 0$$

$$12(3^{2x}) + 17(3^x) - 7 = 0$$

$$\text{Let } y = 3^x$$

$$\text{So } 12y^2 + 17y - 7 = 0$$

$$\text{So } (3y - 1)(4y + 7) = 0$$

$$\text{So } y = \frac{1}{3} \text{ or } y = -\frac{7}{4}$$

$$3^x = -\frac{7}{4} \text{ has no solution}$$

$$\text{Therefore } 3^x = \frac{1}{3}$$

So  $x = -1$  is the only solution

© Pearson Education Ltd 2008

# Solutionbank C2

## Edexcel Modular Mathematics for AS and A-Level

### Exponentials and logarithms

#### Exercise F, Question 1

**Question:**

Find, to 3 decimal places:

(a)  $\log_7 120$

(b)  $\log_3 45$

(c)  $\log_2 19$

(d)  $\log_{11} 3$

(e)  $\log_6 4$

**Solution:**

$$(a) \log_7 120 = \frac{\log_{10} 120}{\log_{10} 7} = 2.460 \text{ (3 d.p.)}$$

$$(b) \log_3 45 = \frac{\log_{10} 45}{\log_{10} 3} = 3.465 \text{ (3 d.p.)}$$

$$(c) \log_2 19 = \frac{\log_{10} 19}{\log_{10} 2} = 4.248 \text{ (3 d.p.)}$$

$$(d) \log_{11} 3 = \frac{\log_{10} 3}{\log_{10} 11} = 0.458 \text{ (3 d.p.)}$$

$$(e) \log_6 4 = \frac{\log_{10} 4}{\log_{10} 6} = 0.774 \text{ (3 d.p.)}$$

© Pearson Education Ltd 2008

# Solutionbank C2

## Edexcel Modular Mathematics for AS and A-Level

### Exponentials and logarithms

#### Exercise F, Question 2

#### Question:

Solve, giving your answer to 3 significant figures:

(a)  $8^x = 14$

(b)  $9^x = 99$

(c)  $12^x = 6$

#### Solution:

(a)  $\log 8^x = \log 14$

$x \log 8 = \log 14$

$$x = \frac{\log_{10} 14}{\log_{10} 8}$$

$x = 1.27$  (3 s.f.)

(b)  $\log 9^x = \log 99$

$x \log 9 = \log 99$

$$x = \frac{\log_{10} 99}{\log_{10} 9}$$

$x = 2.09$  (3 s.f.)

(c)  $\log 12^x = \log 6$

$x \log 12 = \log 6$

$$x = \frac{\log_{10} 6}{\log_{10} 12}$$

$x = 0.721$  (3 s.f.)

# Solutionbank C2

## Edexcel Modular Mathematics for AS and A-Level

### Exponentials and logarithms

#### Exercise F, Question 3

#### Question:

Solve, giving your answer to 3 significant figures:

(a)  $\log_2 x = 8 + 9 \log_x 2$

(b)  $\log_4 x + 2 \log_x 4 + 3 = 0$

(c)  $\log_2 x + \log_4 x = 2$

#### Solution:

(a)  $\log_2 x = 8 + 9 \log_x 2$

$$\log_2 x = 8 + \frac{9}{\log_2 x}$$

Let  $\log_2 x = y$

$$y = 8 + \frac{9}{y}$$

$$y^2 = 8y + 9$$

$$y^2 - 8y - 9 = 0$$

$$(y + 1)(y - 9) = 0$$

So  $y = -1$  or  $y = 9$

If  $y = -1$ ,  $\log_2 x = -1$

$$\Rightarrow x = 2^{-1} = \frac{1}{2}$$

If  $y = 9$ ,  $\log_2 x = 9$

$$\Rightarrow x = 2^9 = 512$$

So  $x = \frac{1}{2}$  or  $x = 512$

(b)  $\log_4 x + 2 \log_x 4 + 3 = 0$

$$\log_4 x + \frac{2}{\log_4 x} + 3 = 0$$

Let  $\log_4 x = y$

$$y + \frac{2}{y} + 3 = 0$$

$$y^2 + 2 + 3y = 0$$

$$y^2 + 3y + 2 = 0$$

$$(y + 1)(y + 2) = 0$$

So  $y = -1$  or  $y = -2$

If  $y = -1$ ,  $\log_4 x = -1$

$$\Rightarrow x = 4^{-1} = \frac{1}{4}$$

If  $y = -2$ ,  $\log_4 x = -2$

$$\Rightarrow x = 4^{-2} = \frac{1}{16}$$

$$\text{So } x = \frac{1}{4} \text{ or } x = \frac{1}{16}$$

$$\text{(c) } \log_2 x + \log_4 x = 2$$

$$\log_2 x + \frac{\log_2 x}{\log_2 4} = 2$$

But  $\log_2 4 = 2$  (because  $2^2 = 4$ ), so

$$\log_2 x + \frac{\log_2 x}{2} = 2$$

$$\frac{3}{2} \log_2 x = 2$$

$$\log_2 x = \frac{4}{3}$$

$$x = 2^{\frac{4}{3}}$$

$$x = 2.52 \text{ (3 s.f.)}$$

© Pearson Education Ltd 2008

# Solutionbank C2

## Edexcel Modular Mathematics for AS and A-Level

### Exponentials and logarithms

#### Exercise G, Question 1

#### Question:

Find the possible values of  $x$  for which  $2^{2x+1} = 3(2^x) - 1$ . **[E]**

#### Solution:

$$2^{2x+1} = 3(2^x) - 1$$

$$2^{2x} \times 2^1 = 3(2^x) - 1$$

$$\text{Let } 2^x = y$$

$$2y^2 = 3y - 1$$

$$2y^2 - 3y + 1 = 0$$

$$(2y - 1)(y - 1) = 0$$

$$\text{So } y = \frac{1}{2} \text{ or } y = 1$$

$$\text{If } y = \frac{1}{2}, 2^x = \frac{1}{2}, x = -1$$

$$\text{If } y = 1, 2^x = 1, x = 0$$

$$\text{So } x = 0 \text{ or } x = -1$$

© Pearson Education Ltd 2008

# Solutionbank C2

## Edexcel Modular Mathematics for AS and A-Level

### Exponentials and logarithms

#### Exercise G, Question 2

#### Question:

(a) Express  $\log_a (p^2q)$  in terms of  $\log_a p$  and  $\log_a q$ .

(b) Given that  $\log_a (pq) = 5$  and  $\log_a (p^2q) = 9$ , find the values of  $\log_a p$  and  $\log_a q$ . **[E]**

#### Solution:

$$(a) \log_a (p^2q) = \log_a (p^2) + \log_a q = 2 \log_a p + \log_a q$$

$$(b) \log_a (pq) = \log_a p + \log_a q$$

So

$$\log_a p + \log_a q = 5 \quad \textcircled{1}$$

$$2 \log_a p + \log_a q = 9 \quad \textcircled{2}$$

Subtracting equation  $\textcircled{1}$  from equation  $\textcircled{2}$ :

$$\log_a p = 4$$

$$\text{So } \log_a q = 1$$

# Solutionbank C2

## Edexcel Modular Mathematics for AS and A-Level

### Exponentials and logarithms

#### Exercise G, Question 3

**Question:**

Given that  $p = \log_q 16$ , express in terms of  $p$ ,

(a)  $\log_q 2$ ,

(b)  $\log_q (8q)$ . [E]

**Solution:**

(a)  $p = \log_q 16$

$$p = \log_q (2^4)$$

$$p = 4 \log_q 2$$

$$\log_q 2 = \frac{p}{4}$$

(b)  $\log_q (8q) = \log_q 8 + \log_q q$

$$= \log_q (2^3) + \log_q q$$

$$= 3 \log_q 2 + \log_q q$$

$$= \frac{3p}{4} + 1$$



# Solutionbank C2

## Edexcel Modular Mathematics for AS and A-Level

### Exponentials and logarithms

#### Exercise G, Question 4

#### Question:

- (a) Given that  $\log_3 x = 2$ , determine the value of  $x$ .
- (b) Calculate the value of  $y$  for which  $2 \log_3 y - \log_3 (y + 4) = 2$ .
- (c) Calculate the values of  $z$  for which  $\log_3 z = 4 \log_z 3$ .

#### [E]

#### Solution:

(a)  $\log_3 x = 2$

$$x = 3^2 = 9$$

(b)  $2 \log_3 y - \log_3 (y + 4) = 2$

$$\log_3 (y^2) - \log_3 (y + 4) = 2$$

$$\log_3 \left( \frac{y^2}{y + 4} \right) = 2$$

$$\frac{y^2}{y + 4} = 9$$

$$y^2 = 9y + 36$$

$$y^2 - 9y - 36 = 0$$

$$(y + 3)(y - 12) = 0$$

$$y = -3 \text{ or } y = 12$$

But  $\log_3 (-3)$  is not defined,

So  $y = 12$

(c)  $\log_3 z = 4 \log_z 3$

$$\log_3 z = \frac{4}{\log_3 z}$$

$$(\log_3 z)^2 = 4$$

Either  $\log_3 z = 2$  or  $\log_3 z = -2$

$$z = 3^2 \text{ or } z = 3^{-2}$$

$$z = 9 \text{ or } z = \frac{1}{9}$$

# Solutionbank C2

## Edexcel Modular Mathematics for AS and A-Level

### Exponentials and logarithms

#### Exercise G, Question 5

#### Question:

(a) Using the substitution  $u = 2^x$ , show that the equation  $4^x - 2^{(x+1)} - 15 = 0$  can be written in the form  $u^2 - 2u - 15 = 0$ .

(b) Hence solve the equation  $4^x - 2^{(x+1)} - 15 = 0$ , giving your answer to 2 decimal places. **[E]**

#### Solution:

$$(a) 4^x - 2^{(x+1)} - 15 = 0$$

$$4^x = (2^2)^x = (2^x)^2$$

$$2^{x+1} = 2^x \times 2^1$$

$$\text{Let } u = 2^x$$

$$u^2 - 2u - 15 = 0$$

$$(b) (u + 3)(u - 5) = 0$$

$$\text{So } u = -3 \text{ or } u = 5$$

If  $u = -3$ ,  $2^x = -3$ . No solution.

$$\text{If } u = 5, 2^x = 5$$

$$\log 2^x = \log 5$$

$$x \log 2 = \log 5$$

$$x = \frac{\log 5}{\log 2}$$

$$x = 2.32 \text{ (2 d.p.)}$$

# Solutionbank C2

## Edexcel Modular Mathematics for AS and A-Level

### Exponentials and logarithms

#### Exercise G, Question 6

#### Question:

Solve, giving your answers as exact fractions, the simultaneous equations:

$$8^y = 4^{2x+3}$$
$$\log_2 y = \log_2 x + 4. \quad \text{[E]}$$

#### Solution:

$$8^y = 4^{2x+3}$$
$$(2^3)^y = (2^2)^{2x+3}$$
$$2^{3y} = 2^{2(2x+3)}$$
$$3y = 4x + 6 \quad \text{①}$$

$$\log_2 y - \log_2 x = 4$$

$$\log_2 \left( \frac{y}{x} \right) = 4$$

$$\frac{y}{x} = 2^4 = 16$$

$$y = 16x \quad \text{②}$$

Substitute ② into ①:

$$48x = 4x + 6$$

$$44x = 6$$

$$x = \frac{3}{22}$$

$$y = 16x = \frac{48}{22} = 2 \frac{2}{11}$$

$$\text{So } x = \frac{3}{22}, y = 2 \frac{2}{11}$$

# Solutionbank C2

## Edexcel Modular Mathematics for AS and A-Level

### Exponentials and logarithms

#### Exercise G, Question 7

#### Question:

Find the values of  $x$  for which  $\log_3 x - 2 \log_x 3 = 1$ . [E]

#### Solution:

$$\log_3 x - 2 \log_x 3 = 1$$

$$\log_3 x - \frac{2}{\log_3 x} = 1$$

$$\text{Let } \log_3 x = y$$

$$y - \frac{2}{y} = 1$$

$$y^2 - 2 = y$$

$$y^2 - y - 2 = 0$$

$$(y + 1)(y - 2) = 0$$

$$\text{So } y = -1 \text{ or } y = 2$$

$$\text{If } y = -1, \log_3 x = -1$$

$$\Rightarrow x = 3^{-1} = \frac{1}{3}$$

$$\text{If } y = 2, \log_3 x = 2$$

$$\Rightarrow x = 3^2 = 9$$

$$\text{So } x = \frac{1}{3} \text{ or } x = 9$$

# Solutionbank C2

## Edexcel Modular Mathematics for AS and A-Level

### Exponentials and logarithms

#### Exercise G, Question 8

#### Question:

Solve the equation

$$\log_3 (2 - 3x) = \log_9 (6x^2 - 19x + 2) . \quad \text{[E]}$$

#### Solution:

$$\log_3 (2 - 3x) = \log_9 (6x^2 - 19x + 2)$$

$$\log_9 \left( 6x^2 - 19x + 2 \right) = \frac{\log_3 (6x^2 - 19x + 2)}{\log_3 9} = \frac{\log_3 (6x^2 - 19x + 2)}{2}$$

So

$$2 \log_3 (2 - 3x) = \log_3 (6x^2 - 19x + 2)$$

$$\log_3 (2 - 3x)^2 = \log_3 (6x^2 - 19x + 2)$$

$$(2 - 3x)^2 = 6x^2 - 19x + 2$$

$$4 - 12x + 9x^2 = 6x^2 - 19x + 2$$

$$3x^2 + 7x + 2 = 0$$

$$(3x + 1)(x + 2) = 0$$

$$x = -\frac{1}{3} \text{ or } x = -2$$

(Both solutions are valid, since they give logs of positive numbers in the original equation.)

# Solutionbank C2

## Edexcel Modular Mathematics for AS and A-Level

### Exponentials and logarithms

#### Exercise G, Question 9

#### Question:

If  $xy = 64$  and  $\log_x y + \log_y x = \frac{5}{2}$ , find  $x$  and  $y$ . [E]

#### Solution:

$$\log_x y + \log_y x = \frac{5}{2}$$

$$\log_x y + \frac{1}{\log_x y} = \frac{5}{2}$$

Let  $\log_x y = u$

$$u + \frac{1}{u} = \frac{5}{2}$$

$$2u^2 + 2 = 5u$$

$$2u^2 - 5u + 2 = 0$$

$$(2u - 1)(u - 2) = 0$$

$$u = \frac{1}{2} \text{ or } u = 2$$

$$\text{If } u = \frac{1}{2}, \log_x y = \frac{1}{2}$$

$$\Rightarrow y = x^{\frac{1}{2}} = \sqrt{x}$$

Since  $xy = 64$ ,

$$x \sqrt{x} = 64 \quad \left( x^{\frac{3}{2}} = 64 \right)$$

$$x = 16$$

$$y = \sqrt{x} = 4$$

If  $u = 2, \log_x y = 2$

$$\Rightarrow y = x^2$$

Since  $xy = 64$ ,

$$x^3 = 64$$

$$x = 4$$

$$y = x^2 = 16$$

So  $x = 16, y = 4$  or  $x = 4, y = 16$

# Solutionbank C2

## Edexcel Modular Mathematics for AS and A-Level

### Exponentials and logarithms

#### Exercise G, Question 10

#### Question:

Prove that if  $a^x = b^y = (ab)^{xy}$ , then  $x + y = 1$ . [E]

#### Solution:

Given that  $a^x = b^y = (ab)^{xy}$

Take logs to base  $a$  for  $a^x = b^y$ :

$$\log_a (a^x) = \log_a (b^y)$$

$$x \log_a a = y \log_a b$$

$$x = y \log_a b \quad \textcircled{1}$$

Take logs to base  $a$  for  $a^x = (ab)^{xy}$

$$x = \log_a (ab)^{xy}$$

$$x = xy \log_a (ab)$$

$$x = xy (\log_a a + \log_a b)$$

$$x = xy (1 + \log_a b)$$

$$1 = y (1 + \log_a b) \quad \textcircled{2}$$

But, from  $\textcircled{1}$ ,  $\log_a b = \frac{x}{y}$

Substitute into  $\textcircled{2}$ :

$$1 = y \left( 1 + \frac{x}{y} \right)$$

$$1 = y + x$$

$$x + y = 1$$

# Solutionbank C2

## Edexcel Modular Mathematics for AS and A-Level

### Exponentials and logarithms

#### Exercise G, Question 11

#### Question:

(a) Show that  $\log_4 3 = \log_2 \sqrt{3}$ .

(b) Hence or otherwise solve the simultaneous equations:

$$2 \log_2 y = \log_4 3 + \log_2 x,$$

$$3^y = 9^x,$$

given that  $x$  and  $y$  are positive. **[E]**

#### Solution:

$$(a) \log_4 3 = \frac{\log_2 3}{\log_2 4} = \frac{\log_2 3}{2}$$

$$\log_4 3 = \frac{1}{2} \log_2 3 = \log_2 3^{\frac{1}{2}} = \log_2 \sqrt{3}$$

$$(b) 3^y = 9^x$$

$$3^y = (3^2)^x = 3^{2x}$$

$$\text{So } y = 2x$$

$$2 \log_2 y = \log_4 3 + \log_2 x$$

$$\log_2 (y^2) = \log_2 \sqrt{3} + \log_2 x = \log_2 (x \sqrt{3})$$

$$\text{So } y^2 = x \sqrt{3}$$

$$\text{Since } y = 2x, (2x)^2 = x \sqrt{3}$$

$$\Rightarrow 4x^2 = x \sqrt{3}$$

$$x \text{ is positive, so } x \neq 0, x = \frac{\sqrt{3}}{4}$$

$$\Rightarrow y = 2x = \frac{\sqrt{3}}{2}$$

$$\text{So } x = \frac{\sqrt{3}}{4}, y = \frac{\sqrt{3}}{2}$$



# Solutionbank C2

## Edexcel Modular Mathematics for AS and A-Level

### Exponentials and logarithms

#### Exercise G, Question 12

#### Question:

- (a) Given that  $3 + 2 \log_2 x = \log_2 y$ , show that  $y = 8x^2$ .
- (b) Hence, or otherwise, find the roots  $\alpha$  and  $\beta$ , where  $\alpha < \beta$ , of the equation  $3 + 2 \log_2 x = \log_2 (14x - 3)$ .
- (c) Show that  $\log_2 \alpha = -2$ .
- (d) Calculate  $\log_2 \beta$ , giving your answer to 3 significant figures. **[E]**

#### Solution:

$$(a) 3 + 2 \log_2 x = \log_2 y$$

$$\log_2 y - 2 \log_2 x = 3$$

$$\log_2 y - \log_2 x^2 = 3$$

$$\log_2 \left( \frac{y}{x^2} \right) = 3$$

$$\frac{y}{x^2} = 2^3 = 8$$

$$y = 8x^2$$

(b) Comparing equations,

$$y = 14x - 3$$

$$8x^2 = 14x - 3$$

$$8x^2 - 14x + 3 = 0$$

$$(4x - 1)(2x - 3) = 0$$

$$x = \frac{1}{4} \text{ or } x = \frac{3}{2}$$

$$\alpha = \frac{1}{4}, \beta = \frac{3}{2}$$

$$(c) \log_2 \alpha = \log_2 \left( \frac{1}{4} \right) = -2,$$

$$\text{since } 2^{-2} = \frac{1}{2^2} = \frac{1}{4}$$

$$(d) \log_2 \beta = \log_2 \left( \frac{3}{2} \right)$$

$$\log_2 1.5 = \frac{\log_{10} 1.5}{\log_{10} 2} = 0.585 \text{ (3 s.f.)}$$