




**EDEXCEL CORE MATHEMATICS C3****PRACTICE PAPER A5 MARK SCHEME**

Question Number	Scheme	Marks
1.	$y = \tan x = \frac{\sin x}{\cos x}$ $\frac{dy}{dx} = \frac{\cos x \cos x - \sin x(-\sin x)}{\cos^2 x}$ (use of quotient rule) $= \frac{1}{\cos^2 x} = \sec^2 x$ *	M1 M1 A1 A1 <b>(4 marks)</b>
2. (a)	$2 + \frac{3}{x+2} \left(= \frac{2(x+2)+3}{x+2}\right) = \underline{\underline{\frac{2x+7}{x+2}}} \text{ or } \underline{\underline{\frac{2(x+2)+3}{x+2}}}$	B1 (1)
(b)	$y = 2 + \frac{3}{x+2}$ or $y = \frac{2x+7}{x+2}$ $y - 2 = \frac{3}{x+2}$ $y(x+2) = 2x+7 = yx - 2x = 7 - 2y$ $x+2 = \frac{3}{y-2}$ $x(y-2) = 7 - 2y$ $x = \frac{3}{y-2} - 2$ $x = \frac{7-2y}{y-2}$ $\therefore f^{-1}(x) = \underline{\underline{\frac{3}{x-2}}} - 2$ $f^{-1}(x) = \underline{\underline{\frac{7-2x}{x-2}}} \text{ o.e}$	M1 M1 A1 (3)
(c)	Domain of $f^{-1}(x)$ is $x \in \mathbb{R}, x \neq 2$ [NB $x \neq +2$ ]	B1 (1)  <b>(5 marks)</b>
3. (a)	$\frac{2}{x-3} + \frac{13}{(x-3)(x+7)}$ $= \frac{2(x+7) + 13}{(x-3)(x+7)} = \frac{2x+27}{(x-3)(x+7)}$	M1 M1 A1 (3)
(b)	$2x+27 = x^2 + 4x - 21$ $x^2 + 2x - 48 = (x+8)(x-6) = 0$ $x = -8, 6$	M1 M1 A1 (3)  <b>(6 marks)</b>

Question Number	Scheme	Marks
4. (a)	$\frac{x^2 + 4x + 3}{x^2 + x} = \frac{(x+3)(x+1)}{x(x+1)}$ $= \frac{x+3}{x} \text{ or } 1 + \frac{3}{x}$	Attempt to factorise numerator or denominator M1 A1 (2)
(b)	LHS = $\log_2\left(\frac{x^2 + 4x + 3}{x^2 + x}\right)$ RHS = $2^4$ or 16 $x + 3 = 16x$ $x = \frac{3}{15}$ or $\frac{1}{5}$ or 0.2	Use of $\log a - \log b$ M1 B1 Linear or quadratic equation in $x$ M1 A1 (4) <b>(6 marks)</b>
5. (i)	Choosing values of $A$ and $B$ and attempting to evaluate LHS and RHS of statement  Showing that LHS $\neq$ RHS + conclusion	M1  A1 (2)
(ii)	Using $\tan \theta \equiv \frac{\sin \theta}{\cos \theta}$  to obtain $\frac{\sin^2 \theta + \cos^2 \theta}{\sin \theta \cos \theta}$  Using $\cos^2 \theta + \sin^2 \theta \equiv 1$  Using $2 \sin \theta \cos \theta \equiv \sin 2\theta$  Leading without any error or fudge to $2 \operatorname{cosec} 2\theta$	M1  A1  M1  M1  A1 cso (5)  <b>(7 marks)</b>

Question Number	Scheme	Marks
6. (a)	$\text{LHS} = \frac{2 \sin^2 \theta}{2 \sin \theta \cos \theta}$ $= \frac{\sin \theta}{\cos \theta} = \tan \theta = \text{RHS}$	M1 A1 A1 c.s.o (3)
(b)	From (a) $\frac{1-\cos 2\theta}{\tan \theta} = \frac{1}{2}$ $\sin 2\theta = \frac{1}{2}$ $2\theta = \frac{\pi}{6}, \frac{5\pi}{6}$ $\theta = \frac{\pi}{12}, \frac{5\pi}{12}$	M1 A1 B1 M1 M1 A1 cao (6) <b>(9 marks)</b>
7. (a) (i)	$x = a^y$	B1 (1)
(ii)	In both sides of (i) i.e $\ln x = \ln a^y$ or ( $y =$ ) $\log_a x = \frac{\ln x}{\ln a}$ $= y \ln a *$ $\Rightarrow y \ln a = \ln x$	B1 cso (1)
(b)	$y = \frac{1}{\ln a} \times \ln x, \Rightarrow \frac{dy}{dx} = \frac{1}{\ln a} \times \frac{1}{x} *$  $\left[ \text{or } \frac{1}{x} = \frac{dy}{dx} \cdot \ln a, \Rightarrow \frac{dy}{dx} = \frac{1}{x \ln a} * \right]$	M1, A1 cso  <b>(2)</b>
(c)	$\log_{10} 10 = 1 \Rightarrow A \text{ is } (10, 1)$ $y_A = 1$	B1
	from(b) $m = \frac{1}{10 \ln a} \text{ or } \frac{1}{10 \ln 10} \text{ or } 0.043 \text{ (or better)}$	B1
	equ of target $y - 1 = m(x - 10)$	M1
	i.e $y - 1 = \frac{1}{10 \ln 10} (x - 10) \text{ or } y = \frac{1}{10 \ln 10} x + 1 - \frac{1}{\ln 10}$ (o.e)	A1 (4)
(d)	$y = 0 \text{ in (c)} \Rightarrow 0 = \frac{x}{10 \ln 10} + 1 - \frac{1}{\ln 10} \Rightarrow x = 10 \ln 10 \left( \frac{1}{\ln 10} - 1 \right)$  $x = 10 - 10 \ln 10 \text{ or } 10 (1 - \ln 10) \text{ or } 10 \ln 10 \left( \frac{1}{\ln 10} - 1 \right)$	M1 A1 (2) <b>(10 marks)</b>

Question Number	Scheme	Marks
8. (a)	 <p>Shape  <math>p = \frac{1}{3}</math> or <math>\{\frac{1}{3}, 0\}</math> seen</p>	B1 B1 (2)
(b)	<p>Gradient of tangent at <math>Q = \frac{1}{q}</math></p> <p>Gradient of normal = <math>-q</math></p> <p>Attempt at equation of <math>OQ</math> [<math>y = -qx</math>] and substituting <math>x = q, y = \ln 3q</math></p> <p>or attempt at equation of tangent [<math>y - 3 \ln q = -q(x - q)</math>] with <math>x = 0, y = 0</math></p> <p>or equating gradient of normal to <math>(\ln 3q)/q</math></p> <p><math>q^2 + \ln 3q = 0</math> (*)</p>	B1 M1 M1 A1 (4)
(c)	$\ln 3x = -x^2 \Rightarrow 3x = e^{-x^2}; \Rightarrow x = \frac{1}{3}e^{-x^2}$	M1; A1 (2)
(d)	$x_1 = 0.298280; x_2 = 0.304957, x_3 = 0.303731, x_4 = 0.303958$ Root = 0.304 (3 decimal places)	M1; A1 A1 (3) <b>(11 marks)</b>

Question Number	Scheme	Marks
9. (a)	 <p>shape intersections with axes <math>(c, 0), (0, d)</math></p>	B1 B1 (2)
(b)	 <p>shape <math>x</math> intersection <math>(\frac{1}{2}d, 0)</math> <math>y</math> intersection <math>(0, 3c)</math></p>	B1 B1 B1 (3)
(c)(i)	$c = 2$	B1
(ii)	$-1 < f(x) \leq$ (candidate's) $c$ value	B1 B1 ft (3)
(d)	$3(2^{-x}) = 1 \Rightarrow 2^{-x} = \frac{1}{3}$ and take logs; $-x = \frac{\ln \frac{1}{3}}{\ln 2}$ $d$ (or $x$ ) = 1.585 (3 decimal places)	M1; A1 A1 (3)
(e)	$fg(x) = f[\log_2 x] = [3(2^{-\log_2 x}) - 1]; = [3(2^{\log_2 \frac{1}{x}}) - 1]$ or $\frac{3}{2^{\log_2 x}} - 1$ $= \frac{3}{x} - 1$	M1; A1 A1 (3)
		<b>(14 marks)</b>