

C3 JAN 11

1. (a) Express  $7 \cos x - 24 \sin x$  in the form  $R \cos(x + \alpha)$  where  $R > 0$  and  $0 < \alpha < \frac{\pi}{2}$ .  
Give the value of  $\alpha$  to 3 decimal places.

(3)

(b) Hence write down the minimum value of  $7 \cos x - 24 \sin x$ .

(1)

(c) Solve, for  $0 \leq x < 2\pi$ , the equation

$$7 \cos x - 24 \sin x = 10$$

giving your answers to 2 decimal places.

(5)

$$\begin{aligned} a) R \cos(x + \alpha) &= R \cos x \cos \alpha - R \sin x \sin \alpha \\ R \cos(x + \alpha) &= 7 \cos x - 24 \sin x \end{aligned}$$

$$\frac{R \sin \alpha = 24}{R \cos \alpha = 7} \Rightarrow \tan \alpha = \frac{24}{7} \Rightarrow \alpha = 1.287^\circ$$

$$b) R^2 = 24^2 + 7^2 \Rightarrow R = 25$$

$$25 \cos(x + 1.287) \quad \text{min} = -25$$

$$c) 25 \cos(x + 1.287) = 10$$

$$x + 1.287 = \cos^{-1}\left(\frac{10}{25}\right) = 1.107, 5.176, 7.176$$

$$-1.287 \Rightarrow x = 3.84^\circ, 6.16^\circ$$

Leave blank

2. (a) Express

$$\frac{4x-1}{2(x-1)} - \frac{3}{2(x-1)(2x-1)}$$

as a single fraction in its simplest form.

(4)

Given that

$$f(x) = \frac{4x-1}{2(x-1)} - \frac{3}{2(x-1)(2x-1)} - 2, \quad x > 1,$$

(b) show that

$$f(x) = \frac{3}{2x-1}$$

(2)

(c) Hence differentiate  $f(x)$  and find  $f'(2)$ .

(3)

$$\begin{aligned} a) \frac{(4x-1)(2x-1) - 3}{2(x-1)(2x-1)} &= \frac{8x^2 - 6x + 1 - 3}{2(x-1)(2x-1)} \\ &= \frac{2(4x+1)(x-1)}{2(x-1)(2x-1)} = \frac{4x+1}{2x-1} \end{aligned}$$

$$b) f(x) = \frac{4x+1}{2x-1} - 2 \times \frac{(2x-1)}{1 \times (2x-1)} = \frac{4x+1 - 4x + 2}{2x-1}$$

$$f(x) = \frac{3}{2x-1} \quad \#$$

$$\begin{aligned} c) f(x) &= 3(2x-1)^{-1} \Rightarrow f'(x) = -3 \times 2(2x-1)^{-2} \\ f'(x) &= \frac{-6}{(2x-1)^2} \Rightarrow f'(2) = \frac{-6}{3^2} = \frac{-2}{3} \end{aligned}$$



3. Find all the solutions of

$$2 \cos 2\theta = 1 - 2 \sin \theta$$

in the interval  $0 \leq \theta < 360^\circ$ .

(6)

$$2 - 4 \sin^2 \theta = 1 - 2 \sin \theta$$

$$\Rightarrow 4 \sin^2 \theta - 2 \sin \theta - 1 = 0$$

$$\Rightarrow \sin^2 \theta - \frac{1}{2} \sin \theta = \frac{1}{4}$$

$$\Rightarrow \left(\sin \theta - \frac{1}{4}\right)^2 = \frac{1}{4} + \frac{1}{16} = \frac{5}{16}$$

$$\Rightarrow \sin \theta = \frac{1}{4} \pm \frac{\sqrt{5}}{4} = \frac{1 \pm \sqrt{5}}{4}$$

$$\theta = \sin^{-1}\left(\frac{1+\sqrt{5}}{4}\right) = \underline{54^\circ}, \underline{126^\circ}$$

$$\theta = \sin^{-1}\left(\frac{1-\sqrt{5}}{4}\right) = \cancel{-18^\circ}, \underline{198^\circ}, \underline{342^\circ}$$

4. Joan brings a cup of hot tea into a room and places the cup on a table. At time  $t$  minutes after Joan places the cup on the table, the temperature,  $\theta^\circ\text{C}$ , of the tea is modelled by the equation

$$\theta = 20 + Ae^{-kt}$$

where  $A$  and  $k$  are positive constants.

Given that the initial temperature of the tea was  $90^\circ\text{C}$ ,

(a) find the value of  $A$ .

(2)

The tea takes 5 minutes to decrease in temperature from  $90^\circ\text{C}$  to  $55^\circ\text{C}$ .

(b) Show that  $k = \frac{1}{5} \ln 2$ .

(3)

(c) Find the rate at which the temperature of the tea is decreasing at the instant when  $t = 10$ . Give your answer, in  $^\circ\text{C}$  per minute, to 3 decimal places.

(3)

$$\text{a) } 90 = 20 + A \Rightarrow \underline{A = 70}$$

$$\text{b) } 55 = 20 + 70e^{-5k} \Rightarrow 35 = 70e^{-5k}$$

$$\Rightarrow e^{-5k} = \frac{1}{2} \Rightarrow -5k = \ln\left(\frac{1}{2}\right) = \ln(2^{-1}) = -\ln 2$$

$$\Rightarrow k = \frac{1}{5} \ln 2 \quad \#$$

$$\text{c) } \frac{d\theta}{dt} = (70 \times -\frac{1}{5} \ln 2) e^{-\frac{1}{5} \ln 2 t} = -14 \ln 2 e^{-\frac{1}{5} \ln 2 t}$$

$$t=10 \quad \frac{d\theta}{dt} = (-14 \ln 2) e^{-2 \ln 2} = -2.426$$

$\Rightarrow$  decreasing at a rate of  $2.426^\circ\text{C}/\text{min}$






Figure 1

Figure 1 shows a sketch of part of the curve with equation  $y = f(x)$ , where

$$f(x) = (8-x)\ln x, \quad x > 0$$

The curve cuts the  $x$ -axis at the points  $A$  and  $B$  and has a maximum turning point at  $Q$ , as shown in Figure 1.

(a) Write down the coordinates of  $A$  and the coordinates of  $B$ .

(2)

(b) Find  $f'(x)$ .

(3)

(c) Show that the  $x$ -coordinate of  $Q$  lies between 3.5 and 3.6

(2)

(d) Show that the  $x$ -coordinate of  $Q$  is the solution of

$$x = \frac{8}{1 + \ln x}$$

(3)

To find an approximation for the  $x$ -coordinate of  $Q$ , the iteration formula

$$x_{n+1} = \frac{8}{1 + \ln x_n}$$

is used.

(e) Taking  $x_0 = 3.55$ , find the values of  $x_1$ ,  $x_2$  and  $x_3$ .

Give your answers to 3 decimal places.

(3)

5a)  $(8-x)\ln x = 0$  when  $x=8, x=1$   $A(1,0)$   $B(8,0)$

b)  $f'(x) = (8-x)' \ln x + (8-x)(\ln x)'$   
 $f'(x) = -\ln x + \frac{8-x}{x}$

c)  $f'(3.5) = 0.033 \Rightarrow$  Increasing function at  $x=3.5$   
 $f'(3.6) = -0.059 \Rightarrow$  decreasing function at  $x=3.6$   
 $\Rightarrow$  turning point between 3.5 and 3.6.

d)  $f'(x) = 0 \Rightarrow \frac{8-x}{x} = \ln x \Rightarrow 8-x = x \ln x$   
 $\Rightarrow 8 = x + x \ln x \Rightarrow 8 = x(1 + \ln x) \Rightarrow x = \frac{8}{1 + \ln x}$

e)  $x_0 = \underline{3.55}$ ;  $x_1 = \underline{3.529}$ ;  $x_2 = \underline{3.538}$ ;  $x_3 = \underline{3.534}$

6. The function  $f$  is defined by

$$f(x) = \frac{3-2x}{x-5}, \quad x \in \mathbb{R}, \quad x \neq 5$$

(a) Find  $f^{-1}(x)$ .




Figure 2

The function  $g$  has domain  $-1 \leq x \leq 8$ , and is linear from  $(-1, -9)$  to  $(2, 0)$  and from  $(2, 0)$  to  $(8, 4)$ . Figure 2 shows a sketch of the graph of  $y = g(x)$ .

(b) Write down the range of  $g$ .

(1)

(c) Find  $gg(2)$ .

(2)

(d) Find  $fg(8)$ .

(2)

(e) On separate diagrams, sketch the graph with equation

(i)  $y = |g(x)|$ ,

(ii)  $y = g^{-1}(x)$ .

Show on each sketch the coordinates of each point at which the graph meets or cuts the axes.

(4)

(f) State the domain of the inverse function  $g^{-1}$ .

(1)

$$6) \quad x = \frac{3-2y}{y-5} \Rightarrow xy - 5x = 3 - 2y \Rightarrow xy + 2y = 5x + 3$$



$$\Rightarrow (x+2)y = 5x+3 \Rightarrow y = f^{-1}(x) = \frac{5x+3}{x+2}$$

b)  $-9 \leq g(x) \leq 4 \quad g(x) \in \mathbb{R}$ .

c)  $gg(2) = g(0) = -6$

d)  $fg(8) = f(4) = \frac{3-8}{4-5} = \frac{-5}{-1} = 5$

e)



f) domain  $g^{-1} = \text{range } g \Rightarrow -9 \leq x \leq 4 \quad x \in \mathbb{R}$



7. The curve  $C$  has equation

$$y = \frac{3 + \sin 2x}{2 + \cos 2x}$$

(a) Show that

$$\frac{dy}{dx} = \frac{6 \sin 2x + 4 \cos 2x + 2}{(2 + \cos 2x)^2}$$

(4)

(b) Find an equation of the tangent to  $C$  at the point on  $C$  where  $x = \frac{\pi}{2}$ .

Write your answer in the form  $y = ax + b$ , where  $a$  and  $b$  are exact constants.

(4)

$$a) \frac{dy}{dx} = \frac{(3 + \sin 2x)'(2 + \cos 2x) - (3 + \sin 2x)(2 + \cos 2x)'}{(2 + \cos 2x)^2}$$

$$= \frac{2 \cos 2x (2 + \cos 2x) - (3 + \sin 2x)(-2 \sin 2x)}{(2 + \cos 2x)^2}$$

$$= \frac{4 \cos 2x + 2 \cos^2 2x + 6 \sin 2x + 2 \sin^2 2x}{(2 + \cos 2x)^2}$$

$$= \frac{6 \sin 2x + 4 \cos 2x + 2(\sin^2 2x + \cos^2 2x)}{(2 + \cos 2x)^2} \quad \overset{=1}{}$$

$$= \frac{6 \sin 2x + 4 \cos 2x + 2}{(2 + \cos 2x)^2} \quad \#$$

$$b) x = \frac{\pi}{2}; y = 3; M_t = -2$$

$$y - 3 = -2(x - \frac{\pi}{2}) \Rightarrow y = -2x + \pi + 3$$

8. (a) Given that

$$\frac{d}{dx}(\cos x) = -\sin x$$

show that  $\frac{d}{dx}(\sec x) = \sec x \tan x$ .

(3)

Given that

$$x = \sec 2y$$

(b) find  $\frac{dx}{dy}$  in terms of  $y$ .

(2)

(c) Hence find  $\frac{dy}{dx}$  in terms of  $x$ .

(4)

$$a) \frac{d}{dx}(\sec x) = \frac{d}{dx}\left(\frac{1}{\cos x}\right) = \frac{d}{dx}(\cos x)^{-1}$$

$$= -(\cos x)^{-2} \times -\sin x = \frac{\sin x}{\cos^2 x} = \frac{1}{\cos x} \times \frac{\sin x}{\cos x}$$

$$= \sec x \tan x \quad \#$$

$$b) x = \sec 2y$$

$$\frac{dx}{dy} = 2 \sec 2y \tan 2y$$

$$c) \frac{dy}{dx} = \frac{1}{2 \sec 2y \tan 2y}$$

$$\sec 2y = x$$

$$\tan^2 2y + 1 = \sec^2 2y$$

$$\frac{dy}{dx} = \frac{1}{2x\sqrt{x^2-1}}$$

$$\tan^2 2y = x^2 - 1$$

$$\tan 2y = \sqrt{x^2 - 1}$$