

C3 June 12

PMT

1. Express

$$\frac{2(3x+2)}{9x^2-4} - \frac{2}{3x+1}$$

as a single fraction in its simplest form.

(4)

$$\frac{2(3x+2)}{(3x-2)(3x+2)} - \frac{2}{3x+1}$$

$$= \frac{2(3x+1) - 2(3x-2)}{(3x-2)(3x+1)} = \frac{6}{(3x-2)(3x+1)}$$

2.

$$f(x) = x^3 + 3x^2 + 4x - 12$$

(a) Show that the equation $f(x) = 0$ can be written as

$$x = \sqrt{\left(\frac{4(3-x)}{3+x}\right)}, \quad x \neq -3 \quad (3)$$

The equation $x^3 + 3x^2 + 4x - 12 = 0$ has a single root which is between 1 and 2

(b) Use the iteration formula

$$x_{n+1} = \sqrt{\left(\frac{4(3-x_n)}{3+x_n}\right)}, \quad n \geq 0$$

with $x_0 = 1$ to find, to 2 decimal places, the value of x_1 , x_2 and x_3 .

(3)

The root of $f(x) = 0$ is α .

(c) By choosing a suitable interval, prove that $\alpha = 1.272$ to 3 decimal places.

(3)

$$a) \quad x^3 + 3x^2 + 4x - 12 = 0$$

$$x^3 + 3x^2 = 12 - 4x$$

$$x^2(x+3) = 4(3-x)$$

$$x^2 = \frac{4(3-x)}{(x+3)} \Rightarrow x = \sqrt{\frac{4(3-x)}{3+x}} \quad \#$$

$$b) \quad x_0 = 1; \quad x_1 = 1.41; \quad x_2 = 1.20; \quad x_3 = 1.31$$

$$c) \quad f(1.2715) = -0.0082 < 0$$

$$f(1.2725) = 0.0083 > 0$$

by sign change rule $\alpha \in (1.2715, 1.2725)$

$$\therefore \alpha = 1.272 \text{ (3dp)}$$

3.




Figure 1

Figure 1 shows a sketch of the curve C which has equation

$$y = e^{x\sqrt{3}} \sin 3x, \quad -\frac{\pi}{3} \leq x \leq \frac{\pi}{3}$$

- (a) Find the x coordinate of the turning point P on C , for which $x > 0$
Give your answer as a multiple of π .

(6)

- (b) Find an equation of the normal to C at the point where $x = 0$

(3)

$$\frac{d}{dx}(uv) = v u' + u v' \quad u = e^{x\sqrt{3}} \quad v = \sin 3x$$

$$u' = \sqrt{3} e^{x\sqrt{3}} \quad v' = 3 \cos 3x$$

$$\frac{dy}{dx} = \sqrt{3} e^{x\sqrt{3}} \sin 3x + 3 e^{x\sqrt{3}} \cos 3x$$

$$= e^{x\sqrt{3}} (\sqrt{3} \sin 3x + 3 \cos 3x)$$

$$\text{at TP } \frac{dy}{dx} = 0 \Rightarrow \sqrt{3} \sin 3x = -3 \cos 3x$$

$$\therefore \tan 3x = -\frac{3}{\sqrt{3}}$$

$$\therefore 3x = -\frac{\pi}{3}, \frac{2\pi}{3}, \frac{5\pi}{3}, \dots \Rightarrow x = -\frac{\pi}{9}, \frac{2\pi}{9}, \frac{5\pi}{9}$$

$$\therefore x = \frac{2\pi}{9} \quad y = e^{\frac{2\sqrt{3}\pi}{9}} \sin\left(3 \times \frac{2\pi}{9}\right) = \frac{\sqrt{3}}{2} e^{\frac{2\sqrt{3}\pi}{9}}$$

(y not asked for!!)

$$b) \quad x=0 \quad y=0$$

$$\left. \frac{dy}{dx} \right|_{x=0} = e^0(0+3) = 3 \quad m_t = 3 \Rightarrow m_n = -\frac{1}{3}$$

$$\therefore y - 0 = -\frac{1}{3}(x - 0) \quad \Rightarrow \quad y = -\frac{1}{3}x$$

4.




Figure 2

Figure 2 shows part of the curve with equation $y = f(x)$
The curve passes through the points $P(-1.5, 0)$ and $Q(0, 5)$ as shown.

On separate diagrams, sketch the curve with equation

(a) $y = |f(x)|$ (2)

(b) $y = f(|x|)$ (2)


(c) $y = 2f(3x)$ (3)

Indicate clearly on each sketch the coordinates of the points at which the curve crosses or meets the axes.

a) $|f(x)|$

b) $f(|x|)$

c) $2f(3x)$
 $\uparrow 2 \rightarrow 3 \leftarrow$



5. (a) Express $4\operatorname{cosec}^2 2\theta - \operatorname{cosec}^2 \theta$ in terms of $\sin \theta$ and $\cos \theta$.

(2)

(b) Hence show that

$$4\operatorname{cosec}^2 2\theta - \operatorname{cosec}^2 \theta = \sec^2 \theta$$

(4)

(c) Hence or otherwise solve, for $0 < \theta < \pi$,

$$4\operatorname{cosec}^2 2\theta - \operatorname{cosec}^2 \theta = 4$$

giving your answers in terms of π .

(3)

$$\frac{4}{\sin^2 2\theta} - \frac{1}{\sin^2 \theta} = \frac{4}{(2\sin\theta\cos\theta)^2} - \frac{1}{\sin^2 \theta}$$

$$= \frac{4}{4\sin^2\theta\cos^2\theta} - \frac{1}{\sin^2\theta} \Rightarrow \frac{1}{\sin^2\theta\cos^2\theta} - \frac{1}{\sin^2\theta}$$

$$\text{b) } \frac{1 - \cos^2\theta}{\sin^2\theta\cos^2\theta} = \frac{\sin^2\theta}{\sin^2\theta\cos^2\theta} = \frac{1}{\cos^2\theta} = \sec^2\theta \quad \#$$

$$\text{c) } \sec^2\theta = 4 \Rightarrow \frac{1}{\cos^2\theta} = 4 \Rightarrow \cos^2\theta = \frac{1}{4}$$

$$\cos\theta = \frac{1}{2}$$

$$\cos\theta = -\frac{1}{2}$$

$$\theta = \frac{\pi}{3}$$

$$\theta = \frac{2\pi}{3}$$

6. The functions f and g are defined by

$$f: x \mapsto e^x + 2, \quad x \in \mathbb{R}$$

$$g: x \mapsto \ln x, \quad x > 0$$

(a) State the range of f .

(1)

(b) Find $fg(x)$, giving your answer in its simplest form.

(2)

(c) Find the exact value of x for which $f(2x+3) = 6$

(4)

(d) Find f^{-1} , the inverse function of f , stating its domain.

(3)

(e) On the same axes sketch the curves with equation $y = f(x)$ and $y = f^{-1}(x)$, giving the coordinates of all the points where the curves cross the axes.

(4)

$$a) \quad y = e^x + 2 \quad \text{range } y > 2$$

$$b) \quad fg(x) = f(\ln x) = e^{\ln x} + 2 = x + 2$$

$$c) \quad f(2x+3) = e^{2x+3} + 2 = 6$$


$$e^{2x+3} = 4 \Rightarrow 2x+3 = \ln 4 \Rightarrow x = \frac{-3 + \ln 4}{2}$$

$$d) \quad y = e^x + 2 \Rightarrow x = e^y + 2$$

$$\Rightarrow e^y = x - 2 \Rightarrow y = \ln(x - 2)$$

domain $x > 2$

e)



7. (a) Differentiate with respect to x ,

(i) $x^{\frac{1}{2}} \ln(3x)$

(ii) $\frac{1-10x}{(2x-1)^5}$, giving your answer in its simplest form.

(6)

(b) Given that $x = 3 \tan 2y$ find $\frac{dy}{dx}$ in terms of x .

(5)

$$\begin{aligned} \text{i) } u &= x^{\frac{1}{2}} & v &= \ln 3x & \frac{dy}{dx} &= \frac{1}{2}x^{-\frac{1}{2}} \ln 3x + \frac{x^{\frac{1}{2}}}{x} \\ u' &= \frac{1}{2}x^{-\frac{1}{2}} & v' &= \frac{3}{3x} = \frac{1}{x} & &= \frac{\ln 3x}{2x^{\frac{1}{2}}} + \frac{1}{\sqrt{x}} \end{aligned}$$

$$\begin{aligned} \text{ii) } u &= 1-10x & v &= (2x-1)^5 & \frac{d}{dx} \left(\frac{u}{v} \right) &= \frac{vu' - uv'}{v^2} \\ u' &= -10 & v' &= 10(2x-1)^4 \end{aligned}$$

$$\frac{dy}{dx} = \frac{-10(2x-1)^5 - 10(1-10x)(2x-1)^4}{(2x-1)^{10}}$$

$$= \frac{-10(2x-1) - 10(1-10x)}{(2x-1)^6} = \frac{80x}{(2x-1)^6}$$

$$\text{iii) } x = 3 \tan 2y$$

$$\frac{dx}{dy} = 6 \sec^2 2y \Rightarrow \frac{dy}{dx} = \frac{1}{6 \sec^2 2y}$$

$$\therefore \frac{dy}{dx} = \frac{1}{6} \cos^2 2y$$

$$\tan 2y = \frac{x}{3} \Rightarrow 2y = \arctan\left(\frac{x}{3}\right) \Rightarrow y = \frac{1}{2} \arctan\left(\frac{x}{3}\right)$$

$$\therefore \frac{dy}{dx} = \frac{1}{6} \cos^2 \left[\arctan\left(\frac{x}{3}\right) \right]$$

$$\text{iii) } x = 3 \tan 2y$$

$$\frac{dx}{dy} = 6 \sec^2 2y \Rightarrow \frac{dy}{dx} = \frac{1}{6 \sec^2 2y}$$

$$\frac{\sin^2}{\cos^2} + \frac{\cos^2}{\cos^2} = \frac{1}{\cos^2} \Rightarrow \tan^2 + 1 = \sec^2$$

$$\frac{x}{3} = \tan 2y \Rightarrow \tan^2 2y = \frac{x^2}{9} \Rightarrow \sec^2 2y = 1 + \frac{x^2}{9}$$

$$\Rightarrow \frac{dy}{dx} = \frac{1}{6\left(1 + \frac{x^2}{9}\right)} = \frac{9}{6(9 + x^2)} = \frac{3}{2(9 + x^2)}$$

$$f(x) = 7 \cos 2x - 24 \sin 2x$$

Given that $f(x) = R \cos(2x + \alpha)$, where $R > 0$ and $0 < \alpha < 90^\circ$,

(a) find the value of R and the value of α .

(3)

(b) Hence solve the equation

$$7 \cos 2x - 24 \sin 2x = 12.5$$

for $0 \leq x < 180^\circ$, giving your answers to 1 decimal place.

(5)

(c) Express $14 \cos^2 x - 48 \sin x \cos x$ in the form $a \cos 2x + b \sin 2x + c$, where a , b , and c are constants to be found.

(2)

(d) Hence, using your answers to parts (a) and (c), deduce the maximum value of

$$14 \cos^2 x - 48 \sin x \cos x$$

(2)

$$\begin{aligned} R(\cos(2x + \alpha)) &= R(\cancel{\cos 2x} \cos \alpha - R \cancel{\sin 2x} \sin \alpha) \\ f(x) &= 7 \cancel{\cos 2x} - 24 \cancel{\sin 2x} \end{aligned}$$

$$\begin{aligned} \frac{R \sin \alpha}{R \cos \alpha} &= \frac{24}{7} \Rightarrow \tan \alpha = \frac{24}{7} & \alpha &= 73.74 \\ & & \alpha &= \underline{73.7} \end{aligned}$$

$$R = \sqrt{24^2 + 7^2} \qquad R = \underline{25}$$

$$f(x) = 25 \cos(2x + 73.74)$$

$$b) \quad 25 \cos(2x + 73.74) = 12.5$$

$$\Rightarrow \cos(2x + 73.74) = \frac{1}{2}$$

$$\Rightarrow 2x + 73.74 = 60^\circ, 300^\circ, 420^\circ$$

$$2x = -13.74, 226.26, 346.26$$

$$\therefore x = \quad \quad \quad , \underline{113.1} \quad ; \quad 113.1$$

$$b) \quad 14 \cos^2 x - 7 - 48 \sin x \cos x + 7$$

$$7(2 \cos^2 x - 1) - 24(2 \sin x \cos x) + 7$$

$$7 \cos 2x - 24 \sin 2x + 7$$

$$d) \quad 25 \cos(2x + 73.74) + 7$$

$$\text{max value} = 25 + 7 = \underline{\underline{32}}$$