




Mathematics in Education and Industry

MEI STRUCTURED MATHEMATICS

METHODS OF ADVANCED MATHEMATICS, C3

Practice Paper C3-A


MARK SCHEME

Qu		Answer	Mark	Comment
Section A				
1		Product of two numbers, one of which is even is always even. Two consecutive numbers contain an even number. <i>OR</i> acceptable alternatives	B1 B1 2	
2		$y = \sqrt{1+x^3}$ Let $u = 1+x^3 \Rightarrow \frac{du}{dx} = 3x^2$ $y = u^{\frac{1}{2}} \Rightarrow \frac{dy}{du} = \frac{1}{2}u^{-\frac{1}{2}}$ $\Rightarrow \frac{dy}{dx} = \frac{dy}{du} \times \frac{du}{dx} = \frac{1}{2}u^{-\frac{1}{2}} \times 3x^2 = \frac{3}{2} \frac{x^2}{\sqrt{1+x^3}}$	M1 A1 A1 3	Chain rule $\frac{dy}{du}$ Answer
3	(i)	Substitute: $0 = a\ln 2b \Rightarrow \ln 2b = 0 \Rightarrow 2b = 1 \Rightarrow b = \frac{1}{2}$ $1 = a\ln 2 \Rightarrow a = \frac{1}{\ln 2}$	M1 A1 A1 3	
	(ii)	$ a\ln(bx) < 2 \Rightarrow \left \frac{\ln \frac{1}{2}x}{\ln 2} \right < 2 \Rightarrow \left \ln \frac{1}{2}x \right < 2\ln 2$ $\Rightarrow -2\ln 2 < \ln \frac{1}{2}x < 2\ln 2$ $\Rightarrow \ln \frac{1}{4} < \ln \frac{1}{2}x < \ln 4$ $\Rightarrow \frac{1}{4} < \frac{1}{2}x < 4 \Rightarrow \frac{1}{2} < x < 8$	M1 M1 A1 A1 4	Modulus Powers of logs
4	(i)	$y = axe^{-x} \Rightarrow \frac{dy}{dx} = ae^{-x} - axe^{-x} = ae^{-x}(1-x)$ $\frac{dy}{dx} = 0 \Rightarrow x = 1$ only at $\left(1, \frac{a}{e}\right)$ $\frac{d^2y}{dx^2} = -ae^{-x}(1-x) - ae^{-x}$: When $x = 1, \frac{d^2y}{dx^2} < 0$ \Rightarrow Maximum	M1 A1 M1 A1 B1 5	Product = 0 or any equivalent argument
	(ii)		B1 B1 2	For curve for stationary point

5	$\int_2^3 xe^{2x} dx \quad u = x, \frac{dv}{dx} = e^{2x}$ $\frac{du}{dx} = 1, v = \frac{1}{2}e^{2x}$ $= \left[\frac{1}{2}xe^{2x} \right]_2^3 - \frac{1}{2} \int_2^3 e^{2x} dx = \left[\frac{1}{2}xe^{2x} - \frac{1}{4}e^{2x} \right]_2^3$ $= \frac{5}{4}e^6 - \frac{3}{4}e^4 = 463.3$	M1 A1 M1 A1 A1	5	Choice of u
6	$\frac{d}{dx}(x \ln x) = \ln x + x \times \frac{1}{x} = \ln x + 1$ $\Rightarrow x \ln x = \int (\ln x + 1) dx = \int \ln x dx + x$ $\Rightarrow \int_2^3 \ln x dx = [x \ln x - x]_2^3 = (3 \ln 3 - 3) - (2 \ln 2 - 2)$ $= 3 \ln 3 - 2 \ln 2 - 1$ $= \ln \frac{27}{4} - 1$	M1 A1 M1 A1 M1 A1	6	Product Integrand limits
7	<p>(i) $x = 5 \sin \frac{\pi}{2} - 4 \cos \frac{\pi}{2} = 5$</p> <p>(ii) $\frac{dx}{d\theta} = 5 \cos \theta + 4 \sin \theta$: When $\theta = \frac{\pi}{2}$, $\frac{dx}{d\theta} = 4$ $\frac{dx}{dt} = \frac{dx}{d\theta} \times \frac{d\theta}{dt} = 4 \times 0.1 = 0.4$</p>	B1 1 M1 A1 A1 M1 E1	1 4 5	

Section B

8	<p>(i) $f(x) = \frac{x}{x^2 + 1} \Rightarrow f'(x) = \frac{(x^2 + 1).1 - x.2x}{(x^2 + 1)^2}$</p> $= \frac{1 - x^2}{(x^2 + 1)^2}$	M1 A1 E1	3	Formula Middle section answer
	<p>(ii) $f(x) = \frac{1 - x^2}{(x^2 + 1)^2} = 0 \Rightarrow 1 - x^2 = 0 \Rightarrow x = \pm 1$</p> <p>When $x = 1$, $f(x) = \frac{1}{1+1} = \frac{1}{2}$ i.e. $\left(1, \frac{1}{2}\right)$</p> <p>Other stationary point is when $x = -1$, $f(x) = \frac{-1}{1+1} = -\frac{1}{2}$</p> <p>i.e. $\left(-1, -\frac{1}{2}\right)$</p>	E1 B1	2	Substitute Find $f(x)$
	<p>(iii) The graph is odd.</p> $f(-x) = \frac{-x}{(-x)^2 + 1} = -\frac{x}{x^2 + 1} = -f(x)$	B1 B1 2		

	(iv)	$u = x^2 \Rightarrow du = 2x dx$ When $x = 1, u = 1$ When $x = 4, u = 16$ $\Rightarrow \int_1^4 \frac{x}{x^2 + 1} dx = \int_1^{16} \frac{1}{u+1} \cdot \frac{1}{2} du$ $\Rightarrow a = 1, b = 16, k = \frac{1}{2}$	M1 B1 B1 A1 A1	5
	(v)	Because the function is odd the area in $[-1,0]$ is equal in magnitude but opposite in sign to the area in $[0,1]$ So shaded area = $\int_1^4 \frac{x}{x^2 + 1} dx + 2 \int_0^1 \frac{x}{x^2 + 1} dx$ $= \int_1^{16} \frac{1}{u+1} \cdot \frac{1}{2} du + 2 \int_0^1 \frac{1}{u+1} \cdot \frac{1}{2} du$ $= \frac{1}{2} [\ln(u+1)]_1^{16} + 2 \frac{1}{2} [\ln(u+1)]_0^1$ $= \frac{1}{2} \ln \frac{17}{2} + 2 \frac{1}{2} \ln 2 = \frac{1}{2} \left(\ln \frac{17}{2} + \ln 4 \right) = \frac{1}{2} \ln 34 \square 1.763$	B1 M1 A1 M1 A1 A1	6
9	(i)	$\sqrt{y} = 1 - \sqrt{x}$ $\Rightarrow y = (1 - \sqrt{x})^2 = 1 + x - 2\sqrt{x}$ This is undefined for $x < 0$ but is defined for $x > 1$ so that part is missing. 	M1 A1 A1 B1 B1 B1 B1	Or: squaring introduces also the negative arm $x < 0$ $x > 0$ Extra range shape 7
	(ii)	A is $\sqrt{-x} + \sqrt{y} = 1$ for $-1 \leq x \leq 0$ B is $\sqrt{-x} + \sqrt{-y} = 1$ for $-1 \leq x \leq 0$ C is $\sqrt{x} + \sqrt{-y} = 1$ for $0 \leq x \leq 1$	B1 B1 B1 B1 B1 B1 B1	6
	(iii)	Area in Fig. 9.1 is $\int_0^1 (1 + x - 2\sqrt{x}) dx$ $= \left[x + \frac{1}{2}x^2 - \frac{4}{3}x^{\frac{3}{2}} \right]_0^1 = 1 + \frac{1}{2} - \frac{4}{3} = \frac{1}{6}$ \Rightarrow Area of shape is $\frac{1}{6} + \frac{1}{6} + 2 \times \frac{1}{6} + 2 \times \frac{1}{6} = 1$	M1 A1 A1 M1 A1	5