



MEI


Mathematics in Education and Industry

MEI STRUCTURED MATHEMATICS

METHODS OF ADVANCED MATHEMATICS, C3





Practice Paper C3-B

MARK SCHEME

Qu	Answer	Mark	Comment
Section A			
1	<p>Call the numbers n, $n + 1$ and $n + 2$ At least one of the numbers is even, and so the product is a multiple of 2. If n is a multiple of 3 then so is the product. If $n = 3k + 1$ then $n + 2$ is a multiple of 3 If $n = 3k + 2$ then $n + 1$ is a multiple of 3.</p> <p>n must have one of the forms $3k$, $3k + 1$ or $3k + 2$. Therefore whichever it is one of the three numbers is a multiple of 3 and so the product is a multiple of 3. Since it is also a multiple of 2 it is a multiple of 6.</p>	B1 M1 M1 E1 4	Algebra Divisibility by 2 Divisibility by 3 conclusion
2	(i) 	B1 B1 2	Right part Left part
	(ii) Line $y = 5$ to be shown on graph. $-1 < x < 4$	M1 A1 2	
3	(i) $y = (x^2 + 3)^5$ Let $u = x^2 + 3 \Rightarrow \frac{du}{dx} = 2x$ $y = u^5 \Rightarrow \frac{dy}{du} = 5u^4$ $\Rightarrow \frac{dy}{dx} = \frac{dy}{du} \times \frac{du}{dx} = 5u^4 \times 2x = 10x(x^2 + 3)^4$	M1 A1 A1 3	Chain rule $\frac{dy}{du}$
	(ii) $y = \frac{\sin 2x}{x}$ Let $u = \sin 2x \Rightarrow \frac{du}{dx} = 2 \cos 2x$ $v = x \Rightarrow \frac{dv}{dx} = 1$ $\Rightarrow \frac{dy}{dx} = \frac{v \frac{du}{dx} - u \frac{dv}{dx}}{v^2} = \frac{2x \cos 2x - \sin 2x}{x^2}$	M1 A1 A1 3	Quotient rule
4	$y^2 = 5x - 4 \Rightarrow 2y \frac{dy}{dx} = 5 \Rightarrow \frac{dy}{dx} = \frac{5}{2y}$ When $x = 8$, $y^2 = 36 \Rightarrow y = \pm 6$ \Rightarrow gradients = $\frac{5}{12}$ and $-\frac{5}{12}$	M1 A1 A1 A1 A1 5	

5		$x = x_0 e^{-3t} \Rightarrow e^{3t} = \frac{x_0}{x}$ $\Rightarrow 3t = \ln\left(\frac{x_0}{x}\right) \Rightarrow t = \frac{1}{3} \ln\left(\frac{x_0}{x}\right)$ $\Rightarrow t = \ln\left(\frac{x_0}{x}\right)^{\frac{1}{3}}$ <p>i.e. $a = x_0$, $b = \frac{1}{3}$</p>	M1 A1 A1 A1	Take logs or any equivalent method	4
6	(i)	$\int (2x-3)^7 dx. \quad \text{Let } u = 2x-3, \frac{du}{dx} = 2 \Rightarrow dx = \frac{1}{2} du$ $= \int \frac{1}{2} u^7 du = \frac{u^8}{2 \times 8} = \frac{1}{16} (2x-3)^8 + c$	M1 A1 A1	or B3 cao	3
	(ii)	<p>The substitution $u = x^2 + 1$ gives $\frac{du}{dx} = 2x$</p> $\Rightarrow \int_1^2 x(x^2 + 1)^3 dx = \int_2^5 \frac{1}{2} u^3 du$ $= \left[\frac{u^4}{8} \right]_2^5$ $= \frac{609}{8} (= 76\frac{1}{8})$	M1 A1 A1 A1 A1	Using sub Correct int Correct limits Int Ans	5
7	(i)	$f^2(x) = 4x$	B1		1
	(ii)	$fgh(x) = fg(x+2)$ $= f(x+2)^2$ $= 2(x+2)^2$	M1 A1 A1	correct order of functions	3
	(iii)	$y = h(x)$ $= x+2$ $\Rightarrow x = y-2$ $h^{-1}(x) = x-2$	B1		1

Section B				
8	(i)	$0 = (x+2)e^{-x}$ $\Rightarrow x = -2$ so $(-2,0)$ and $(0,2)$	B1 B1	2
	(ii)	$y = (x+2)e^{-x}$ $\Rightarrow \frac{dy}{dx} = -e^{-x}(x+1) = 0 \Rightarrow x = -1$ SP is $(-1,e)$	M1 A1 M1 A1	Product rule = 0 4
	(iii)	$\Rightarrow \frac{d^2y}{dx^2} = xe^{-x}$ At $(-1,e)$ this is negative, so SP is a maximum.	M1 A1 A1	3
	(iv)		B1	1
	(v)	At $(0,2)$ gradient is -1 so gradient of normal is 1 Normal is $y = x + 2$. $y = x + 2, y = (x+2)e^{-x}$ $\Rightarrow 0 = (x+2)(1 - e^{-x})$ $\Rightarrow x = -2$ (or 0) New intersection point is $(-2,0)$.	B1 M1 A1	3
	(vi)	Required area is $\int_1^3 (x+2)e^{-x} dx$ $= [-e^{-x}(x+2)]_1^3 + \int_1^3 e^{-x} dx$ $= [-e^{-x}(x+2)]_1^3 + [-e^{-x}]_1^3$ $= \frac{-6}{e^3} + \frac{4}{e}$	B1 M1 A1 A1 A1	or equivalent 5

9	(i) (A)		The transformation is a stretch with the x -axis invariant and of scale factor 2.	B1 B1 2	Same orientation y values doubled
	(i) (B)		The transformation is a reflection in the y -axis.	B1 B2 3	same shape Inversion
	(i) (C)		The transformation is a translation of 2 units parallel to the x -axis, ie $\begin{pmatrix} 2 \\ 0 \end{pmatrix}$	B1 B2 3	Same shape Moved 2 to the right
	(ii)	There is a set of values of y (for example, $y = 1$) for which there are three corresponding values of x (so an inverse would be multivalued).	B1 B1 2		
	(iii)		B1 1		
	(iv)	$f(x) = x^2(x+2)$ $\Rightarrow f'(x) = 3x^2 + 4x$ So the gradient at (1,3) is 7. The gradient on the inverse (which is a reflection of the original in $y = x$) is therefore $^{-1}/7$.	M1 A1 M1 A1 4		
	(v)	The graph and its reflection must intersect on the axis of reflection, ie $y = x$, so solve $y = x, y = x^2(x+2)$ $\Rightarrow x = x^2(x+2)$ $\Rightarrow 0 = x(x^2 + 2x - 1)$ $\Rightarrow x = 0, -1 \pm \sqrt{2}$ The positive non-zero root is as given.	M1 M1 E1 3		