

4723 Core Mathematics 3

- 1 (i) State $y = \sec x$ B1
 (ii) State $y = \cot x$ B1
 (iii) State $y = \sin^{-1} x$ B1 3
3

- 2 Either: State or imply $\int \pi(2x-3)^4 dx$ B1 or unsimplified equiv
 Obtain integral of form $k(2x-3)^5$ M1 any constant k involving π or not
 Obtain $\frac{1}{10}(2x-3)^5$ or $\frac{1}{10}\pi(2x-3)^5$ A1
 Attempt evaluation using 0 and $\frac{3}{2}$ M1 subtraction correct way round
 Obtain $\frac{243}{10}\pi$ A1 5 or exact equiv
- Or: State or imply $\int \pi(2x-3)^4 dx$ B1 or unsimplified equiv
 Expand and obtain integral of order 5 M1 with at least three terms correct
 Ob'n $\frac{16}{5}x^5 - 24x^4 + 72x^3 - 108x^2 + 81x$ A1 with or without π
 Attempt evaluation using (0 and) $\frac{3}{2}$ M1
 Obtain $\frac{243}{10}\pi$ A1 (5) or exact equiv
5

- 3 (i) Attempt use of identity for $\sec^2 \alpha$ M1 using $\pm \tan^2 \alpha \pm 1$
 Obtain $1 + (m+2)^2 - (1+m^2)$ A1 absent brackets implied by subsequent correct working
 Obtain $4m + 4 = 16$ and hence $m = 3$ A1 3
-
- (ii) Attempt subn in identity for $\tan(\alpha + \beta)$ M1 using $\frac{\pm \tan \alpha \pm \tan \beta}{1 \pm \tan \alpha \tan \beta}$
 Obtain $\frac{5+3}{1-15}$ or $\frac{m+2+m}{1-m(m+2)}$ A1√ following their m
 Obtain $-\frac{4}{7}$ A1 3 or exact equiv
6

- 4 (i) Obtain $\frac{1}{3}e^{3x} + e^x$ B1
 Substitute to obtain $\frac{1}{3}e^{9a} + e^{3a} - \frac{1}{3}e^{3a} - e^a$ B1 or equiv
 Equate definite integral to 100 and attempt rearrangement M1 as far as $e^{9a} = \dots$
 Introduce natural logarithm M1 using correct process
 Obtain $a = \frac{1}{9} \ln(300 + 3e^a - 2e^{3a})$ A1 5 AG; necessary detail needed
-
- (ii) Obtain correct first iterate B1 allow for 4 dp rounded or truncated
 Show correct iteration process M1 with at least one more step
 Obtain at least three correct iterates in all A1 allowing recovery after error
 Obtain 0.6309 A1 4 following at least three correct steps; answer required to exactly 4 dp
 [0.6 \rightarrow 0.631269 \rightarrow 0.630884 \rightarrow 0.630889]
9

5 (i)	<u>Either:</u> Show correct process for comp'n	M1	correct way round and in terms of x
	Obtain $y = 3(3x + 7) - 2$	A1	or equiv
	Obtain $x = -\frac{19}{9}$	A1 3	or exact equiv; condone absence of $y = 0$
	<u>Or:</u> Use $fg(x) = 0$ to obtain $g(x) = \frac{2}{3}$	B1	
	Attempt solution of $g(x) = \frac{2}{3}$	M1	
	Obtain $x = -\frac{19}{9}$	A1 (3)	or exact equiv; condone absence of $y = 0$

(ii)	Attempt formation of one of the equations		
	$3x + 7 = \frac{x-7}{3}$ or $3x + 7 = x$ or $\frac{x-7}{3} = x$	M1	or equiv
	Obtain $x = -\frac{7}{2}$	A1	or equiv
	Obtain $y = -\frac{7}{2}$	A1√ 3	or equiv; following their value of x

(iii)	Attempt solution of modulus equation	M1	squaring both sides to obtain 3-term quadratics or forming linear equation with signs of $3x$ different on each side
	Obtain $-12x + 4 = 42x + 49$ or $3x - 2 = -3x - 7$	A1	or equiv
	Obtain $x = -\frac{5}{6}$	A1	or exact equiv; as final answer
	Obtain $y = \frac{9}{2}$	A1 4	or equiv; and no other pair of answers
10			

6 (i)	Obtain derivative $k(37 + 10y - 2y^2)^{-\frac{1}{2}}f(y)$	M1	any constant k ; any linear function for f
	Obtain $\frac{1}{2}(10 - 4y)(37 + 10y - 2y^2)^{-\frac{1}{2}}$	A1 2	or equiv

(ii)	<u>Either:</u> Sub'te $y = 3$ in expression for $\frac{dx}{dy}$	*M1	
	Take reciprocal of expression/value	*M1	and without change of sign
	Obtain -7 for gradient of tangent	A1	
	Attempt equation of tangent	M1	dep *M *M
	Obtain $y = -7x + 52$	A1 5	and no second equation
	<u>Or:</u> Sub'te $y = 3$ in expression for $\frac{dx}{dy}$	M1	
	Attempt formation of eq'n $x = m'y + c$	M1	where m' is attempt at $\frac{dx}{dy}$
	Obtain $x - 7 = -\frac{1}{7}(y - 3)$	A1	or equiv
	Attempt rearrangement to required form	M1	
	Obtain $y = -7x + 52$	A1 (5)	and no second equation
7			

7 (i)	State $R = 10$ Attempt to find value of α Obtain 36.9 or $\tan^{-1} \frac{3}{4}$	B1 M1 A1 3	or equiv implied by correct answer or its complement; allow sin/cos muddles or greater accuracy 36.8699...

(ii)(a)	Show correct process for finding one angle Obtain $(64.16 + 36.87)$ and hence) 101 Show correct process for finding second angle Obtain $(115.84 + 36.87)$ and hence) 153	M1 A1 M1 A1√ 4	 or greater accuracy 101.027... following their value of α ; or greater accuracy 152.711...; and no other between 0 and 360

(b)	Recognise link with part (i) Use fact that maximum and minimum values of sine are 1 and -1 Obtain 60	M1 M1 A1 3 10	signalled by $40 \dots - 20 \dots$ may be implied; or equiv

8 (i)	Refer to translation and stretch State translation in x direction by 6 State stretch in y direction by 2 [SC: if M0 but one transformation completely correct, give B1]	M1 A1 A1 3	in either order; allow here equiv informal terms such as 'move', ... or equiv; now with correct terminology or equiv; now with correct terminology

(ii)	State $2 \ln(x - 6) = \ln x$ Show correct use of logarithm property Attempt solution of 3-term quadratic Obtain 9 only	B1 *M1 M1 A1 4	or $2 \ln(a - 6) = \ln a$ or equiv dep *M following correct solution of equation

(iii)	Attempt evaluation of form $k(y_0 + 4y_1 + y_2)$ Obtain $\frac{1}{3} \times (2 \ln 1 + 8 \ln 2 + 2 \ln 3)$ Obtain 2.58	M1 A1 A1 3 10	any constant k ; maybe with $y_0 = 0$ implied or equiv or greater accuracy 2.5808...

9 (a)	Attempt use of quotient rule Obtain $\frac{(kx^2 + 1)2kx - (kx^2 - 1)2kx}{(kx^2 + 1)^2}$ Obtain correct simplified numerator $4kx$ Equate numerator of first derivative to zero State $x = 0$ <u>or</u> refer to $4kx$ being linear <u>or</u> observe that, with $k \neq 0$, only one sol'n	*M1 A1 A1 M1 A1√ 5	or equiv; allow numerator wrong way round and denominator errors or equiv; with absent brackets implied by subsequent correct working dep *M AG or equiv; following numerator of form $k'kx = 0$, any constant k'

4723

Mark Scheme

June 2009

(b)	Attempt use of product rule	*M1	
	Obtain $me^{mx}(x^2 + mx) + e^{mx}(2x + m)$	A1	or equiv
	Equate to zero and either factorise with factor e^{mx} or divide through by e^{mx}	M1	dep *M
	Obtain $mx^2 + (m^2 + 2)x + m = 0$ or equiv		
	and observe that e^{mx} cannot be zero	A1	
	Attempt use of discriminant	M1	using correct $b^2 - 4ac$ with their a, b, c
	Simplify to obtain $m^4 + 4$	A1	or equiv
	Observe that this is positive for all m and hence two roots	A1	7 or equiv; AG
			12