

Core Mathematics 3 Paper D

1. (i) Show that

$$\sin(x + 30)^\circ + \sin(x - 30)^\circ \equiv a \sin x^\circ,$$

where a is a constant to be found. [3]

- (ii) Hence find the exact value of $\sin 75^\circ + \sin 15^\circ$, giving your answer in the form $b\sqrt{6}$. [3]

2. Solve each equation, giving your answers in exact form.

(i) $\ln(2x - 3) = 1$ [2]

(ii) $3e^y + 5e^{-y} = 16$ [5]

3. The curve C has the equation $y = 2e^x - 6 \ln x$ and passes through the point P with x -coordinate 1.

- (i) Find an equation for the tangent to C at P . [4]

The tangent to C at P meets the coordinate axes at the points Q and R .


- (ii) Show that the area of triangle OQR , where O is the origin, is $\frac{9}{3-e}$. [4]

4. The finite region R is bounded by the curve with equation $y = \frac{1}{2x-1}$, the x -axis and the lines $x = 1$ and $x = 2$.

- (i) Find the exact area of R . [4]

- (ii) Show that the volume of the solid formed when R is rotated through four right angles about the x -axis is $\frac{1}{3}\pi$. [4]

5.



The diagram shows the graph of $y = f(x)$. The graph has a minimum at $(\frac{\pi}{2}, -1)$, a maximum at $(\frac{3\pi}{2}, -5)$ and an asymptote with equation $x = \pi$.

(i) Showing the coordinates of any stationary points, sketch the graph of $y = |f(x)|$. [2]

Given that

$$f: x \rightarrow a + b \operatorname{cosec} x, \quad x \in \mathbb{R}, \quad 0 < x < 2\pi, \quad x \neq \pi,$$

(ii) find the values of the constants a and b , [3]

(iii) find, to 2 decimal places, the x -coordinates of the points where the graph of $y = f(x)$ crosses the x -axis. [3]

6. (i) Prove the identity

$$2 \cot 2x + \tan x \equiv \cot x, \quad x \neq \frac{n}{2}\pi, \quad n \in \mathbb{Z}. \quad [5]$$

(ii) Solve, for $0 \leq x < \pi$, the equation

$$2 \cot 2x + \tan x = \operatorname{cosec}^2 x - 7,$$

giving your answers to 2 decimal places. [6]

Turn over

7. The function f is defined by

$$f : x \rightarrow 3e^{x-1}, \quad x \in \mathbb{R}.$$

(i) State the range of f . [1]

(ii) Find an expression for $f^{-1}(x)$ and state its domain. [3]

The function g is defined by

$$g : x \rightarrow 5x - 2, \quad x \in \mathbb{R}.$$

Find, in terms of e ,

(iii) the value of $gf(\ln 2)$, [3]

(iv) the solution of the equation

$$f^{-1}g(x) = 4. \quad [4]$$

8. A curve has the equation $y = x^2 - \sqrt{4 + \ln x}$.

(i) Show that the tangent to the curve at the point where $x = 1$ has the equation

$$7x - 4y = 11. \quad [5]$$

The curve has a stationary point with x -coordinate α .

(ii) Show that $0.3 < \alpha < 0.4$ [3]

(iii) Show that α is a solution of the equation

$$x = \frac{1}{2}(4 + \ln x)^{-\frac{1}{4}}. \quad [2]$$

(iv) Use the iterative formula

$$x_{n+1} = \frac{1}{2}(4 + \ln x_n)^{-\frac{1}{4}},$$

with $x_0 = 0.35$, to find α correct to 5 decimal places.

You should show the result of each iteration. [3]