

# Core Mathematics 3 Paper J

1. Use Simpson's rule with four strips to estimate the value of the integral


$$\int_0^3 e^{\cos x} dx. \quad [4]$$

2. Giving your answers to 1 decimal place, solve the equation

$$5 \tan^2 2\theta - 13 \sec 2\theta = 1,$$

for  $\theta$  in the interval  $0 \leq \theta \leq 360^\circ$ . [7]

- 3.



The diagram shows the curve  $y = f(x)$  which has a maximum point at  $(-3, 2)$  and a minimum point at  $(2, -4)$ .

- (a) Showing the coordinates of any stationary points, sketch on separate diagrams the graphs of

(i)  $y = |f(x)|$ , [2]

(ii)  $y = 3f(2x)$ . [3]

- (b) Write down the values of the constants  $a$  and  $b$  such that the curve with equation  $y = a + f(x + b)$  has a minimum point at the origin  $O$ . [2]

4. Find the values of  $x$  in the interval  $-180 < x < 180$  for which

$$\tan(x + 45)^\circ - \tan x^\circ = 4,$$

giving your answers to 1 decimal place. [7]

5. The finite region  $R$  is bounded by the curve with equation  $y = \sqrt[3]{3x-1}$ , the  $x$ -axis and the lines  $x = \frac{2}{3}$  and  $x = 3$ .
- (i) Find the area of  $R$ . [4]
- (ii) Find, in terms of  $\pi$ , the volume of the solid formed when  $R$  is rotated through four right angles about the  $x$ -axis. [4]

6. The functions  $f$  and  $g$  are defined by

$$f : x \rightarrow 1 - ax, \quad x \in \mathbb{R},$$

$$g : x \rightarrow x^2 + 2ax + 2, \quad x \in \mathbb{R},$$

where  $a$  is a constant.

Find, in terms of  $a$ ,

- (i) an expression for  $f^{-1}(x)$ , [2]
- (ii) the range of  $g$ . [3]

Given that  $gf(3) = 7$ ,

- (iii) find the two possible values of  $a$ . [4]

7. The curve with equation  $y = x^{\frac{5}{2}} \ln \frac{x}{4}$ ,  $x > 0$  crosses the  $x$ -axis at the point  $P$ .

- (i) Write down the coordinates of  $P$ . [1]

The normal to the curve at  $P$  crosses the  $y$ -axis at the point  $Q$ .

- (ii) Find the area of triangle  $OPQ$  where  $O$  is the origin. [6]

The curve has a stationary point at  $R$ .

- (iii) Find the  $x$ -coordinate of  $R$  in exact form. [2]

**Turn over**

8. (i) Solve the equation

$$\pi - 3 \cos^{-1} \theta = 0. \quad [2]$$

- (ii) Sketch on the same diagram the curves  $y = \cos^{-1}(x - 1)$ ,  $0 \leq x \leq 2$  and  $y = \sqrt{x+2}$ ,  $x \geq -2$ . [3]

Given that  $\alpha$  is the root of the equation

$$\cos^{-1}(x - 1) = \sqrt{x+2},$$

- (iii) show that  $0 < \alpha < 1$ , [2]

- (iv) use the iterative formula

$$x_{n+1} = 1 + \cos \sqrt{x_n + 2}$$

with  $x_0 = 1$  to find  $\alpha$  correct to 3 decimal places.

You should show the result of each iteration. [3]

9. The number of bacteria present in a culture at time  $t$  hours is modelled by the continuous variable  $N$  and the relationship

$$N = 2000e^{kt},$$

where  $k$  is a constant.

Given that when  $t = 3$ ,  $N = 18\,000$ , find

- (i) the value of  $k$  to 3 significant figures, [3]
- (ii) how long it takes for the number of bacteria present to double, giving your answer to the nearest minute, [4]
- (iii) the rate at which the number of bacteria is increasing when  $t = 3$ . [4]