

Exercise 1C

$$1 \quad \frac{x^3 + 2x^2 + 3x - 4}{x+1} \equiv Ax^2 + Bx + C + \frac{D}{x+1}$$

$$\begin{array}{r}
 \overline{) x^3 + 2x^2 + 3x - 4} \\
 \underline{x^3 + x^2} \\
 x^2 + 3x \\
 \underline{x^2 + x} \\
 2x - 4 \\
 \underline{2x + 2} \\
 -6
 \end{array}$$

$$\frac{x^3 + 2x^2 + 3x - 4}{x+1} \equiv x^2 + x + 2 - \frac{6}{x+1}$$

So $A = 1, B = 1, C = 2, D = -6$

2 Using algebraic long division:

$$\begin{array}{r}
 \overline{) 2x^3 + 3x^2 - 4x + 5} \\
 \underline{2x^3 + 6x^2} \\
 -3x^2 - 4x + 5 \\
 \underline{-3x^2 - 9x} \\
 5x + 5 \\
 \underline{5x + 15} \\
 -10
 \end{array}$$

$$\frac{2x^3 + 3x^2 - 4x + 5}{x+3} = 2x^2 - 3x + 5 - \frac{10}{x+3}$$

So $a = 2, b = -3, c = 5$ and $d = -10$

3 Using algebraic long division:

$$\begin{array}{r}
 x^2 + 2x + 4 \\
 x - 2 \overline{) x^3 + 0x^2 + 0x - 8} \\
 \underline{x^3 - 2x^2} \\
 2x^2 + 0x \\
 \underline{2x^2 - 4x} \\
 4x - 8 \\
 \underline{4x - 8} \\
 0
 \end{array}$$

$$\text{So } \frac{x^3 - 8}{x - 2} = x^2 + 2x + 4$$

$$p = 1, q = 2 \text{ and } r = 4$$

4 Using algebraic long division:

$$\begin{array}{r}
 2 \\
 x^2 - 1 \overline{) 2x^2 + 4x + 5} \\
 \underline{2x^2 + 0x - 2} \\
 4x + 7
 \end{array}$$

$$\frac{2x^2 + 4x + 5}{x^2 - 1} \equiv 2 + \frac{4x + 7}{x^2 - 1}$$

$$\text{So } m = 2, n = 4 \text{ and } p = 7$$

5 Using algebraic long division:

$$\begin{array}{r}
 4x + 1 \\
 2x^2 + 2 \overline{) 8x^3 + 2x^2 + 0x + 5} \\
 \underline{8x^3 + 8x} \\
 2x^2 - 8x + 5 \\
 \underline{2x^2 + 2} \\
 -8x + 3
 \end{array}$$

$$8x^3 + 2x^2 + 5 \equiv (4x + 1)(2x^2 + 2) - 8x + 3$$

$$\text{So } A = 4, B = 1, C = -8 \text{ and } D = 3.$$

6 Using algebraic long division:

$$\begin{array}{r}
 4x-13 \\
 x^2+2x-1 \overline{)4x^3-5x^2+3x-14} \\
 \underline{4x^3+8x^2-4x} \\
 -13x^2+7x-14 \\
 \underline{-13x^2-26x+13} \\
 33x-27
 \end{array}$$

$$\frac{4x^3-5x^2+3x-14}{x^2+2x-1} \equiv 4x-13 + \frac{33x-27}{x^2+2x-1}$$

So $A = 4$, $B = -13$, $C = 33$ and $D = -27$.

7 Using algebraic long division:

$$\begin{array}{r}
 x^2+2 \\
 x^2+1 \overline{)x^4+3x^2-4} \\
 \underline{x^4+x^2} \\
 2x^2-4 \\
 \underline{2x^2+2} \\
 -6
 \end{array}$$

$$\frac{x^4+3x^2-4}{x^2+1} \equiv x^2+2 - \frac{6}{x^2+1}$$

So $p = 1$, $q = 0$, $r = 2$, $s = 0$ and $t = -6$.

8 Using algebraic long division:

$$\begin{array}{r}
 2x^2+x+1 \\
 x^2+x-2 \overline{)2x^4+3x^3-2x^2+4x-6} \\
 \underline{2x^4+2x^3-4x^2} \\
 x^3+2x^2+4x \\
 \underline{x^3+x^2-2x} \\
 x^2+6x-6 \\
 \underline{x^2+x-2} \\
 5x-4
 \end{array}$$

$$\frac{2x^4+3x^3-2x^2+4x-6}{x^2+x-2} \equiv 2x^2+x+1 + \frac{5x-4}{x^2+x-2}$$

So $a = 2$, $b = 1$, $c = 1$, $d = 5$ and $e = -4$.

$$9 \quad 3x^4 - 4x^3 - 8x^2 + 16x - 2 \equiv (Ax^2 + Bx + C)(x^2 - 3) + Dx + E$$

Compare coefficients of x^4 :

$$A = 3$$

Compare coefficients of x^3 :

$$B = -4$$

Compare coefficients of x^2 :

$$-8 = -3A + C$$

$$-8 = -9 + C \quad (\text{substituting } A = 3)$$

$$C = 1$$

Compare coefficients of x :

$$16 = -3B + D$$

$$16 = 12 + D \quad (\text{substituting } B = -4)$$

$$D = 4$$

Equate constant terms:

$$-2 = -3C + E$$

$$-2 = -3 + E \quad (\text{substituting } C = 1)$$

$$E = 1$$

$$\text{Hence } 3x^4 - 4x^3 - 8x^2 + 16x - 2 \equiv (3x^2 - 4x + 1)(x^2 - 3) + 4x + 1$$

Note: After lots of comparing coefficients, it is a good idea to check your answer by substituting a simple value of x into both sides of the identity to check that your answers are correct. For example,

Substitute $x = 1$ into LHS

$$\Rightarrow 3 - 4 - 8 + 16 - 2 = 5$$

Substitute $x = 1$ into RHS

$$\begin{aligned} \Rightarrow (3 - 4 + 1) \times (1 - 3) + 4 + 1 \\ = 0 \times -2 + 4 + 1 = 5 \end{aligned}$$

LHS = RHS, so you can be fairly sure the identity is correct.

$$10 \text{ a } \quad x^4 - 1 \equiv (x^2 - 1)(x^2 + 1)$$

$$\equiv (x - 1)(x + 1)(x^2 + 1)$$

$$\text{b } \quad \frac{x^4 - 1}{x + 1} \equiv \frac{(x - 1)(x + 1)(x^2 + 1)}{(x + 1)}$$

$$\equiv (x - 1)(x^2 + 1)$$

So $a = 1$, $b = -1$, $c = 1$, $d = 0$ and $e = 1$.