

Chapter review

1 a $\sin 40^\circ \cos 10^\circ - \cos 40^\circ \sin 10^\circ$

$$= \sin(40^\circ - 10^\circ) = \sin 30^\circ = \frac{1}{2}$$

b $\frac{1}{\sqrt{2}} \cos 15^\circ - \frac{1}{\sqrt{2}} \sin 15^\circ$

$$= \cos 45^\circ \cos 15^\circ - \sin 45^\circ \sin 15^\circ$$

$$= \cos(45^\circ + 15^\circ) = \cos 60^\circ = \frac{1}{2}$$


c $\frac{1 - \tan 15^\circ}{1 + \tan 15^\circ} = \frac{\tan 45^\circ - \tan 15^\circ}{1 + \tan 45^\circ \tan 15^\circ}$

$$= \tan(45^\circ - 15^\circ) = \tan 30^\circ = \frac{\sqrt{3}}{3}$$

2 As $\cos(x - y) = \sin y$
 $\cos x \cos y + \sin x \sin y = \sin y \quad (1)$

Draw a right-angled triangle,

where $\sin x = \frac{1}{\sqrt{5}}$



Using Pythagoras' theorem,

$$a^2 = (\sqrt{5})^2 - 1 = 4 \Rightarrow a = 2$$

So $\cos x = \frac{2}{\sqrt{5}}$

Substitute into (1):

$$\frac{2}{\sqrt{5}} \cos y + \frac{1}{\sqrt{5}} \sin y = \sin y$$

$$\Rightarrow 2 \cos y + \sin y = \sqrt{5} \sin y$$

$$\Rightarrow 2 \cos y = \sin y (\sqrt{5} - 1)$$

$$\Rightarrow \frac{2}{\sqrt{5} - 1} = \tan y \quad \left(\tan y = \frac{\sin y}{\cos y} \right)$$

$$\Rightarrow \tan y = \frac{2(\sqrt{5} + 1)}{(\sqrt{5} - 1)(\sqrt{5} + 1)}$$

$$= \frac{2(\sqrt{5} + 1)}{4} = \frac{\sqrt{5} + 1}{2}$$

3 a $\tan A = 2, \tan B = \frac{1}{3}$ since $y = \frac{1}{3}x - \frac{1}{3}$


b The angle required is $(A - B)$.

Using $\tan(A - B) = \frac{\tan A - \tan B}{1 + \tan A \tan B}$

$$= \frac{2 - \frac{1}{3}}{1 + 2 \times \frac{1}{3}} = \frac{\frac{5}{3}}{\frac{5}{3}} = 1$$

$$\Rightarrow A - B = 45^\circ$$

4



$$\begin{aligned} \text{Using } \frac{\sin B}{b} &= \frac{\sin C}{c} \\ \Rightarrow \frac{\sin(\theta-30^\circ)}{4} &= \frac{\sin(\theta+30^\circ)}{5} \\ \Rightarrow 5\sin(\theta-30^\circ) &= 4\sin(\theta+30^\circ) \\ \Rightarrow 5(\sin\theta\cos 30^\circ - \cos\theta\sin 30^\circ) &= 4(\sin\theta\cos 30^\circ + \cos\theta\sin 30^\circ) \\ \Rightarrow \sin\theta\cos 30^\circ &= 9\cos\theta\sin 30^\circ \\ \Rightarrow \frac{\sin\theta}{\cos\theta} &= 9 \frac{\sin 30^\circ}{\cos 30^\circ} = 9\tan 30^\circ \\ \Rightarrow \tan\theta &= 9 \times \frac{\sqrt{3}}{3} = 3\sqrt{3} \end{aligned}$$


- 5 As the three values are consecutive terms of an arithmetic progression,

$$\begin{aligned} \sin(\theta-30^\circ) - \sqrt{3}\cos\theta &= \sin\theta - \sin(\theta-30^\circ) \\ \Rightarrow 2\sin(\theta-30^\circ) &= \sin\theta + \sqrt{3}\cos\theta \\ \Rightarrow 2(\sin\theta\cos 30^\circ - \cos\theta\sin 30^\circ) &= \sin\theta + \sqrt{3}\cos\theta \\ \Rightarrow \sqrt{3}\sin\theta - \cos\theta &= \sin\theta + \sqrt{3}\cos\theta \\ \Rightarrow \sin\theta(\sqrt{3}-1) &= \cos\theta(\sqrt{3}+1) \\ \Rightarrow \tan\theta &= \frac{\sqrt{3}+1}{\sqrt{3}-1} \end{aligned}$$

$$\text{Calculator value is } \theta = \tan^{-1} \frac{\sqrt{3}+1}{\sqrt{3}-1} = 75^\circ$$

No other values as θ is acute.

6 a



$$\sin A = \frac{3}{5}, \cos A = \frac{4}{5} \quad \sin B = \frac{5}{13}, \cos B = \frac{12}{13}$$

$$\mathbf{i} \quad \sin(A+B) = \sin A \cos B + \cos A \sin B$$

$$= \frac{3}{5} \times \frac{12}{13} + \frac{4}{5} \times \frac{5}{13} = \frac{56}{65}$$

$$\mathbf{ii} \quad \tan 2B = \frac{2 \tan B}{1 - \tan^2 B}$$

$$\begin{aligned} &= \frac{2 \times \frac{5}{12}}{1 - \left(\frac{5}{12}\right)^2} = \frac{\frac{5}{6}}{\frac{119}{144}} \\ &= \frac{5}{6} \times \frac{144}{119} = \frac{120}{119} \end{aligned}$$

$$\mathbf{b} \quad \cos C = \cos(180^\circ - (A+B))$$

$$= -\cos(A+B)$$

$$= -(\cos A \cos B - \sin A \sin B)$$

$$= -\left(\frac{4}{5} \times \frac{12}{13} - \frac{3}{5} \times \frac{5}{13}\right)$$

$$= -\frac{33}{65}$$


$$\mathbf{7} \quad \mathbf{a} \quad \cos 2x \equiv 1 - 2\sin^2 x$$

$$= 1 - 2\left(\frac{2}{\sqrt{5}}\right)^2 = 1 - \frac{8}{5} = -\frac{3}{5}$$

$$\mathbf{b} \quad \cos 2y \equiv 2\cos^2 y - 1$$

$$= 2\left(\frac{3}{\sqrt{10}}\right)^2 - 1 = 2\left(\frac{9}{10}\right) - 1 = \frac{4}{5}$$

7 c



$$\begin{aligned} \text{i} \quad \tan(x+y) &= \frac{\tan x + \tan y}{1 - \tan x \tan y} \\ &= \frac{2 + \frac{1}{3}}{1 - \frac{2}{3}} = \frac{\frac{7}{3}}{\frac{1}{3}} = 7 \end{aligned}$$

$$\text{ii} \quad \tan(x-y) = \frac{\tan x - \tan y}{1 + \tan x \tan y} = \frac{\frac{5}{3}}{\frac{5}{3}} = 1$$

As x and y are acute, and $x > y$,

$x - y$ is acute

$$\text{So } x - y = \frac{\pi}{4} \quad \left(\text{it cannot be } \frac{5\pi}{4} \right)$$

$$\text{8 a} \quad \sin(x+y) \equiv \sin x \cos y + \cos x \sin y$$

$$= \frac{1}{2} + \frac{1}{3} = \frac{5}{6}$$

$$5 \sin(x-y) \equiv 5(\sin x \cos y - \cos x \sin y)$$

$$= 5\left(\frac{1}{2} - \frac{1}{3}\right) = 5 \times \frac{1}{6} = \frac{5}{6}$$

$$\text{b} \quad \frac{\sin x \cos y}{\cos x \sin y} = \frac{\frac{1}{2}}{\frac{1}{3}} = \frac{3}{2}$$

$$\Rightarrow \frac{\tan x}{\tan y} = \frac{3}{2}$$

$$\text{so } \tan x = \frac{3 \tan y}{2} = \frac{3k}{2}$$

$$\text{c} \quad \tan 2x = \frac{2 \tan x}{1 - \tan^2 x} = \frac{3k}{1 - \frac{9}{4}k^2}$$

$$= \frac{12k}{4 - 9k^2}$$

$$\text{9 a} \quad \sqrt{3} \sin 2\theta + 2 \sin^2 \theta = 1$$

$$\sqrt{3} \sin 2\theta = 1 - 2 \sin^2 \theta = \cos 2\theta$$

$$\frac{\sin 2\theta}{\cos 2\theta} = \frac{1}{\sqrt{3}} \Rightarrow \tan 2\theta = \frac{1}{\sqrt{3}}$$

$$\text{9 b} \quad \tan 2\theta = \frac{1}{\sqrt{3}}, \text{ for } 0 \leq 2\theta \leq 2\pi$$

$$2\theta = \frac{\pi}{6}, \frac{7\pi}{6} \Rightarrow \theta = \frac{\pi}{12}, \frac{7\pi}{12}$$

$$\text{10 a} \quad \cos 2\theta = 5 \sin \theta$$

$$\Rightarrow \cos 2\theta - 5 \sin \theta = 0$$

$$\Rightarrow 1 - 2 \sin^2 \theta - 5 \sin \theta = 0$$

$$\Rightarrow 2 \sin^2 \theta + 5 \sin \theta - 1 = 0$$

$$a = 2, b = 5 \text{ and } c = -1$$

$$\text{b} \quad 2 \sin^2 \theta + 5 \sin \theta - 1 = 0$$

Using the quadratic formula

$$\sin \theta = \frac{-5 \pm \sqrt{5^2 - 4(2)(-1)}}{2(2)}$$

$$= \frac{-5 \pm \sqrt{33}}{4}$$

$$\sin \theta = 0.1861, \text{ for } -\pi \leq \theta \leq \pi$$

$\sin \theta$ is positive so solutions in the first and second quadrants

$$\theta = \sin^{-1} 0.1861, \pi - \sin^{-1} 0.1861$$

$$\theta = 0.187, 2.954 \text{ (3 d.p.)}$$

$$\text{11 a} \quad \cos(x-60^\circ) = \cos x \cos 60^\circ + \sin x \sin 60^\circ$$

$$= \frac{1}{2} \cos x + \frac{\sqrt{3}}{2} \sin x$$

$$\text{So } 2 \sin x = \frac{1}{2} \cos x + \frac{\sqrt{3}}{2} \sin x$$

$$\Rightarrow \left(2 - \frac{\sqrt{3}}{2}\right) \sin x = \frac{1}{2} \cos x$$

$$\Rightarrow \tan x = \frac{\frac{1}{2}}{2 - \frac{\sqrt{3}}{2}} = \frac{1}{2} \times \frac{2}{4 - \sqrt{3}} = \frac{1}{4 - \sqrt{3}}$$

$$\text{b} \quad \tan x = \frac{1}{4 - \sqrt{3}} = 0.44 \text{ (2 d.p.)}, \text{ in the}$$

interval $0^\circ \leq \theta \leq 360^\circ$

$\tan x$ is positive so solutions in the first and third quadrants

$$x = 23.8^\circ, 203.8^\circ$$

Pure Mathematics 3

Solution Bank



12 a

$$\begin{aligned} \cos(x+20^\circ) &= \sin(90^\circ - 20^\circ - x) \\ &= \sin(70^\circ - x) \\ &= \sin 70^\circ \cos x - \cos 70^\circ \sin x \quad (1) \end{aligned}$$

$$\begin{aligned} 4\sin(70^\circ + x) &= 4\sin 70^\circ \cos x \\ &\quad + 4\cos 70^\circ \sin x \quad (2) \end{aligned}$$

As (1) = (2)

$$\begin{aligned} 4\sin 70^\circ \cos x + 4\cos 70^\circ \sin x \\ = \sin 70^\circ \cos x - \cos 70^\circ \sin x \end{aligned}$$

$$5\sin x \cos 70^\circ = -3\sin 70^\circ \cos x$$

$$\tan x = -\frac{3}{5} \tan 70^\circ$$

b $\tan x = -\frac{3}{5} \tan 70^\circ$, for $0^\circ \leq \theta \leq 180^\circ$


$\tan \theta$ is negative so the solution is in the second quadrant

$$x = 180^\circ + \tan^{-1}(-\frac{3}{5} \tan 70^\circ)$$

$$x = 180^\circ - \tan^{-1}(-1.648)$$

$$x = 180^\circ - (-58.8^\circ) = 121.2^\circ$$

- 13 a** Draw a right-angled triangle and find $\sin \alpha$ and $\cos \alpha$.



$$\Rightarrow \sin \alpha = \frac{3}{5}, \cos \alpha = \frac{4}{5}$$

$$\begin{aligned} 3\sin(\theta+\alpha) + 4\cos(\theta+\alpha) \\ \equiv 3(\sin \theta \cos \alpha + \cos \theta \sin \alpha) \\ + 4(\cos \theta \cos \alpha - \sin \theta \sin \alpha) \\ \equiv 3\left(\frac{4}{5}\sin \theta + \frac{3}{5}\cos \theta\right) \\ + 4\left(\frac{4}{5}\cos \theta - \frac{3}{5}\sin \theta\right) \\ \equiv \frac{12}{5}\sin \theta + \frac{9}{5}\cos \theta + \frac{16}{5}\cos \theta - \frac{12}{5}\sin \theta \\ \equiv \frac{25}{5}\cos \theta \equiv 5\cos \theta \end{aligned}$$

13 b

$$\begin{aligned} \cos(x+270^\circ) &\equiv \cos x^\circ \cos 270^\circ - \sin x^\circ \sin 270^\circ \\ &= (-0.8)(0) - (0.6)(-1) \\ &= 0 + 0.6 = 0.6 \end{aligned}$$

$$\begin{aligned} \cos(x+540^\circ) &\equiv \cos x^\circ \cos 540^\circ - \sin x^\circ \sin 540^\circ \\ &= (-0.8)(-1) - (0.6)(0) \\ &= 0.8 - 0 = 0.8 \end{aligned}$$

- 14 a** One example is sufficient to disprove a statement. Let $A = 60^\circ$, $B = 0^\circ$

$$\sec(A+B) = \sec(60^\circ + 0^\circ)$$

$$= \sec 60^\circ = \frac{1}{\cos 60^\circ} = 2$$

$$\sec A = \sec 60^\circ = \frac{1}{\cos 60^\circ} = 2$$

$$\sec B = \sec 0^\circ = \frac{1}{\cos 0^\circ} = 1$$

$$\text{So } \sec A + \sec B = 2 + 1 = 3$$

$$\text{So } \sec(60^\circ + 0^\circ) \neq \sec 60^\circ + \sec 0^\circ$$

$\Rightarrow \sin(A+B) = \sec A + \sec B$ is not true for all values of A, B .

b

$$\begin{aligned} \text{LHS} &\equiv \tan \theta + \cot \theta \\ &\equiv \frac{\sin \theta}{\cos \theta} + \frac{\cos \theta}{\sin \theta} \\ &\equiv \frac{\sin^2 \theta + \cos^2 \theta}{\sin \theta \cos \theta} \\ &\equiv \frac{1}{\frac{1}{2}\sin 2\theta} \end{aligned}$$

Using $\sin^2 \theta + \cos^2 \theta \equiv 1$, and
 $\sin 2\theta \equiv 2\sin \theta \cos \theta$

$$\begin{aligned} \text{So LHS} &\equiv \frac{2}{\sin 2\theta} \\ &\equiv 2 \operatorname{cosec} 2\theta \\ &\equiv \text{RHS} \end{aligned}$$

Pure Mathematics 3

Solution Bank



15 a Using $\tan 2\theta = \frac{2\tan\theta}{1-\tan^2\theta}$ with $\theta = \frac{\pi}{8}$

$$\Rightarrow \tan \frac{\pi}{4} = \frac{2\tan \frac{\pi}{8}}{1-\tan^2 \frac{\pi}{8}}$$

Let $t = \tan \frac{\pi}{8}$

So $1 = \frac{2t}{1-t^2}$

$$\Rightarrow 1-t^2 = 2t$$

$$\Rightarrow t^2 + 2t - 1 = 0$$

$$\Rightarrow t = \frac{-2 \pm \sqrt{8}}{2} = \frac{-2 \pm 2\sqrt{2}}{2}$$

$$= -1 \pm \sqrt{2}$$

As $\frac{\pi}{8}$ is acute, $\tan \frac{\pi}{8}$ is positive,

$$\text{so } \tan \frac{\pi}{8} = \sqrt{2} - 1$$

b $\tan \frac{3\pi}{8} = \tan \left(\frac{\pi}{4} + \frac{\pi}{8} \right) = \frac{\tan \frac{\pi}{4} + \tan \frac{\pi}{8}}{1 - \tan \frac{\pi}{4} \tan \frac{\pi}{8}}$

$$= \frac{1 + (\sqrt{2} - 1)}{1 - (\sqrt{2} - 1)} = \frac{\sqrt{2}}{2 - \sqrt{2}}$$

$$= \frac{\sqrt{2}(2 + \sqrt{2})}{(2 - \sqrt{2})(2 + \sqrt{2})}$$

$$= \frac{\sqrt{2}}{2} (2 + \sqrt{2}) = \sqrt{2} + 1$$

16 a Let $\sin x - \sqrt{3} \cos x \equiv R \sin(x - \alpha)$

$$\equiv R \sin x \cos \alpha - R \cos x \sin \alpha$$

$$R > 0, 0 < \alpha < 90^\circ$$

Compare $\sin x$: $R \cos \alpha = 1$ (1)

Compare $\cos x$: $R \sin \alpha = \sqrt{3}$ (2)


Divide (2) by (1): $\tan \alpha = \sqrt{3}$

$$\Rightarrow \alpha = 60^\circ$$

$$R^2 = (\sqrt{3})^2 + 1^2 = 4 \Rightarrow R = 2$$

So $\sin x - \sqrt{3} \cos x \equiv 2 \sin(x - 60^\circ)$

16 b Sketch $y = 2 \sin(x - 60^\circ)$ by first translating $y = \sin x$ by 60° to the right and then stretching the result in the y direction by scale factor 2.



Graph meets y -axis when $x = 0$,
i.e. $y = 2 \sin(-60^\circ) = -\sqrt{3}$, at $(0, -\sqrt{3})$

Graph meets x -axis when $y = 0$,
i.e. $(-300^\circ, 0), (-120^\circ, 0), (60^\circ, 0), 240^\circ, 0)$

17 a Let $7 \cos 2\theta + 24 \sin 2\theta \equiv R \cos(2\theta - \alpha)$
 $\equiv R \cos 2\theta \cos \alpha + R \sin 2\theta \sin \alpha$

$$R > 0, 0 < \alpha < \frac{\pi}{2}$$

Compare $\cos 2\theta$: $R \cos \alpha = 7$ (1)

Compare $\sin 2\theta$: $R \sin \alpha = 24$ (2)

Divide (2) by (1): $\tan \alpha = \frac{24}{7}$
 $\Rightarrow \alpha = 1.29$

$$R^2 = 24^2 + 7^2 \Rightarrow R = 25$$

So $7 \cos 2\theta + 24 \sin 2\theta \equiv 25 \cos(2\theta - 1.29)$

b $14 \cos 2\theta + 48 \sin \theta \cos \theta$
 $\equiv 14 \left(\frac{1 + \cos 2\theta}{2} \right) + 24(2 \sin \theta \cos \theta)$
 $\equiv 7(1 + \cos 2\theta) + 24 \sin 2\theta$
 $\equiv 7 + 7 \cos 2\theta + 24 \sin 2\theta$

The maximum value of

$$7 \cos 2\theta + 24 \sin 2\theta \text{ is } 25$$

(using (a) with $\cos(2\theta - 1.29) = 1$)

So maximum value of

$$7 + 7 \cos 2\theta + 24 \sin 2\theta = 7 + 25 = 32$$

Pure Mathematics 3

Solution Bank



17 c Using the answer to part a:

$$\text{Solve } 25\cos(2\theta - 1.29) = 12.5$$

$$\cos(2\theta - 1.29) = \frac{1}{2}$$

$$2\theta - 1.29 = -\frac{\pi}{3}, \frac{\pi}{3}$$

$$\theta = 0.119902..., 1.167099...$$

$$\theta = 0.12, 1.17$$

18 a Let $1.5\sin 2x + 2\cos 2x \equiv R\sin(2x + \alpha)$

$$\equiv R\sin 2x \cos \alpha + R\cos 2x \sin \alpha$$

$$R > 0, 0 < \alpha < \frac{\pi}{2}$$

$$\text{Compare } \sin 2x : R\cos \alpha = 1.5 \quad (1)$$

$$\text{Compare } \cos 2x : R\sin \alpha = 2 \quad (2)$$

$$\text{Divide (2) by (1)} : \tan \alpha = \frac{4}{3}$$

$$\Rightarrow \alpha = 0.927$$

$$R^2 = 2^2 + 1.5^2 \Rightarrow R = 2.5$$

b $3\sin x \cos x + 4\cos^2 x$

$$\equiv \frac{3}{2}(2\sin x \cos x) + 4\left(\frac{1 + \cos 2x}{2}\right)$$

$$\equiv \frac{3}{2}\sin 2x + 2 + 2\cos 2x$$

$$\equiv \frac{3}{2}\sin 2x + 2\cos 2x + 2$$

c From part (a) $1.5\sin 2x + 2\cos 2x \equiv 2.5\sin(2x + 0.927)$

So maximum value of

$$1.5\sin 2x + 2\cos 2x = 2.5 \times 1 = 2.5$$

So maximum value of

$$3\sin x \cos x + 4\cos^2 x = 2.5 + 2 = 4.5$$

$$\mathbf{19 a} \quad \sin^2 \frac{\theta}{2} = 2\sin \theta$$

$$\frac{1 - \cos \theta}{2} = 2\sin \theta$$

$$1 - \cos \theta = 4\sin \theta$$

$$4\sin \theta + \cos \theta = 1$$

$$\text{Let } 4\sin \theta + \cos \theta = R\sin(\theta + \alpha)$$

$$= R\sin \theta \cos \alpha + R\cos \theta \sin \alpha$$

$$\text{So } R\cos \alpha = 4 \text{ and } R\sin \alpha = 1$$

$$\frac{R\sin \alpha}{R\cos \alpha} = \tan \alpha = \frac{1}{4}$$

$$\alpha = \tan^{-1}\left(\frac{1}{4}\right) = \tan^{-1} 0.25 = 14.04^\circ \text{ (2 d.p.)}$$

$$R^2 = 4^2 + 1^2 = \sqrt{17}$$

$$4\sin \theta + \cos \theta = \sqrt{17} \sin(\theta + 14.04^\circ) = 1$$

$$\mathbf{b} \quad \sqrt{17} \sin(\theta + 14.04^\circ) = 1, \text{ for } 0^\circ \leq \theta \leq 360^\circ$$

$$\sin(\theta + 14.04^\circ) = \frac{1}{\sqrt{17}} = 0.24 \text{ (2 d.p.)}$$

$$\theta + 14.04^\circ = \sin^{-1} 0.24 = 14.04^\circ, \text{ for}$$

$$14.04^\circ \leq \theta + 14.04^\circ \leq 374.04^\circ$$

$$\theta + 14.04^\circ = 14.04^\circ, 165.96^\circ, 374.04^\circ$$

$$\theta = 0^\circ, 151.9^\circ, 360^\circ$$

$$\mathbf{20 a} \quad 2\cos \theta = 1 + 3\sin \theta$$

$$\text{So } 2\cos \theta - 3\sin \theta = 1$$

$$\text{Let } 2\cos \theta - 3\sin \theta = R\cos(\theta + \alpha)$$

$$= R\cos \theta \cos \alpha - R\sin \theta \sin \alpha$$

$$\text{So } R\cos \alpha = 2 \text{ and } R\sin \alpha = 3$$

$$\frac{R\sin \alpha}{R\cos \alpha} = \tan \alpha = \frac{3}{2}$$

$$\alpha = \tan^{-1}\left(\frac{3}{2}\right) = 56.3^\circ \text{ (1 d.p.)}$$

$$R^2 = 2^2 + 3^2 = 13$$

$$R = \sqrt{13}$$

$$\text{So } 2\cos \theta - 3\sin \theta = \sqrt{13} \cos(\theta + 56.3^\circ) = 1$$

20 b $\sqrt{13} \cos(\theta + 56.3^\circ) = 1$, for $0^\circ \leq \theta \leq 360^\circ$

$$\cos(\theta + 56.3^\circ) = \frac{1}{\sqrt{13}},$$

$$\text{for } 56.3^\circ \leq \theta + 56.3^\circ \leq 416.3^\circ$$

$$\theta + 56.3^\circ = 73.9^\circ, 286.1^\circ \text{ (1 d.p.)}$$

$$\theta = 17.6^\circ, 229.8^\circ \text{ (1 d.p.)}$$

21 a LHS $\equiv \frac{1}{\cos \theta} \times \frac{1}{\sin \theta} \equiv \frac{1}{\frac{1}{2} \sin 2\theta}$
 $\equiv \frac{2}{\sin 2\theta} \equiv 2 \operatorname{cosec} 2\theta \equiv \text{RHS}$

b LHS $\equiv \frac{\tan \frac{\pi}{4} + \tan x}{1 - \tan \frac{\pi}{4} \tan x} - \frac{\tan \frac{\pi}{4} - \tan x}{1 + \tan \frac{\pi}{4} \tan x}$
 $\equiv \frac{1 + \tan x}{1 - \tan x} - \frac{1 - \tan x}{1 + \tan x}$
 $\equiv \frac{(1 + \tan x)^2 - (1 - \tan x)^2}{(1 - \tan x)(1 + \tan x)}$
 $\equiv \frac{(1 + 2 \tan x + \tan^2 x)}{1 - \tan^2 x} - \frac{(1 - 2 \tan x + \tan^2 x)}{1 - \tan^2 x}$
 $\equiv \frac{4 \tan x}{1 - \tan^2 x}$
 $\equiv 2 \left(\frac{2 \tan x}{1 - \tan^2 x} \right)$
 $\equiv 2 \tan 2x \equiv \text{RHS}$

c LHS $\equiv (\sin x \cos y + \cos x \sin y) \times (\sin x \cos y - \cos x \sin y)$
 $\equiv \sin^2 x \cos^2 y - \cos^2 x \sin^2 y$
 $\equiv (1 - \cos^2 x) \cos^2 y - \cos^2 x (1 - \cos^2 y)$
 $\equiv \cos^2 y - \cos^2 x \cos^2 y - \cos^2 x + \cos^2 x \cos^2 y$
 $\equiv \cos^2 y - \cos^2 x = \text{RHS}$

21 d LHS $\equiv 1 + 2 \cos 2\theta + (2 \cos^2 2\theta - 1)$
 $\equiv 2 \cos 2\theta + 2 \cos^2 2\theta$
 $\equiv 2 \cos 2\theta (1 + \cos 2\theta)$
 $\equiv 2 \cos 2\theta (2 \cos^2 \theta)$
 $\equiv 4 \cos^2 \theta \cos 2\theta = \text{RHS}$

22 a LHS $\equiv \frac{1 - \cos 2x}{1 + \cos 2x} \equiv \frac{1 - (1 - 2 \sin^2 x)}{1 + (2 \cos^2 x - 1)}$
 $\equiv \frac{2 \sin^2 x}{2 \cos^2 x} \equiv \tan^2 x = \text{RHS}$

b $\tan^2 x = 3$

$$\tan x = \pm \sqrt{3}, \text{ for } -\pi \leq x \leq \pi$$

$$\tan x = \sqrt{3} \Rightarrow x = \frac{\pi}{3}, -\frac{2\pi}{3}$$

$$\tan x = -\sqrt{3} \Rightarrow x = -\frac{\pi}{3}, \frac{2\pi}{3}$$

$$x = -\frac{2\pi}{3}, -\frac{\pi}{3}, \frac{\pi}{3}, \frac{2\pi}{3}$$

23 a LHS $\equiv \cos^4 2\theta - \sin^4 2\theta$
 $\equiv (\cos^2 2\theta - \sin^2 2\theta)(\cos^2 2\theta + \sin^2 2\theta)$
 $\equiv (\cos^2 2\theta - \sin^2 2\theta)(1)$
 $\equiv \cos 4\theta \equiv \text{RHS}$

b $\cos 4\theta = \frac{1}{2}$, for $0^\circ \leq 4\theta \leq 720^\circ$
 $4\theta = 60^\circ, 300^\circ, 420^\circ, 660^\circ$
 $\theta = 15^\circ, 75^\circ, 105^\circ, 165^\circ$

24 a LHS $\equiv \frac{1 - (1 - 2 \sin^2 \theta)}{2 \sin \theta \cos \theta}$
 $\equiv \frac{2 \sin^2 \theta}{2 \sin \theta \cos \theta}$
 $\equiv \frac{\sin \theta}{\cos \theta} \equiv \tan \theta = \text{RHS}$

b When $\theta = 180^\circ$, $\sin 2\theta = \sin 360^\circ = 0$ and $2 - 2 \cos 360^\circ = 2 - 2 = 0$ therefore $\theta = 180^\circ$ is a solution of the equation $\sin 2\theta = 2 - 2 \cos 2\theta$

24 c Rearrange $\sin 2\theta = 2 - 2 \cos 2\theta$ to give

$$\frac{2(1-\cos 2\theta)}{\sin 2\theta} = 1$$

Using the identity in part (a) gives
 $2 \tan \theta = 1$

$$\Rightarrow \tan \theta = \frac{1}{2}, \text{ for } 0 < \theta < 360^\circ$$

$$\theta = 26.6^\circ, 206.6^\circ \text{ (1 d.p.)}$$

Challenge

1 a Using $\cos P + \cos Q$

$$\equiv 2 \cos\left(\frac{P+Q}{2}\right) \cos\left(\frac{P-Q}{2}\right)$$

and $\sin P - \sin Q$

$$\equiv 2 \cos\left(\frac{P+Q}{2}\right) \sin\left(\frac{P-Q}{2}\right)$$

$$\text{LHS} \equiv \frac{\cos 2\theta + \cos 4\theta}{\sin 2\theta - \sin 4\theta}$$

$$\equiv \frac{2 \cos\left(\frac{6\theta}{2}\right) \cos\left(\frac{2\theta}{2}\right)}{2 \cos\left(\frac{6\theta}{2}\right) \sin\left(\frac{-2\theta}{2}\right)}$$

$$\equiv \frac{2 \cos 3\theta \cos \theta}{2 \cos 3\theta \sin(-\theta)}$$

$$\equiv \frac{\cos \theta}{\sin(-\theta)} \equiv -\cot \theta$$

b LHS $\equiv \cos x + 2 \cos 3x + \cos 5x$

$$\equiv \cos 5x + \cos x + 2 \cos 3x$$

$$\equiv 2 \cos\left(\frac{6x}{2}\right) \cos\left(\frac{4x}{2}\right) + 2 \cos 3x$$


$$\equiv 2 \cos 3x \cos 2x + 2 \cos 3x$$

$$\equiv 2 \cos 3x (\cos 2x + 1)$$

$$\equiv 2 \cos 3x (2 \cos^2 x)$$

$$\equiv 4 \cos^2 x \cos 3x \equiv \text{RHS}$$

2 a As $\angle OAB = \angle OBA \Rightarrow \angle AOB = \pi - 2\theta$, so
 $\angle BOD = 2\theta$



$$OB = 1$$

$$OD = \cos 2\theta$$

$$BD = \sin 2\theta$$

$$AB = 2 \cos \theta$$

$$\sin \theta = \frac{BD}{AB} = \frac{BD}{2 \cos \theta}$$

$$\text{So } BD = 2 \sin \theta \cos \theta$$

$$\text{But } BD = \sin 2\theta$$

$$\text{So } \sin 2\theta \equiv 2 \sin \theta \cos \theta$$

b $AB = 2 \cos \theta$

$$AD = (2 \cos \theta) \cos \theta = 2 \cos^2 \theta$$

$$OD = 2 \cos^2 \theta - 1$$

$$\text{From part (a)} \quad OD = \cos 2\theta$$

$$\text{So } \cos 2\theta \equiv 2 \cos^2 \theta - 1$$