

Solutionbank

Edexcel AS and A Level Modular Mathematics

Exercise A, Question 1

Question:

Sketch the graphs of

(a) $y = e^x + 1$

(b) $y = 4e^{-2x}$

(c) $y = 2e^x - 3$


(d) $y = 4 - e^x$

(e) $y = 6 + 10e^{\frac{1}{2}x}$

(f) $y = 100e^{-x} + 10$


Solution:

(a) $y = e^x + 1$



This is the normal $y = e^x$ 'moved up' (translated) 1 unit.

(b) $y = 4e^{-2x}$




$$x = 0 \Rightarrow y = 4$$

$$\text{As } x \rightarrow -\infty, y \rightarrow \infty$$

$$\text{As } x \rightarrow \infty, y \rightarrow 0$$

This is an exponential decay type graph.

$$(c) y = 2e^x - 3$$




$$x = 0 \Rightarrow y = 2 \times 1 - 3 = -1$$

$$\text{As } x \rightarrow \infty, y \rightarrow \infty$$

$$\text{As } x \rightarrow -\infty, y \rightarrow 2 \times 0 - 3 = -3$$

$$(d) y = 4 - e^x$$




$$x = 0 \Rightarrow y = 4 - 1 = 3$$

As $x \rightarrow \infty$, $y \rightarrow 4 - \infty$, i.e. $y \rightarrow -\infty$

As $x \rightarrow -\infty$, $y \rightarrow 4 - 0 = 4$

(e) $y = 6 + 10 \frac{1}{2}^x$




$$x = 0 \Rightarrow y = 6 + 10 \times 1 = 16$$

As $x \rightarrow \infty$, $y \rightarrow \infty$

As $x \rightarrow -\infty$, $y \rightarrow 6 + 10 \times 0 = 6$

(f) $y = 100e^{-x} + 10$



$$x = 0 \Rightarrow y = 100 \times 1 + 10 = 110$$

$$\text{As } x \rightarrow \infty, y \rightarrow 100 \times 0 + 10 = 10$$

$$\text{As } x \rightarrow -\infty, y \rightarrow \infty$$

© Pearson Education Ltd 2008

Solutionbank

Edexcel AS and A Level Modular Mathematics

Exercise A, Question 2

Question:

The value of a car varies according to the formula

$$V = 20\,000e^{-\frac{t}{12}}$$

where V is the value in £'s and t is its age in years from new.

- State its value when new.
- Find its value (to the nearest £) after 4 years.
- Sketch the graph of V against t .

Solution:

$$V = 20\,000e^{-\frac{t}{12}}$$

- The new value is when $t = 0$.


$$\Rightarrow V = 20\,000 \times e^{-\frac{0}{12}} = 20\,000 \times 1 = 20\,000$$

New value = £20 000

- Value after 4 years is given when $t = 4$.

$$\Rightarrow V = 20\,000 \times e^{-\frac{4}{12}} = 20\,000 \times e^{-\frac{1}{3}} = 14\,330.63$$

Value after 4 years is £14 331 (to nearest £)



Solutionbank

Edexcel AS and A Level Modular Mathematics

Exercise A, Question 3

Question:

The population of a country is increasing according to the formula

$$P = 20 + 10e^{\frac{t}{50}}$$

where P is the population in thousands and t is the time in years after the year 2000.

- (a) State the population in the year 2000.
- (b) Use the model to predict the population in the year 2020.
- (c) Sketch the graph of P against t for the years 2000 to 2100.

Solution:

$$P = 20 + 10e^{\frac{t}{50}}$$

- (a) The year 2000 corresponds to $t = 0$.

Substitute $t = 0$ into $P = 20 + 10e^{\frac{t}{50}}$

$$P = 20 + 10 \times e^0 = 20 + 10 \times 1 = 30$$


Population = 30 thousand

- (b) The year 2020 corresponds to $t = 20$.

Substitute $t = 20$ into $P = 20 + 10e^{\frac{t}{50}}$

$$P = 20 + 10e^{\frac{20}{50}} = 20 + 14.918 = 34.918 \text{ thousand}$$

Population in 2020 will be 34 918



Year 2100 is $t = 100$

$$P = 20 + 10e^{\frac{100}{50}} = 20 + 10e^2 = 93.891 \text{ thousand}$$

© Pearson Education Ltd 2008

Solutionbank

Edexcel AS and A Level Modular Mathematics

Exercise A, Question 4

Question:

The number of people infected with a disease varies according to the formula
 $N = 300 - 100e^{-0.5t}$
 where N is the number of people infected with the disease and t is the time in years after detection.

- (a) How many people were first diagnosed with the disease?
 (b) What is the long term prediction of how this disease will spread?
 (c) Graph N against t .

Solution:

$$N = 300 - 100e^{-0.5t}$$

(a) The number *first* diagnosed means when $t = 0$.


Substitute $t = 0$ in $N = 300 - 100e^{-0.5t}$

$$N = 300 - 100 \times e^{-0.5 \times 0} = 300 - 100 \times 1 = 200$$

(b) The long term prediction suggests $t \rightarrow \infty$.

As $t \rightarrow \infty$, $e^{-0.5t} \rightarrow 0$

$$\text{So } N \rightarrow 300 - 100 \times 0 = 300$$



Solutionbank

Edexcel AS and A Level Modular Mathematics

Exercise A, Question 5

Question:

The value of an investment varies according to the formula

$$V = A e^{\frac{t}{12}}$$

where V is the value of the investment in £'s, A is a constant to be found and t is the time in years after the investment was made.

- (a) If the investment was worth £8000 after 3 years find A to the nearest £.
- (b) Find the value of the investment after 10 years.
- (c) By what factor will the original investment have increased by after 20 years?

Solution:

$$V = A e^{\frac{t}{12}}$$

- (a) We are given that $V = 8000$ when $t = 3$.

Substituting gives

$$8000 = A e^{\frac{3}{12}}$$

$$8000 = A e^{\frac{1}{4}} \quad (\div e^{\frac{1}{4}})$$

$$A = \frac{8000}{e^{\frac{1}{4}}}$$

$$A = 8000 e^{-\frac{1}{4}}$$

$$A = 6230.41$$

$$A = \text{£}6230 \text{ (to the nearest £)}$$

- (b) Hence $V = (8000 \times e^{-\frac{1}{4}}) e^{\frac{t}{12}}$ (use real value)

After 10 years

$$V = 8000 \times e^{-\frac{1}{4}} \times e^{\frac{10}{12}} \quad (\text{use laws of indices})$$

$$= 8000 \times e^{\frac{10}{12} - \frac{3}{12}}$$

$$= 8000 e^{\frac{7}{12}}$$

$$= £14\,336.01$$

Investment is worth £14 336 (to nearest £) after 10 years.

(c) After 20 years $V = Ae^{\frac{20}{12}}$

This is $e^{\frac{20}{12}}$ times the original amount A
 $= 5.29$ times.

© Pearson Education Ltd 2008

Solutionbank

Edexcel AS and A Level Modular Mathematics

Exercise B, Question 1

Question:

Solve the following equations giving exact solutions:

(a) $e^x = 5$

(b) $\ln x = 4$

(c) $e^{2x} = 7$

(d) $\ln \frac{x}{2} = 4$

(e) $e^{x-1} = 8$

(f) $\ln (2x + 1) = 5$

(g) $e^{-x} = 10$

(h) $\ln (2 - x) = 4$

(i) $2e^{4x} - 3 = 8$

Solution:

(a) $e^x = 5 \Rightarrow x = \ln 5$

(b) $\ln x = 4 \Rightarrow x = e^4$

(c) $e^{2x} = 7 \Rightarrow 2x = \ln 7 \Rightarrow x = \frac{\ln 7}{2}$

(d) $\ln \left(\frac{x}{2} \right) = 4 \Rightarrow \frac{x}{2} = e^4 \Rightarrow x = 2e^4$

(e) $e^{x-1} = 8 \Rightarrow x - 1 = \ln 8 \Rightarrow x = \ln 8 + 1$

(f) $\ln (2x + 1) = 5$
 $\Rightarrow 2x + 1 = e^5$

$$\Rightarrow 2x = e^5 - 1$$

$$\Rightarrow x = \frac{e^5 - 1}{2}$$

$$(g) e^{-x} = 10$$

$$\Rightarrow -x = \ln 10$$

$$\Rightarrow x = -\ln 10$$

$$\Rightarrow x = \ln 10^{-1}$$

$$\Rightarrow x = \ln (0.1)$$

$$(h) \ln (2 - x) = 4$$

$$\Rightarrow 2 - x = e^4$$

$$\Rightarrow 2 = e^4 + x$$

$$\Rightarrow x = 2 - e^4$$

$$(i) 2e^{4x} - 3 = 8$$

$$\Rightarrow 2e^{4x} = 11$$

$$\Rightarrow e^{4x} = \frac{11}{2}$$

$$\Rightarrow 4x = \ln \left(\frac{11}{2} \right)$$

$$\Rightarrow x = \frac{1}{4} \ln \left(\frac{11}{2} \right)$$

Solutionbank

Edexcel AS and A Level Modular Mathematics

Exercise B, Question 2

Question:

Solve the following giving your solution in terms of $\ln 2$:

(a) $e^{3x} = 8$

(b) $e^{-2x} = 4$

(c) $e^{2x+1} = 0.5$

Solution:

(a) $e^{3x} = 8$

$$\Rightarrow 3x = \ln 8$$

$$\Rightarrow 3x = \ln 2^3$$

$$\Rightarrow 3x = 3 \ln 2$$

$$\Rightarrow x = \ln 2$$

(b) $e^{-2x} = 4$

$$\Rightarrow -2x = \ln 4$$

$$\Rightarrow -2x = \ln 2^2$$

$$\Rightarrow -2x = 2 \ln 2$$

$$\Rightarrow x = \frac{2 \ln 2}{-2}$$

$$\Rightarrow x = -1 \ln 2$$

(c) $e^{2x+1} = 0.5$

$$\Rightarrow 2x + 1 = \ln (0.5)$$

$$\Rightarrow 2x + 1 = \ln 2^{-1}$$

$$\Rightarrow 2x + 1 = -\ln 2$$

$$\Rightarrow 2x = -\ln 2 - 1$$

$$\Rightarrow x = \frac{-\ln 2 - 1}{2}$$

Solutionbank

Edexcel AS and A Level Modular Mathematics

Exercise B, Question 3

Question:

Sketch the following graphs stating any asymptotes and intersections with axes:

(a) $y = \ln(x + 1)$

(b) $y = 2 \ln x$

(c) $y = \ln(2x)$


(d) $y = (\ln x)^2$

(e) $y = \ln(4 - x)$

(f) $y = 3 + \ln(x + 2)$

Solution:

(a) $y = \ln(x + 1)$




When $x = 0$, $y = \ln(1) = 0$

When $x \rightarrow -1$, $y \rightarrow -\infty$

y wouldn't exist for values of $x < -1$

When $x \rightarrow \infty$, $y \rightarrow \infty$ (slowly)

(b) $y = 2 \ln x$




When $x = 1$, $y = 2 \ln(1) = 0$

When $x \rightarrow 0$, $y \rightarrow -\infty$

y wouldn't exist for values of $x < 0$

When $x \rightarrow \infty$, $y \rightarrow \infty$ (slowly)

(c) $y = \ln(2x)$




When $x = \frac{1}{2}$, $y = \ln(1) = 0$

When $x \rightarrow 0$, $y \rightarrow -\infty$

y wouldn't exist for values of $x < 0$

When $x \rightarrow \infty$, $y \rightarrow \infty$ (slowly)

(d) $y = (\ln x)^2$




When $x = 1$, $y = (\ln 1)^2 = 0$

For $0 < x < 1$, $\ln x$ is negative, but $(\ln x)^2$ is positive.

When $x \rightarrow 0$, $y \rightarrow \infty$

When $x \rightarrow \infty$, $y \rightarrow \infty$

(e) $y = \ln(4 - x)$



When $x = 3$, $y = \ln 1 = 0$


When $x \rightarrow 4$, $y \rightarrow -\infty$

y doesn't exist for values of $x > 4$

When $x \rightarrow -\infty$, $y \rightarrow \infty$ (slowly)

When $x = 0$, $y = \ln 4$

(f) $y = 3 + \ln(x + 2)$



When $x = -1$, $y = 3 + \ln 1 = 3 + 0 = 3$

When $x \rightarrow -2$, $y \rightarrow -\infty$

y doesn't exist for values of $x < -2$

When $x \rightarrow \infty$, $y \rightarrow \infty$ slowly

When $x = 0$, $y = 3 + \ln(0 + 2) = 3 + \ln 2$

When $y = 0$,

$$0 = 3 + \ln(x + 2)$$

$$-3 = \ln(x + 2)$$

$$e^{-3} = x + 2$$

$$x = e^{-3} - 2$$

Solutionbank

Edexcel AS and A Level Modular Mathematics

Exercise B, Question 4

Question:

The price of a new car varies according to the formula

$$P = 15\,000e^{-\frac{t}{10}}$$

where P is the price in £'s and t is the age in years from new.

- State its new value.
- Calculate its value after 5 years (to the nearest £).
- Find its age when its price falls below £5 000.
- Sketch the graph showing how the price varies over time. Is this a good model?

Solution:

$$P = 15\,000e^{-\frac{t}{10}}$$

$$(a) \text{ New value is when } t = 0 \Rightarrow P = 15\,000 \times e^0 = 15\,000$$

The new value is £15 000

$$(b) \text{ Value after 5 years is when } t = 5$$

$$\Rightarrow P = 15\,000 \times e^{-\frac{5}{10}} = 15\,000e^{-0.5} = 9097.96$$

Value after 5 years is £9 098 (to nearest £)

$$(c) \text{ Find when price is } £5\,000$$

Substitute $P = 5\,000$:

$$5\,000 = 15\,000e^{-\frac{t}{10}} \quad (\div 15\,000)$$

$$\frac{5\,000}{15\,000} = e^{-\frac{t}{10}}$$

$$\frac{1}{3} = e^{-\frac{t}{10}}$$

$$\ln\left(\frac{1}{3}\right) = -\frac{t}{10}$$


$$t = -10\ln\left(\frac{1}{3}\right)$$

$$t = 10 \ln \left(\frac{1}{3} \right) - 1$$

$$t = 10 \ln 3$$

$$t = 10.99 \text{ years}$$

The price falls below £5 000 after 11 years.



A fair model! Perhaps the price should be lower after 11 years.

© Pearson Education Ltd 2008

Solutionbank

Edexcel AS and A Level Modular Mathematics


Exercise B, Question 5

Question:

The graph below is of the function

$$f(x) = \ln(2 + 3x) \quad \{x \in \mathbb{R}, x > a\}.$$

- (a) State the value of a .
- (b) Find the value of s for which $f(s) = 20$.
- (c) Find the function $f^{-1}(x)$ stating its domain.
- (d) Sketch the graphs $f(x)$ and $f^{-1}(x)$ on the same axes stating the relationship between them.



Solution:

(a) $x = a$ is the asymptote to the curve. It will be where

$$2 + 3x = 0$$

$$3x = -2$$

$$x = -\frac{2}{3}$$

Hence $a = -\frac{2}{3}$

(b) If $f(s) = 20$ then

$$\ln(2 + 3s) = 20$$

$$2 + 3s = e^{20}$$

$$3s = e^{20} - 2$$

$$s = \frac{e^{20} - 2}{3}$$

(c) To find $f^{-1}(x)$, change the subject of the formula.

$$y = \ln(2 + 3x)$$


$$e^y = 2 + 3x$$

$$e^y - 2 = 3x$$

$$x = \frac{e^y - 2}{3}$$

$$\text{Therefore } f^{-1}(x) = \frac{e^x - 2}{3}$$

domain of $f^{-1}(x) = \text{range of } f(x)$, so $x \in \mathbb{R}$



$f^{-1}(x)$ is a reflection of $f(x)$ in the line $y = x$.

Solutionbank

Edexcel AS and A Level Modular Mathematics


Exercise B, Question 6

Question:

The graph below is of the function

$$g(x) = 2e^{2x} + 4 \quad \{ x \in \mathbb{R} \} .$$

- (a) Find the range of the function.
- (b) Find the value of p to 2 significant figures.
- (c) Find $g^{-1}(x)$ stating its domain.
- (d) Sketch $g(x)$ and $g^{-1}(x)$ on the same set of axes stating the relationship between them.



Solution:

(a) $g(x) = 2e^{2x} + 4$

As $x \rightarrow -\infty$, $g(x) \rightarrow 2 \times 0 + 4 = 4$

Therefore the range of $g(x)$ is $g(x) > 4$

(b) If $(p, 10)$ lies on $g(x) = 2e^{2x} + 4$

$$2e^{2p} + 4 = 10$$

$$2e^{2p} = 6$$

$$e^{2p} = 3$$

$$2p = \ln 3$$

$$p = \frac{1}{2} \ln 3$$

$$p = 0.55 \text{ (2 s.f.)}$$

(c) $g^{-1}(x)$ is found by changing the subject of the formula.

$$\text{Let } y = 2e^{2x} + 4$$

$$y - 4 = 2e^{2x}$$

$$\frac{y-4}{2} = e^{2x}$$


$$\ln \left(\frac{y-4}{2} \right) = 2x$$

$$x = \frac{1}{2} \ln \left(\frac{y-4}{2} \right)$$

$$\text{Hence } g^{-1}(x) = \frac{1}{2} \ln \left(\frac{x-4}{2} \right)$$

Its domain is the same as the range of $g(x)$.

$g^{-1}(x)$ has a domain of $x > 4$



$g^{-1}(x)$ is a reflection of $g(x)$ in the line $y = x$.

Solutionbank

Edexcel AS and A Level Modular Mathematics

Exercise B, Question 7

Question:

The number of bacteria in a culture grows according to the following equation:

$$N = 100 + 50e^{\frac{t}{30}}$$

where N is the number of bacteria present and t is the time in days from the start of the experiment.

- State the number of bacteria present at the start of the experiment.
- State the number after 10 days.
- State the day on which the number first reaches 1 000 000.
- Sketch the graph showing how N varies with t .

Solution:

$$N = 100 + 50e^{\frac{t}{30}}$$

- (a) At the start $t = 0$

$$\Rightarrow N = 100 + 50e^{\frac{0}{30}} = 100 + 50 \times 1 = 150$$

There are 150 bacteria present at the start.

- (b) After 10 days $t = 10$

$$\Rightarrow N = 100 + 50e^{\frac{10}{30}} = 100 + 50e^{\frac{1}{3}} = 170$$

There are 170 bacteria present after 10 days.

- (c) When $N = 1\,000\,000$

$$1\,000\,000 = 100 + 50e^{\frac{t}{30}} \quad (- 100)$$

$$999\,900 = 50e^{\frac{t}{30}} \quad (\div 50)$$


$$19\,998 = e^{\frac{t}{30}}$$

$$\ln (19\,998) = \frac{t}{30}$$

$$t = 30 \ln (19\,998)$$

$$t = 297.10$$

The number of bacteria reaches 1 000 000 on the 298th day (to the nearest day).



© Pearson Education Ltd 2008

Solutionbank

Edexcel AS and A Level Modular Mathematics


Exercise B, Question 8

Question:

The graph below shows the function

$$h(x) = 40 - 10e^{3x} \quad \{ x > 0, x \in \mathbb{R} \} .$$

- State the range of the function.
- Find the exact coordinates of A in terms of $\ln 2$.
- Find $h^{-1}(x)$ stating its domain.




Solution:

$$(a) h(x) = 40 - 10e^{3x}$$

The range is the set of values that y can take.

$$h(0) = 40 - 10e^0 = 40 - 10 = 30$$

Hence range is $h(x) < 30$



(b) A is where $y = 0$

$$\text{Solve } 40 - 10e^{3x} = 0$$

$$40 = 10e^{3x} \quad (\div 10)$$

$$4 = e^{3x}$$

$$\ln 4 = 3x$$

$$x = \frac{1}{3} \ln 4$$

$$x = \frac{1}{3} \ln 2^2$$

$$x = \frac{2}{3} \ln 2$$

$$\text{A is } \left(\frac{2}{3} \ln 2, 0 \right)$$

(c) To find $h^{-1}(x)$ change the subject of the formula.

$$\text{Let } y = 40 - 10e^{3x}$$

$$10e^{3x} = 40 - y$$

$$e^{3x} = \frac{40 - y}{10}$$

$$3x = \ln \left(\frac{40 - y}{10} \right)$$

$$x = \frac{1}{3} \ln \left(\frac{40 - y}{10} \right)$$

The domain of the inverse function is the same as the range of the function.

$$\text{Hence } h^{-1}(x) = \frac{1}{3} \ln \left(\frac{40 - x}{10} \right) \quad \{ x \in \mathbb{R}, x < 30 \}$$

© Pearson Education Ltd 2008

Solutionbank

Edexcel AS and A Level Modular Mathematics

Exercise C, Question 1

Question:

Sketch the following functions stating any asymptotes and intersections with axes:

(a) $y = e^x + 3$

(b) $y = \ln(-x)$

(c) $y = \ln(x + 2)$


(d) $y = 3e^{-2x} + 4$

(e) $y = e^{x+2}$

(f) $y = 4 - \ln x$

Solution:


(a) $y = e^x + 3$



This is the graph of $y = e^x$ 'moved up' 3 units.

$$x = 0, \quad y = e^0 + 3 = 1 + 3 = 4$$

(b) $y = \ln(-x)$




$$x = -1, \quad y = \ln(-(-1)) = \ln(1) = 0$$

y will not exist for values of $x > 0$

$$x \rightarrow -\infty, \quad y \rightarrow \infty \text{ (slowly)}$$

The graph will be a reflection of $y = \ln(x)$ in the y axis.

(c) $y = \ln(x + 2)$



$$x = -1, \quad y = \ln(-1 + 2) = \ln(1) = 0$$


y will not exist for values of $x < -2$

$$x \rightarrow -2, \quad y \rightarrow -\infty$$

$$x \rightarrow \infty, \quad y \rightarrow \infty \text{ (slowly)}$$

$$x = 0, \quad y = \ln(0 + 2) = \ln 2$$

(d) $y = 3e^{-2x} + 4$




$$x = 0, \quad y = 3e^0 + 4 = 3 + 4 = 7$$

$$x \rightarrow \infty, \quad y \rightarrow 3 \times 0 + 4 = 4$$

$$x \rightarrow -\infty, \quad y \rightarrow \infty$$

(e) $y = e^{x+2}$




$$x = -2, \quad y = e^{-2+2} = e^0 = 1$$

$$x \rightarrow -\infty, \quad y \rightarrow 0$$

$$x \rightarrow \infty, \quad y \rightarrow \infty$$

$$x = 0, \quad y = e^2$$

(f) $y = 4 - \ln x$



$$x = 1, \quad y = 4 - \ln(1) = 4$$

$$x \rightarrow 0, \quad y \rightarrow 4 - (-\infty), \text{ so } y \rightarrow +\infty$$

y will not exist for values of $x < 0$

$$y = 0 \Rightarrow 4 - \ln x = 0 \Rightarrow \ln x = 4 \Rightarrow x = e^4$$

Solutionbank

Edexcel AS and A Level Modular Mathematics

Exercise C, Question 2

Question:

Solve the following equations, giving exact solutions:

(a) $\ln (2x - 5) = 8$

(b) $e^{4x} = 5$

(c) $24 - e^{-2x} = 10$

(d) $\ln x + \ln (x - 3) = 0$

(e) $e^x + e^{-x} = 2$

(f) $\ln 2 + \ln x = 4$

Solution:

(a) $\ln (2x - 5) = 8$ (inverse of \ln)

$$2x - 5 = e^8 \quad (+ 5)$$

$$2x = e^8 + 5 \quad (\div 2)$$

$$x = \frac{e^8 + 5}{2}$$

(b) $e^{4x} = 5$ (inverse of e)

$$4x = \ln 5 \quad (\div 4)$$

$$x = \frac{\ln 5}{4}$$

(c) $24 - e^{-2x} = 10$ ($+ e^{-2x}$)

$$24 = 10 + e^{-2x} \quad (- 10)$$

$$14 = e^{-2x} \quad (\text{inverse of } e)$$

$$\ln (14) = - 2x \quad (\div - 2)$$

$$- \frac{1}{2} \ln (14) = x$$

$$x = - \frac{1}{2} \ln (14)$$

$$(d) \ln(x) + \ln(x - 3) = 0$$

$$\ln[x(x - 3)] = 0$$

$$x(x - 3) = e^0$$

$$x(x - 3) = 1$$

$$x^2 - 3x - 1 = 0$$

$$x = \frac{3 \pm \sqrt{9 + 4}}{2}$$

$$x = \frac{3 \pm \sqrt{13}}{2}$$

$$x = \frac{3 + \sqrt{13}}{2}$$

(x cannot be negative because of initial equation)

$$(e) e^x + e^{-x} = 2$$

$$e^x + \frac{1}{e^x} = 2 \quad (\times e^x)$$

$$(e^x)^2 + 1 = 2e^x$$

$$(e^x)^2 - 2e^x + 1 = 0$$

$$(e^x - 1)^2 = 0$$

$$e^x = 1$$

$$x = \ln 1 = 0$$

$$(f) \ln 2 + \ln x = 4$$

$$\ln 2x = 4$$

$$2x = e^{\text{hairsp};4}$$

$$x = \frac{e^{\text{hairsp};4}}{2}$$


Solutionbank

Edexcel AS and A Level Modular Mathematics

Exercise C, Question 3

Question:

The function $c(x) = 3 + \ln(4 - x)$ is shown below.



- State the exact coordinates of point A.
- Calculate the exact coordinates of point B.
- Find the inverse function $c^{-1}(x)$ stating its domain.
- Sketch $c(x)$ and $c^{-1}(x)$ on the same set of axes stating the relationship between them.

Solution:

(a) A is where $x = 0$

Substitute $x = 0$ into $y = 3 + \ln(4 - x)$ to give

$$y = 3 + \ln 4$$

$$A = (0, 3 + \ln 4)$$

(b) B is where $y = 0$

Substitute $y = 0$ into $y = 3 + \ln(4 - x)$ to give

$$0 = 3 + \ln(4 - x)$$

$$-3 = \ln(4 - x)$$

$$e^{-3} = 4 - x$$

$$x = 4 - e^{-3}$$

$$B = (4 - e^{-3}, 0)$$

(c) To find $c^{-1}(x)$ change the subject of the formula.

$$y = 3 + \ln(4 - x)$$


$$y - 3 = \ln(4 - x)$$

$$e^{y-3} = 4 - x$$

$$x = 4 - e^{y-3}$$

The domain of the inverse function is the range of the function. Looking at graph this is all the real numbers. So

$$c^{-1}(x) = 4 - e^{x-3} \quad \{ x \in \mathbb{R} \}$$



© Pearson Education Ltd 2008

Solutionbank

Edexcel AS and A Level Modular Mathematics

Exercise C, Question 4

Question:

The price of a computer system can be modelled by the formula

$$P = 100 + 850e^{-\frac{t}{2}}$$

where P is the price of the system in £s and t is the age of the computer in years after being purchased.

- Calculate the new price of the system.
- Calculate its price after 3 years.
- When will it be worth less than £200?
- Find its price as $t \rightarrow \infty$.
- Sketch the graph showing P against t .
Comment on the appropriateness of this model.

Solution:

$$P = 100 + 850e^{-\frac{t}{2}}$$

- New price is when $t = 0$.

Substitute $t = 0$ into $P = 100 + 850e^{-\frac{t}{2}}$ to give

$$P = 100 + 850e^{-\frac{0}{2}} \quad (e^0 = 1)$$

$$= 100 + 850 = 950$$

The new price is £950

- After 3 years $t = 3$.

Substitute $t = 3$ into $P = 100 + 850e^{-\frac{t}{2}}$ to give

$$P = 100 + 850e^{-\frac{3}{2}} = 289.66$$

Price after 3 years is £290 (to nearest £)

- It is worth less than £200 when $P < 200$

Substitute $P = 200$ into $P = 100 + 850e^{-\frac{t}{2}}$ to give

$$200 = 100 + 850e^{-\frac{t}{2}}$$

$$100 = 850e^{-\frac{t}{2}}$$

$$\frac{100}{850} = e^{-\frac{t}{2}}$$

$$\ln\left(\frac{100}{850}\right) = -\frac{t}{2}$$

$$t = -2\ln\left(\frac{100}{850}\right)$$


$$t = 4.28$$

It is worth less than £200 after 4.28 years.

(d) As $t \rightarrow \infty$, $e^{-\frac{t}{2}} \rightarrow 0$

Hence $P \rightarrow 100 + 850 \times 0 = 100$

The computer will be worth £100 eventually.



© Pearson Education Ltd 2008

Solutionbank

Edexcel AS and A Level Modular Mathematics

Exercise C, Question 5

Question:

The function f is defined by

$$f : x \rightarrow \ln (5x - 2) \quad \left\{ x \in \mathbb{R}, x > \frac{2}{5} \right\} .$$

- (a) Find an expression for $f^{-1} (x)$.
- (b) Write down the domain of $f^{-1} (x)$.
- (c) Solve, giving your answer to 3 decimal places,
 $\ln (5x - 2) = 2$.

[E]

Solution:

(a) Let $y = \ln (5x - 2)$

$$e^y = 5x - 2$$

$$e^y + 2 = 5x$$

$$\frac{e^y + 2}{5} = x$$

The range of $y = \ln (5x - 2)$ is $y \in \mathbb{R}$

$$\text{So } f^{-1} (x) = \frac{e^x + 2}{5} \quad \{ x \in \mathbb{R} \}$$

(b) Domain is $x \in \mathbb{R}$

(c) $\ln (5x - 2) = 2$

$$5x - 2 = e^2$$

$$5x = e^2 + 2$$

$$x = \frac{e^2 + 2}{5} = 1.8778 \quad \dots$$

$$x = 1.878 \text{ (to 3d.p.)}$$

Solutionbank

Edexcel AS and A Level Modular Mathematics

Exercise C, Question 6

Question:

The functions f and g are given by

$$f : x \rightarrow 3x - 1 \quad \{ x \in \mathbb{R} \}$$

$$g : x \rightarrow e^{\frac{x}{2}} \quad \{ x \in \mathbb{R} \}$$

- (a) Find the value of $fg(4)$, giving your answer to 2 decimal places.
- (b) Express the inverse function $f^{-1}(x)$ in the form $f^{-1} : x \rightarrow \dots$.
- (c) Using the same axes, sketch the graphs of the functions f and gf . Write on your sketch the value of each function at $x = 0$.
- (d) Find the values of x for which $f^{-1}(x) = \frac{5}{f(x)}$.

[E]


Solution:

$$\begin{aligned} \text{(a) } fg(4) &= f\left(e^{\frac{4}{2}}\right) = f(e^2) = 3e^2 - 1 \\ &= 21.17 \text{ (2d.p.)} \end{aligned}$$

$$\text{(b) If } f : x \rightarrow 3x - 1 \quad \{ x \in \mathbb{R} \}$$


$$\text{then } f^{-1} : x \rightarrow \frac{x+1}{3} \quad \left\{ x \in \mathbb{R} \right\}$$

by using flow diagram method:



$$\text{(c) } gf(x) = g(3x - 1) = e^{\frac{3x-1}{2}} \quad f(x) = 3x - 1$$

At $x = 0$, $gf(x) = e^{\frac{0-1}{2}} = e^{-\frac{1}{2}}$ and $f(x) = 3 \times 0 - 1 = -1$



$$(d) f^{-1}(x) = \frac{5}{f(x)}$$

$$\frac{x+1}{3} = \frac{5}{3x-1} \quad (\text{cross multiply})$$

$$(x+1)(3x-1) = 5 \times 3$$

$$3x^2 + 2x - 1 = 15$$

$$3x^2 + 2x - 16 = 0$$

$$(3x+8)(x-2) = 0$$

$$x = 2, -\frac{8}{3}$$

Solutionbank

Edexcel AS and A Level Modular Mathematics

Exercise C, Question 7

Question:


The points P and Q lie on the curve with equation $y = e^{\frac{1}{2}x}$.
The x -coordinates of P and Q are $\ln 4$ and $\ln 16$ respectively.

- Find an equation for the line PQ.
- Show that this line passes through the origin O.
- Calculate the length, to 3 significant figures, of the line segment PQ.

[E]

Solution:

(a)



Q has y coordinate $e^{\frac{1}{2}\ln 16} = e^{\ln 16 \frac{1}{2}} = 16^{\frac{1}{2}} = 4$

P has y coordinate $e^{\frac{1}{2}\ln 4} = e^{\ln 4 \frac{1}{2}} = 4^{\frac{1}{2}} = 2$

$$\text{Gradient of the line PQ} = \frac{\text{change in } y}{\text{change in } x} = \frac{4 - 2}{\ln 16 - \ln 4} = \frac{2}{\ln \frac{16}{4}} = \frac{2}{\ln 4}$$

Using $y = mx + c$ the equation of the line PQ is

$$y = \frac{2}{\ln 4}x + c$$

$(\ln 4, 2)$ lies on line so

$$2 = \frac{2}{\ln 4} \times \ln 4 + c$$

$$2 = 2 + c$$

$$c = 0$$

$$\text{Equation of PQ is } y = \frac{2x}{\ln 4}$$

(b) The line passes through the origin as $c = 0$.

(c) Length from $(\ln 4, 2)$ to $(\ln 16, 4)$ is

$$\sqrt{(\ln 16 - \ln 4)^2 + 4 - 2)^2} = \sqrt{\left(\ln \frac{16}{4}\right)^2 + 2^2} = \sqrt{(\ln 4) + 4} = 2.43$$

Solutionbank

Edexcel AS and A Level Modular Mathematics

Exercise C, Question 8

Question:

The functions f and g are defined over the set of real numbers by

$$f : x \rightarrow 3x - 5$$

$$g : x \rightarrow e^{-2x}$$

(a) State the range of $g(x)$.


(b) Sketch the graphs of the inverse functions f^{-1} and g^{-1} and write on your sketches the coordinates of any points at which a graph meets the coordinate axes.

(c) State, giving a reason, the number of roots of the equation

$$f^{-1}(x) = g^{-1}(x).$$

(d) Evaluate $fg\left(-\frac{1}{3}\right)$, giving your answer to 2 decimal places.


Solution:



$$g(x) > 0$$

$$(b) f^{-1}(x) = \frac{x+5}{3}$$

$$g^{-1}(x) = -\frac{1}{2} \ln x$$



(c) $f^{-1}(x) = g^{-1}(x)$ would have 1 root because there is 1 point of intersection.

$$(d) fg\left(-\frac{1}{3}\right) = f\left(e^{-2 \times -\frac{1}{3}}\right) = f\left(e^{\frac{2}{3}}\right) = 3 \times e^{\frac{2}{3}} - 5 = 0.84$$

Solutionbank

Edexcel AS and A Level Modular Mathematics

Exercise C, Question 9

Question:

The function f is defined by $f : x \rightarrow e^x + k$, $x \in \mathbb{R}$ and k is a positive constant.


- (a) State the range of $f(x)$.
- (b) Find $f(\ln k)$, simplifying your answer.
- (c) Find f^{-1} , the inverse function of f , in the form $f^{-1} : x \rightarrow \dots$, stating its domain.
- (d) On the same axes, sketch the curves with equations $y = f(x)$ and $y = f^{-1}(x)$, giving the coordinates of all points where the graphs cut the axes.

[E]

Solution:

(a) $f : x \rightarrow e^x + k$

As $x \rightarrow -\infty$, $f(x) \rightarrow 0 + k = k$



Range of $f(x)$ is $f(x) > k$

(b) $f(\ln k) = e^{\ln k} + k = k + k = 2k$


(c) Let $y = e^x + k$

$$y - k = e^x$$

$$\ln (y - k) = x$$

Hence $f^{-1} : x \rightarrow \ln (x - k) , x > k$

(d)



© Pearson Education Ltd 2008

Solutionbank

Edexcel AS and A Level Modular Mathematics

Exercise C, Question 10

Question:

The function f is given by

$$f : x \rightarrow \ln(4 - 2x) \quad \{ x \in \mathbb{R}, x < 2 \}$$

(a) Find an expression for $f^{-1}(x)$.

(b) Sketch the curve with equation $y = f^{-1}(x)$, showing the coordinates of the points where the curve meets the axes.

(c) State the range of $f^{-1}(x)$.

The function g is given by

$$g : x \rightarrow e^x \quad \{ x \in \mathbb{R} \}$$

(d) Find the value of $gf(0.5)$.

[E]

Solution:

$$f(x) = \ln(4 - 2x) \quad \{ x \in \mathbb{R}, x < 2 \}$$


(a) Let $y = \ln(4 - 2x)$ and change the subject of the formula.

$$e^y = 4 - 2x$$

$$2x = 4 - e^y$$

$$x = \frac{4 - e^y}{2}$$

$$f^{-1} : x \rightarrow \frac{4 - e^x}{2} \quad \{ x \in \mathbb{R} \}$$



$$x = 0 \Rightarrow f^{-1}(x) = \frac{4-1}{2} = \frac{3}{2}$$

$$y = 0 \Rightarrow \frac{4-e^x}{2} = 0 \Rightarrow e^x = 4 \Rightarrow x = \ln 4$$

$$x \rightarrow \infty, y \rightarrow -\infty$$

$$x \rightarrow -\infty, y \rightarrow \frac{4-0}{2} = 2$$

(c) Range of $f^{-1}(x)$ is $f^{-1}(x) < 2$

$$(d) gf(0.5) = g[\ln(4 - 2 \times 0.5)] = g(\ln 3) = e^{\ln 3} = 3$$

Solutionbank

Edexcel AS and A Level Modular Mathematics

Exercise C, Question 11

Question:

The function $f(x)$ is defined by

$$f(x) = 3x^3 - 4x^2 - 5x + 2$$

- (a) Show that $(x + 1)$ is a factor of $f(x)$.
- (b) Factorise $f(x)$ completely.
- (c) Solve, giving your answers to 2 decimal places, the equation
 $3[\ln(2x)]^3 - 4[\ln(2x)]^2 - 5\ln(2x) + 2 = 0 \quad x > 0$

[E]

Solution:

$$f(x) = 3x^3 - 4x^2 - 5x + 2$$

$$\begin{aligned} \text{(a) } f(-1) &= 3 \times (-1)^3 - 4 \times (-1)^2 - 5 \times (-1) \\ &+ 2 = -3 - 4 + 5 + 2 = 0 \end{aligned}$$

As $f(-1) = 0$ then $(x + 1)$ is a factor.

$$\text{(b) } f(x) = 3x^3 - 4x^2 - 5x + 2$$

$$f(x) = (x + 1)(3x^2 - 7x + 2) \quad (\text{by inspection})$$

$$f(x) = (x + 1)(3x - 1)(x - 2)$$

$$\begin{aligned} \text{(c) If } 3[\ln(2x)]^3 - 4[\ln(2x)]^2 - 5[\ln(2x)] + 2 &= 0 \\ \Rightarrow [\ln(2x) + 1][3\ln(2x) - 1][\ln(2x) - 2] &= 0 \end{aligned}$$

$$\Rightarrow \ln(2x) = -1, \frac{1}{3}, 2$$

$$\Rightarrow 2x = e^{-1}, e^{\frac{1}{3}}, e^2$$

$$\Rightarrow x = \frac{1}{2}e^{-1}, \frac{1}{2}e^{\frac{1}{3}}, \frac{1}{2}e^2$$

$$\Rightarrow x = 0.18, 0.70, 3.69$$