

C3**DIFFERENTIATION****Answers - Worksheet D**

1 $x = \frac{1}{2} \therefore y = \frac{1}{4}$

$$\frac{dy}{dx} = 2x + \frac{1}{4x-1} \times 4 = 2x + \frac{4}{4x-1}$$

$$\text{grad} = 1 + 4 = 5$$

$$\therefore y - \frac{1}{4} = 5(x - \frac{1}{2})$$

$$[y = 5x - \frac{9}{4}]$$

2 **a** $\sqrt{8-e^{2x}} = 2$

$$8 - e^{2x} = 4$$

$$x = \frac{1}{2} \ln 4 = \ln 2$$

b $\frac{dy}{dx} = \frac{1}{2}(8-e^{2x})^{-\frac{1}{2}} \times (-2e^{2x})$

$$= \frac{-e^{2x}}{\sqrt{8-e^{2x}}}$$

$$\text{grad} = -2$$

$$\therefore y - 2 = -2(x - \ln 2)$$

$$2x + y = 2 + 2 \ln 2$$

$$2x + y = 2 + \ln 2^2$$

$$2x + y = 2 + \ln 4$$

3 **a** $\frac{dy}{dx} = 2 + \frac{1}{4-2x} \times (-2) = 2 - \frac{1}{2-x}$

$$\frac{d^2y}{dx^2} = (2-x)^{-2} \times (-1) = \frac{-1}{(2-x)^2}$$

b SP: $2 - \frac{1}{2-x} = 0$

$$2-x = \frac{1}{2}$$

$$x = \frac{3}{2} \therefore (\frac{3}{2}, 4)$$

c $x = \frac{3}{2}, \frac{d^2y}{dx^2} = -4 \therefore \text{maximum}$

4 **a** $\frac{dy}{dx} = -3(2x+1)^{-2} \times 2 = \frac{-6}{(2x+1)^2}$

$$x = 1, \text{ grad} = -\frac{2}{3}, \therefore \text{grad of normal} = \frac{3}{2}$$

$$\therefore y - 1 = \frac{3}{2}(x - 1)$$

$$[y = \frac{3}{2}x - \frac{1}{2}]$$

b at $Q \quad \frac{3x-1}{2} = \frac{3}{2x+1}$

$$(3x-1)(2x+1) = 6$$

$$6x^2 + x - 7 = 0$$

$$(6x+7)(x-1) = 0$$

$$x = 1 \text{ (at } P \text{) or } -\frac{7}{6}$$

$$\therefore Q(-\frac{7}{6}, -\frac{9}{4})$$

5 **a** $t = 0, N = 20 \therefore a = 20$

$$t = 8, N = 60 \therefore 60 = 20e^{8k}$$

$$k = \frac{1}{8} \ln 3 = 0.137 \text{ (3sf)}$$

b $N = 20e^{0.1373t}$

$$t = 12, N = 104 \text{ (3sf)}$$

c $\frac{dN}{dt} = 20 \times 0.1373e^{0.1373t} = 2.747e^{0.1373t}$

$$t = 12, \frac{dN}{dt} = 14.3$$

$\therefore N$ increasing at 14.3 per second (3sf)

6 **a** $= 3(5-2x^2)^2 \times (-4x)$

$$= -12x(5-2x^2)^2$$

b SP: $-12x(5-2x^2)^2 = 0$

$$x = 0 \text{ or } x^2 = \frac{5}{2}$$

$$x = 0, \pm \frac{1}{2}\sqrt{10}$$

$$\therefore (-\frac{1}{2}\sqrt{10}, 0), (0, 125), (\frac{1}{2}\sqrt{10}, 0)$$

c $x = \frac{3}{2}, y = \frac{1}{8}$

$$\text{grad} = -18 \times \frac{1}{4} = -\frac{9}{2}$$

$$\therefore y - \frac{1}{8} = -\frac{9}{2}(x - \frac{3}{2})$$

$$8y - 1 = -36x + 54$$

$$36x + 8y - 55 = 0$$

C3 DIFFERENTIATION*Answers - Worksheet D page 2*

7 **a** $\frac{dy}{dx} = 4 - e^{2x}$
 SP: $4 - e^{2x} = 0$
 $x = \frac{1}{2} \ln 4 = \ln 2$
 $\therefore (\ln 2, 4 \ln 2 - 2)$

b $\frac{d^2y}{dx^2} = -2e^{2x}$
 $x = \ln 2; \frac{d^2y}{dx^2} = -8 \therefore \text{maximum}$

8 **a** $f'(x) = \frac{3}{x} - 2$
b grad of curve = 4
 $\therefore \frac{3}{x} - 2 = 4$
 $x = \frac{1}{2}$
c SP: $\frac{3}{x} - 2 = 0$
 $x = \frac{3}{2} \quad \therefore (\frac{3}{2}, 3 \ln \frac{15}{2} - 3)$
d $x \geq \frac{3}{2}$

9 **a** $\frac{dy}{dx} = \frac{1}{2}(x^2 + 3)^{-\frac{1}{2}} \times 2x = \frac{x}{\sqrt{x^2 + 3}}$
 at A, grad = $-\frac{1}{2}$
 $\therefore y - 2 = -\frac{1}{2}(x + 1)$
 $[y = \frac{3}{2} - \frac{1}{2}x]$

b at B, grad = $\frac{1}{2}$
 grad of normal = -2
 $\therefore y - 2 = -2(x - 1)$
 $[y = 4 - 2x]$

c $\frac{3}{2} - \frac{1}{2}x = 4 - 2x$
 $x = \frac{5}{3}$

11 **a** $f'(x) = 2x - 7 + \frac{4}{x} = 0$
 $2x^2 - 7x + 4 = 0$
 $x = \frac{7 \pm \sqrt{49 - 32}}{4} = \frac{7 \pm \sqrt{17}}{4}$
 $x = 0.72, 2.78$

b $x = 2 \therefore y = -10, \text{grad} = -1$
 $\therefore y + 10 = -(x - 2)$
 $[y = -x - 8]$

10 **a** 80°C
b 20°C , as $t \rightarrow \infty, T \rightarrow 20$
c $30 = 20 + 60e^{-25k}$
 $e^{-25k} = \frac{30-20}{60} = \frac{1}{6}$
 $k = \frac{-1}{25} \ln \frac{1}{6} = 0.0717 \text{ (3sf)}$
d $T = 20 + 60e^{-0.07167t}$
 $\frac{dT}{dt} = 60 \times (-0.07167)e^{-0.07167t}$
 $= -4.300e^{-0.07167t}$
 $t = 40, \frac{dT}{dt} = -0.245$
 $\therefore \text{temp. decreasing at } 0.245^\circ\text{C min}^{-1} \text{ (3sf)}$

12 **a** $\frac{dy}{dx} = 2x + 8(x - 1)^{-2}$
 SP: $2x + \frac{8}{(x-1)^2} = 0$
 $2x(x - 1)^2 + 8 = 0$
 $2x(x^2 - 2x + 1) + 8 = 0$
 $2x^3 - 4x^2 + 2x + 8 = 0$
 $x^3 - 2x^2 + x + 4 = 0$
b let $f(x) = x^3 - 2x^2 + x + 4$
 $f(1) = 4, f(2) = 6, f(-1) = 0$
 $\therefore (x + 1) \text{ is a factor}$
 $\therefore (x + 1)(x^2 - 3x + 4) = 0$
 $x = -1 \text{ or } x^2 - 3x + 4 = 0$
 $b^2 - 4ac = 9 - 16 = -7$
 $b^2 - 4ac < 0 \therefore \text{no real roots}$
 $\therefore \text{exactly one SP}$
 $(-1, 5)$

c $\frac{d^2y}{dx^2} = 2 - 16(x - 1)^{-3}$
 when $x = -1, \frac{d^2y}{dx^2} = 4$
 $\frac{d^2y}{dx^2} > 0 \therefore \text{minimum}$