C3 DIFFERENTIATION

Worksheet D

- Find an equation for the tangent to the curve with equation $y = x^2 + \ln(4x 1)$ at the point on the curve where $x = \frac{1}{2}$.
- 2 A curve has the equation $y = \sqrt{8 e^{2x}}$.


The point *P* on the curve has *y*-coordinate 2.

- **a** Find the x-coordinate of P.
- **b** Show that the tangent to the curve at *P* has equation

$$2x + y = 2 + \ln 4$$
.

- 3 A curve has the equation $y = 2x + 1 + \ln(4 2x)$, x < 2.
 - **a** Find and simplify expressions for $\frac{dy}{dx}$ and $\frac{d^2y}{dx^2}$.
 - **b** Find the coordinates of the stationary point of the curve.
 - c Determine the nature of this stationary point.

4



The diagram shows the curve with equation $y = \frac{3}{2x+1}$.

a Find an equation for the normal to the curve at the point P(1, 1).

The normal to the curve at P intersects the curve again at the point Q.

- **b** Find the exact coordinates of Q.
- 5 A quantity *N* is increasing such that at time *t* seconds,

$$N = ae^{kt}$$
.

Given that at time t = 0, N = 20 and that at time t = 8, N = 60, find

- **a** the values of the constants a and k,
- **b** the value of *N* when t = 12,
- **c** the rate at which *N* is increasing when t = 12.

6


$$f(x) \equiv (5 - 2x^2)^3$$
.

- a Find f'(x).
- **b** Find the coordinates of the stationary points of the curve y = f(x).
- **c** Find the equation for the tangent to the curve y = f(x) at the point with x-coordinate $\frac{3}{2}$, giving your answer in the form ax + by + c = 0, where a, b and c are integers.

C3 DIFFERENTIATION

- 7 A curve has the equation $y = 4x \frac{1}{2}e^{2x}$.
 - **a** Find the coordinates of the stationary point of the curve, giving your answers in terms of natural logarithms.
 - **b** Determine the nature of the stationary point.

8



The diagram shows the curve y = f(x) where $f(x) = 3 \ln 5x - 2x$, x > 0.

- a Find f'(x).
- **b** Find the x-coordinate of the point on the curve at which the gradient of the normal to the curve is $-\frac{1}{4}$.
- c Find the coordinates of the maximum turning point of the curve.
- **d** Write down the set of values of x for which f(x) is a decreasing function.
- 9 The curve C has the equation $y = \sqrt{x^2 + 3}$.
 - **a** Find an equation for the tangent to C at the point A(-1, 2).
 - **b** Find an equation for the normal to C at the point B(1, 2).
 - **c** Find the x-coordinate of the point where the tangent to C at A meets the normal to C at B.
- A bucket of hot water is placed outside and allowed to cool. The surface temperature of the water, T °C, after t minutes is given by

$$T = 20 + 60e^{-kt}$$
.

where k is a positive constant.

- a State the initial surface temperature of the water.
- **b** State, with a reason, the air temperature around the bucket.

Given that T = 30 when t = 25,

- **c** find the value of k,
- **d** find the rate at which the surface temperature of the water is decreasing when t = 40.

11
$$f(x) \equiv x^2 - 7x + 4 \ln\left(\frac{x}{2}\right), \ x > 0.$$

- a Solve the equation f'(x) = 0, giving your answers correct to 2 decimal places.
- **b** Find an equation for the tangent to the curve y = f(x) at the point on the curve where x = 2.
- 12 A curve has the equation $y = x^2 \frac{8}{x-1}$.
 - a Show that the x-coordinate of any stationary point of the curve satisfies the equation $x^3 2x^2 + x + 4 = 0$.
 - **b** Hence, show that the curve has exactly one stationary point and find its coordinates.
 - c Determine the nature of this stationary point.