



GCE

Mathematics

Advanced GCE

Unit 4724: Core Mathematics 4

Mark Scheme for June 2012

OCR (Oxford Cambridge and RSA) is a leading UK awarding body, providing a wide range of qualifications to meet the needs of candidates of all ages and abilities. OCR qualifications include AS/A Levels, Diplomas, GCSEs, OCR Nationals, Functional Skills, Key Skills, Entry Level qualifications, NVQs and vocational qualifications in areas such as IT, business, languages, teaching/training, administration and secretarial skills.

It is also responsible for developing new specifications to meet national requirements and the needs of students and teachers. OCR is a not-for-profit organisation; any surplus made is invested back into the establishment to help towards the development of qualifications and support, which keep pace with the changing needs of today's society.

This mark scheme is published as an aid to teachers and students, to indicate the requirements of the examination. It shows the basis on which marks were awarded by examiners. It does not indicate the details of the discussions which took place at an examiners' meeting before marking commenced.

All examiners are instructed that alternative correct answers and unexpected approaches in candidates' scripts must be given marks that fairly reflect the relevant knowledge and skills demonstrated.

Mark schemes should be read in conjunction with the published question papers and the report on the examination.

OCR will not enter into any discussion or correspondence in connection with this mark scheme.

© OCR 2012

Any enquiries about publications should be addressed to:

OCR Publications
PO Box 5050
Annesley
NOTTINGHAM
NG15 0DL

Telephone: 0870 770 6622
Facsimile: 01223 552610
E-mail: publications@ocr.org.uk

Annotations

Annotation in scoris	Meaning
✓and ✖	
BOD	Benefit of doubt
FT	Follow through
ISW	Ignore subsequent working
M0, M1	Method mark awarded 0, 1
A0, A1	Accuracy mark awarded 0, 1
B0, B1	Independent mark awarded 0, 1
SC	Special case
^	Omission sign
MR	Misread
Highlighting	
Other abbreviations in mark scheme	Meaning
E1	Mark for explaining
U1	Mark for correct units
G1	Mark for a correct feature on a graph
M1 dep*	Method mark dependent on a previous mark, indicated by *
cao	Correct answer only
oe	Or equivalent
rot	Rounded or truncated
soi	Seen or implied
www	Without wrong working
B2	Independent mark awarded 2
A1 dep*	Accuracy mark dependent on a previous mark, indicated by *

Subject-specific Marking Instructions for GCE Mathematics Pure strand

- a Annotations should be used whenever appropriate during your marking.

The A, M and B annotations must be used on your standardisation scripts for responses that are not awarded either 0 or full marks. It is vital that you annotate standardisation scripts fully to show how the marks have been awarded.

For subsequent marking you must make it clear how you have arrived at the mark you have awarded.

- b An element of professional judgement is required in the marking of any written paper. Remember that the mark scheme is designed to assist in marking incorrect solutions. Correct *solutions* leading to correct answers are awarded full marks but work must not be judged on the answer alone, and answers that are given in the question, especially, must be validly obtained; key steps in the working must always be looked at and anything unfamiliar must be investigated thoroughly.

Correct but unfamiliar or unexpected methods are often signalled by a correct result following an *apparently* incorrect method. Such work must be carefully assessed. When a candidate adopts a method which does not correspond to the mark scheme, award marks according to the spirit of the basic scheme; if you are in any doubt whatsoever (especially if several marks or candidates are involved) you should contact your Team Leader.

- c The following types of marks are available.

M

A suitable method has been selected and *applied* in a manner which shows that the method is essentially understood. Method marks are not usually lost for numerical errors, algebraic slips or errors in units. However, it is not usually sufficient for a candidate just to indicate an intention of using some method or just to quote a formula; the formula or idea must be applied to the specific problem in hand, eg by substituting the relevant quantities into the formula. In some cases the nature of the errors allowed for the award of an M mark may be specified.

A

Accuracy mark, awarded for a correct answer or intermediate step correctly obtained. Accuracy marks cannot be given unless the associated Method mark is earned (or implied). Therefore M0 A1 cannot ever be awarded.

B

Mark for a correct result or statement independent of Method marks.

E

A given result is to be established or a result has to be explained. This usually requires more working or explanation than the establishment of an unknown result.

Unless otherwise indicated, marks once gained cannot subsequently be lost, eg wrong working following a correct form of answer is ignored. Sometimes this is reinforced in the mark scheme by the abbreviation isw. However, this would not apply to a case where a candidate passes through the correct answer as part of a wrong argument.

- d When a part of a question has two or more 'method' steps, the M marks are in principle independent unless the scheme specifically says otherwise; and similarly where there are several B marks allocated. (The notation 'dep *' is used to indicate that a particular mark is dependent on an earlier, asterisked, mark in the scheme.) Of course, in practice it may happen that when a candidate has once gone wrong in a part of a question, the work from there on is worthless so that no more marks can sensibly be given. On the other hand, when two or more steps are successfully run together by the candidate, the earlier marks are implied and full credit must be given.
- e The abbreviation ft implies that the A or B mark indicated is allowed for work correctly following on from previously incorrect results. Otherwise, A and B marks are given for correct work only — differences in notation are of course permitted. A (accuracy) marks are not given for answers obtained from incorrect working. When A or B marks are awarded for work at an intermediate stage of a solution, there may be various alternatives that are equally acceptable. In such cases, exactly what is acceptable will be detailed in the mark scheme rationale. If this is not the case please consult your Team Leader.

Sometimes the answer to one part of a question is used in a later part of the same question. In this case, A marks will often be 'follow through'. In such cases you must ensure that you refer back to the answer of the previous part question even if this is not shown within the image zone. You may find it easier to mark follow through questions candidate-by-candidate rather than question-by-question.

- f Wrong or missing units in an answer should not lead to the loss of a mark unless the scheme specifically indicates otherwise. Candidates are expected to give numerical answers to an appropriate degree of accuracy, with 3 significant figures often being the norm. Small variations in the degree of accuracy to which an answer is given (e.g. 2 or 4 significant figures where 3 is expected) should not normally be penalised, while answers which are grossly over- or under-specified should normally result in the loss of a mark. The situation regarding any particular cases where the accuracy of the answer may be a marking issue should be detailed in the mark scheme rationale. If in doubt, contact your Team Leader.
- g **Rules for replaced work**
If a candidate attempts a question more than once, and indicates which attempt he/she wishes to be marked, then examiners should do as the candidate requests.
If there are two or more attempts at a question which have not been crossed out, examiners should mark what appears to be the last (complete) attempt and ignore the others.
NB Follow these maths-specific instructions rather than those in the assessor handbook.

- h For a *genuine* misreading (of numbers or symbols) which is such that the object and the difficulty of the question remain unaltered, mark according to the scheme but following through from the candidate's data. A penalty is then applied; 1 mark is generally appropriate, though this may differ for some units. This is achieved by withholding one A mark in the question.

Note that a miscopy of the candidate's own working is not a misread but an accuracy error.

Question		Answer	Marks	Guidance
1	(i)	$x^2 - 3x + 2 = (x-1)(x-2)$ or $(1-x)(2-x)$ oe Obtain $-\frac{1}{x-2}$ or $\frac{1}{2-x}$ or $\frac{-1}{x-2}$ or $\frac{1}{-(x-2)}$ ISW If Partial Fractions are used, apply normal mark scheme.	B1 B1 [2]	Not $\frac{-1}{-(2-x)}$ Accept WW
1	(ii)	Attempt single fraction or 2 fractions with same relevant denom Fully correct fraction(s) before any simplification Relevant numerator = $3x-9$ or $3x^2-18x+27$ Final answer = $\frac{3}{(x-1)(x-4)}$ or $\frac{3}{x^2-5x+4}$ ISW S.R. If partial fractions are used on each fraction $-\frac{1}{x-1} + \frac{2}{x-3}$ $\frac{2}{x-3} - \frac{1}{x-4}$ $-\frac{1}{x-1} + \frac{1}{x-4}$ ISW	M1 A1 B1 A1 [4] (M1) (A1) (A1) (A1)	e.g. $(x-1)(x-4)[(x-3)$ or $(x-3)^2]$ Can award if no denominator
2		Write (or imply as) $\int 1 \cdot \ln(x+2)(dx)$ ($\ln x + \ln 2 \rightarrow M0$) Correct 'by parts' 1 st stage $x \ln(x+2) - \int \frac{x}{x+2}(dx)$ Any suitable <u>starting idea</u> for integrating $\frac{x}{x+2}$ [e.g. change num to $x+2-2$ or use substitution $x+2 = u$] $\int \frac{x}{x+2}(dx) = x - 2 \ln(x+2)$ or $x+2 - 2 \ln(x+2)$ Overall result = $x \ln(x+2) - x + 2 \ln(x+2)$ [(+c) or (-2+c)] ISW SR: Correct answer with no working	M1 A1 M1 A1 A1 [5] (B2)	OR: $t = \ln(x+2)$ and attempt to connect dx and dt $\int te^t (dt)$ Attempt by parts with $u = t$, $\frac{dv}{dt} = e^t$ $te^t - e^t$

Question	Answer	Marks	Guidance
3 (i)	<p>The first 5 marks are awarded for expansions of either $(1+4x)^{-\frac{1}{2}}$ or $(1+4x)^{\frac{1}{2}}$</p> <p>Expansion of $(1+4x)^{-\frac{1}{2}}$; First 2 terms = $1-2x$</p> <p>3rd term = $\frac{-\frac{1}{2} \cdot (-\frac{1}{2} - 1)}{2} \cdot 16x^2$ [Accept $4x^2$ for $16x^2$]</p> <p>= + $6x^2$</p> <p>4th term = $\frac{-\frac{1}{2} \cdot (-\frac{1}{2} - 1) \cdot (-\frac{1}{2} - 2)}{2 \cdot 3} \cdot 64x^3$ [Accept $4x^3$ for $64x^3$]</p> <p>= $-20x^3$</p> <p>$1-2x+7x^2-22x^3$; $1+ax+(b+1)x^2+(a+c)x^3$</p>	<p>B1</p> <p>M1</p> <p>A1</p> <p>M1</p> <p>A1</p> <p>A1 ft</p> <p>[6]</p>	<p><u>Or</u> $(1+4x)^{\frac{1}{2}} = 1+2x.....$</p> <p>3rd term = $\frac{\frac{1}{2} \cdot -\frac{1}{2}}{2} \cdot 16x^2$ [ditto]</p> <p>= $-2x^2$</p> <p>4th tm = $\frac{\frac{1}{2} \cdot -\frac{1}{2} \cdot -\frac{3}{2}}{2 \cdot 3} \cdot 64x^3$ [ditto]</p> <p>= + $4x^3$</p> <p>ft only $(1+4x)^{-\frac{1}{2}} = 1+ax+bx^2+cx^3$ provided a, b and c attempted and at least one @ M1 obtained</p>
3 (ii)	<p>$x < \frac{1}{4}$; $-\frac{1}{4} < x < \frac{1}{4}$; $\{-\frac{1}{4} < x, x < \frac{1}{4}\}$ no equality</p>	<p>B1</p> <p>[1]</p>	<p>But not $\{-\frac{1}{4} < x$ OR $x < \frac{1}{4}\}$ If choice mark what appears to be the final answer.</p>
4	<p>$+/- \int e^{2y} (dy)$ and $+/- \int \tan x (dx)$ seen</p> <p>$\int e^{2y} (dy) = \frac{1}{2} e^{2y}$</p> <p>$\int \tan x (dx) = \ln \sec x$ or $-\ln \cos x$</p> <p>Subst $x = 0, y = 0$ into their equation containing $f(x), g(y)$ and c</p> <p>$c = \frac{1}{2}$ WWW (or poss $-\frac{1}{2}$ if c on LHS)</p> <p>$y = \frac{1}{2} \ln(1 - 2 \ln \sec x)$ or $\frac{1}{2} \ln(1 + 2 \ln \cos x)$ oe WWW</p>	<p>M1</p> <p>B1</p> <p>B1</p> <p>M1</p> <p>A1</p> <p>A1</p> <p>[6]</p>	<p>may be implied later</p> <p>Accept $\ln \sec x$ or $-\ln \cos x$</p> <p>S.R. Using def integrals: M1 $\int_0^x = \int_0^y$ followed by A2 or A0</p> <p>Accept omission of modulus</p>

Question		Answer	Marks	Guidance	
5	(i)	Use $\cos \theta = \frac{a \cdot b}{ a b }$	M1	Better: 1.0471976 (rot)	
		Obtain $\left(\cos \theta = \frac{6}{12} \right) \theta = 60$ or $\frac{1}{3}\pi$ or 1.05 or better	A1		
			[2]		
5	(ii)	Indicate $\mathbf{a} - \mathbf{b}$ is vector joining ends of \mathbf{a} and \mathbf{b} or equiv $ \mathbf{a} - \mathbf{b} = \mathbf{a} - \mathbf{b} $, or anything similar, \rightarrow M0	M1	Or any other correct method 3.6055513 (rot)	
		Use cosine rule correctly on 3, 4 and included (i) angle	M1		
		Obtain $\sqrt{13}$ or 3.61 or better (No ft from wrong θ)	A1		
			[3]		
6		<u>Attempt</u> diff to connect du and dx or find $\frac{du}{dx}$ or $\frac{dx}{du}$	M1	<u>no</u> accuracy, <u>not</u> just $du = dx$	
		Correct e.g. $\frac{du}{dx} = \frac{1}{2}x^{-\frac{1}{2}}$ or $dx = (2u - 2)du$ AEF	*A1		
		Indefinite integral <u>in terms of u</u> = $\int \frac{2u-2}{u} (du)$	A1dep*		
		Provided of form $\int \frac{au+b}{u} (du)$, change to $\int a + \frac{b}{u} (du)$	M1		Or by parts
		Integrate to $au + b \ln u $ or $au + b \ln u$	A1 ft		
		Use correct variable for limits after attempt at integral of $f(u)$	M1		i.e. use new values of u (usually) or orig values of x (if resubst)
		Show as $8 - 2 \ln 4 - 6 + 2 \ln 3$ (oe) = $2 + 2 \ln \frac{3}{4}$ AG WWW	A1		Some 'numerical' working must be shown before giving final ans
			[7]		

Question	Answer	Marks	Guidance
7	<p>Satisfactory start method eg <u>attempt</u> square of $(1 - \sin 3x)$</p> <p>[N.B. The squaring process might include a term $\sin^2 9x$]</p> <p><u>The next 2 marks are awarded for integrating $-2\sin 3x$</u></p> <p>Obtain $\int -2 \sin 3x \, dx = \frac{2}{3} \cos 3x$</p> <p>Obtain $-\frac{2}{3}$ or $(\dots + 0\dots) - (\dots + \frac{2}{3}\dots)$</p> <p><u>The next 3 marks are awarded for integrating $\sin^2 3x$</u></p> <p>Use $\sin^2 3x = k(+/-1 +/- \cos 6x)$</p> <p>Correct version = $\frac{1}{2}(1 - \cos 6x)$</p> <p>$\int \cos 6x \, dx = \frac{1}{6} \sin 6x$, seen anywhere, indep</p> <p>Final answer = $\frac{1}{4}\pi + \textit{their} - \frac{2}{3}$</p>	<p>M1</p> <p>*A1</p> <p>A1dep*</p> <p>M1</p> <p>A1</p> <p>B1</p> <p>A1</p> <p>[7]</p>	<p>Not e.g. $\frac{(1 - \sin 3x)^3}{3}$.</p> <p>or for integrating $\sin^2 ax$ where $a = 6$ or 9 only</p> <p>$\sin^2 ax = k(+/-1 +/- \cos 2ax)$</p> <p>Correct = $\frac{1}{2}(1 - \cos 2ax)$</p> <p>or $\int \cos 2ax \, dx = \frac{1}{2a} \sin 2ax$</p> <p>Check that the $\frac{1}{4}\pi$ is from $\left[\frac{3}{2}x - \frac{1}{12} \sin 6x \right]_0^{\frac{1}{6}\pi}$</p>

Question			Answer	Marks	Guidance
8	(a)		$\frac{d}{dx}(xy) = x \frac{dy}{dx} + y$ $\frac{d}{dx}(y^2) = 2y \frac{dy}{dx}$ Substitute $(-1,-1)$ for (x, y) & attempt to solve for $\frac{dy}{dx}$ Obtain $\frac{dy}{dx} = -1$ WWW	B1 B1 M1 A1 [4]	or solve then substitute
8	(b)	(i)	Tangent parallel y -axis $\rightarrow \frac{dx}{dt} = 0$ or $\frac{dy}{dx} \rightarrow \infty$ or $\frac{dy}{dx} = \infty$ Obtain $t = 0$ $(-1,0)$ with no other possibilities	M1 A1 A1 [3]	Accept clear intention Accept $x = -1, y = 0$
8	(b)	(ii)	State or imply or use $\frac{dy}{dt} = \frac{dx}{dt}$ Produce $3t^2 + 1 = 4t$ oe $t = \frac{1}{3}$ or 1	M1 A1 A1 [3]	

Question	Answer	Marks	Guidance
9 (i)	$\frac{A}{x+1} + \frac{B}{x-2} + \frac{C}{(x-2)^2}$ $A(x-2)^2 + B(x+1)(x-2) + C(x+1) = x^2 - x - 11$ $A = -1$ $B = 2$ $C = -3$ <p><u>Special Cases</u></p> <p>The problems arise when we see how candidates deal with the denominator $(x-2)^2$:</p> $\frac{A}{x+1} + \frac{Bx+C}{(x-2)^2}$; allow B1 for PF format, M1 for associated identity, B1 for $A = -1$ (max 3) $\frac{A}{x+1} + \frac{B}{x-2} + \frac{Cx+D}{(x-2)^2}$; allow B1 for PF format, M1 for assoc identity, B1 for $A = -1$ (max 3) $\frac{A}{x+1} + \frac{Bx}{(x-2)^2}$; allow B0 for PF format, M1 for associated identity (max 1, even if $A = -1$) $\frac{A}{x+1} + \frac{B}{(x-2)^2}$; allow B0 for PF format, M1 for associated identity (max 1, even if $A = -1$)	<p>B1</p> <p>M1</p> <p>A1</p> <p>A1</p> <p>A1</p> <p>[5]</p>	<p>i.e. correct partial fractions</p> <p>or equivalent identity or method</p> <p>B1 if cover up method used</p> <p>B1 if cover up method used</p>
9 (ii)	<p>No marks are to be awarded for integrating a fraction with a zero numerator. Irrespective of the format used for the Partial Fractions in part (i), award marks as follow:</p> $\int \frac{\lambda}{x+1} dx = \left(\lambda \text{ or } \frac{1}{\lambda} \right) \ln(x+1) \quad \text{or....}$ $\int \frac{\mu}{(x-2)^2} dx = - \left(\mu \text{ or } \frac{1}{\mu} \right) \cdot \frac{1}{x-2}$ $-\frac{3}{2}$ <p>..... + $\ln \frac{16}{5}$ ISW for either term</p>	<p>B1</p> <p>B1</p> <p>B1 ft</p> <p>B1 ft</p> <p>[4]</p>	$\int \frac{\lambda}{x-2} dx = \left(\lambda \text{ or } \frac{1}{\lambda} \right) \ln(x-2)$ <p>ft $\frac{C}{2}$</p> <p>ft + $\ln \left\{ \left(\frac{5}{4} \right)^A \cdot 2^B \right\}$</p>

Question	Answer	Marks	Guidance
<p>10</p> <p>(i)</p>	<p>If MR, mark according to the scheme & follow-through from candidate's data. Award M, A & B marks (where possible) & apply penalty of 1 mark (by withholding one A mark in the question). E.g. in (i), product to be 'correct' & 'not perpendicular' to be stated.</p> <p>α. Full justification that $t = -1$. May be 'by inspection'. [No equations not satisfied by $t = -1$ to be shown] [‘unusual’ attempts must be carefully checked; if convinced, award the B1 e.g. displacement vector between $(-3\mathbf{i} + 6\mathbf{k})$ and $(-\mathbf{i} + 2\mathbf{j} + 7\mathbf{k}) = \pm(2\mathbf{i} + 2\mathbf{j} + \mathbf{k})$]</p> <p>β. Consider scalar product $\begin{pmatrix} -3 \\ 0 \\ 6 \end{pmatrix} \cdot \begin{pmatrix} 2 \\ 2 \\ 1 \end{pmatrix}$</p> <p>Show $-6 + (0) + 6 = 0$ and somewhere state perpendicularity oe</p> <p>[If $\cos \theta = \frac{\mathbf{a} \cdot \mathbf{b}}{ \mathbf{a} \mathbf{b} }$ quoted, ignore accuracy of work involving \mathbf{a} and \mathbf{b}]</p>	<p>B1</p> <p>M1</p> <p>A1</p> <p>[3]</p>	<p>No other $t =$ to be mentioned</p>
<p>10</p> <p>(ii)</p>	<p>Use $\mathbf{r} = v(-3\mathbf{i} + 6\mathbf{k})$ and ℓ_2</p> <p>Attempt to produce at least two relevant equations</p> <p>Solve two equations & produce $(v, s) = (\frac{1}{3}, -3)$ soi</p> <p>Demonstrate clearly that these satisfy third equation</p>	<p>*M1</p> <p>M1dep*</p> <p>A1</p> <p>B1</p> <p>[4]</p>	<p>or $(-3\mathbf{i} + 6\mathbf{k}) + v(-3\mathbf{i} + 6\mathbf{k})$</p> <p>$(v, s) = (-\frac{2}{3}, -3)$</p> <p>Numerical proof required</p>
<p>10</p> <p>(iii)</p>	<p>Method for finding \overrightarrow{OB} or \overrightarrow{OA} or \overrightarrow{AB}</p> <p>$\overrightarrow{OB} = \sqrt{5}$ or $\overrightarrow{OA} = \sqrt{45}$ oe or $\overrightarrow{BA} = \sqrt{20}$ oe</p> <p>Obtain 3:2 oe</p>	<p>M1</p> <p>A1</p> <p>A1</p> <p>[3]</p>	<p>Method for finding \overrightarrow{OB} or \overrightarrow{BO} or \overrightarrow{AB} or \overrightarrow{BA}</p> <p>$\overrightarrow{OB} = \begin{pmatrix} -1 \\ 0 \\ 2 \end{pmatrix}$ or $\overrightarrow{BA} = \begin{pmatrix} -2 \\ 0 \\ 4 \end{pmatrix}$</p> <p>Answer 3:2 WW \rightarrow B3</p>

OCR (Oxford Cambridge and RSA Examinations)
1 Hills Road
Cambridge
CB1 2EU

OCR Customer Contact Centre

Education and Learning

Telephone: 01223 553998

Facsimile: 01223 552627

Email: general.qualifications@ocr.org.uk

www.ocr.org.uk

For staff training purposes and as part of our quality assurance programme your call may be recorded or monitored

Oxford Cambridge and RSA Examinations
is a Company Limited by Guarantee
Registered in England
Registered Office; 1 Hills Road, Cambridge, CB1 2EU
Registered Company Number: 3484466
OCR is an exempt Charity

OCR (Oxford Cambridge and RSA Examinations)
Head office
Telephone: 01223 552552
Facsimile: 01223 552553

© OCR 2012

