

### Exercise 1A

- 1 B At least one multiple of three is odd.
- 2 a At least one rich person is not happy.
- b There is at least one prime number between 10 million and 11 million.
- c There exists prime numbers  $p$  and  $q$  such that  $(pq + 1)$  is not prime.
- d There is a number of the form  $2^n - 1$  that is either not prime or not a multiple of 3.
- e None of the above statements is true.
- 3 a There exists a number  $n$  such that  $n^2$  is odd but  $n$  is even.
- b Assume  $n$  is even, so write  $n = 2k$ , for some integer  $k$ .  
 $n^2 = (2k)^2 = 4k^2 = 2(2k^2)$   
 Since  $2k^2$  is an integer,  $n^2$  is even.  
 This contradicts the assumption that  $n^2$  is odd.  
 Therefore, if  $n^2$  is odd then  $n$  must be odd.
- 4 a Assumption: there is a greatest even integer,  $2n$ , for some integer  $n$ .  
 $2(n + 1)$  is also an even integer, since  $n + 1$  is an integer, and  
 $2(n + 1) > 2n$   
 So there exists an even integer greater than  $2n$ .  
 This contradicts the assumption that the greatest even integer is  $2n$ .  
 Therefore there is no greatest even integer.
- b Assumption: there exists a number  $n$  such that  $n^3$  is even but  $n$  is odd.  
 $n$  is odd so write  $n = 2k + 1$ , where  $k$  is an integer.  
 $n^3 = (2k + 1)^3$   
 $= 8k^3 + 12k^2 + 6k + 1$   
 $= 2(4k^3 + 6k^2 + 3k) + 1$   
 So  $n^3$  is odd, since  $4k^3 + 6k^2 + 3k$  is an integer.  
 This contradicts the assumption that  $n^3$  is even.  
 Therefore, if  $n^3$  is even then  $n$  must be even.
- c Assumption: if  $pq$  is even then neither  $p$  nor  $q$  is even.  
 Assume  $p$  is odd,  $p = 2k + 1$ , where  $k$  is an integer.  
 Assume  $q$  is odd,  $q = 2m + 1$ , where  $m$  is an integer.  
 $pq = (2k + 1)(2m + 1)$   
 $= 2km + 2k + 2m + 1$   
 $= 2(km + k + m) + 1$   
 So  $pq$  is odd, since  $km + k + m$  is an integer.  
 This contradicts the assumption that  $pq$  is even.  
 Therefore, if  $pq$  is even then at least one of  $p$  and  $q$  is even.

- 4 d** Assumption: if  $p + q$  is odd then neither  $p$  nor  $q$  is odd.  
 Assume  $p$  is even,  $p = 2k$ , where  $k$  is an integer.  
 Assume  $q$  is even,  $q = 2m$ , where  $m$  is an integer.  
 $p + q = 2k + 2m = 2(k + m)$   
 So  $p + q$  is even, since  $k + m$  is an integer.  
 This contradicts the assumption that  $p + q$  is odd.  
 Therefore, if  $p + q$  is odd then at least one of  $p$  and  $q$  is odd.
- 5 a** Assumption: if  $ab$  is an irrational number then neither  $a$  nor  $b$  is irrational.  
 If  $a$  is rational, then  $a = \frac{c}{d}$  where  $c$  and  $d$  are integers.  
 If  $b$  is rational, then  $b = \frac{e}{f}$  where  $e$  and  $f$  are integers.  
 Then  $ab = \frac{ce}{df}$ , where  $ce$  is an integer,  $df$  is an integer.  
 Therefore  $ab$  is a rational number.  
 This contradicts assumption that  $ab$  is irrational.  
 Therefore, if  $ab$  is an irrational number then at least one of  $a$  and  $b$  is an irrational number.
- b** Assumption: if  $a + b$  is an irrational number then neither  $a$  nor  $b$  is irrational.  
 If  $a$  is rational, then  $a = \frac{c}{d}$  where  $c$  and  $d$  are integers.  
 If  $b$  is rational, then  $b = \frac{e}{f}$  where  $e$  and  $f$  are integers.  
 Then  $a + b = \frac{cf + de}{df}$ , where  $cf$ ,  $de$  and  $df$  are integers.  
 So  $a + b$  is rational. This contradicts the assumption that  $a + b$  is irrational.  
 Therefore if  $a + b$  is irrational then at least one of  $a$  and  $b$  is irrational.
- c** Many possible answers  
 e.g.  $a = 2 - \sqrt{2}$ ,  $b = \sqrt{2}$ , or  $a = e$ ,  $b = -e$ .
- 6** Assumption: there exist integers  $a$  and  $b$  such that  $21a + 14b = 1$ .  
 Since 21 and 14 are multiples of 7, divide both sides by 7.  
 So now  $3a + 2b = \frac{1}{7}$   
 $3a$  is also an integer.  $2b$  is also an integer.  
 The sum of two integers will always be an integer, so  $3a + 2b$  is an integer.  
 This contradicts the statement that  
 $3a + 2b = \frac{1}{7}$   
 Therefore there exist no integers  $a$  and  $b$  for which  $21a + 14b = 1$ .

- 7 a Assumption: There exists a number  $n$  such that  $n^2$  is a multiple of 3, but  $n$  is not a multiple of 3.

All multiples of 3 can be written in the form  $n = 3k$  where  $k$  is an integer, therefore  $3k + 1$  and  $3k + 2$  are not multiples of 3.

Let  $n = 3k + 1$

$$\begin{aligned} n^2 &= (3k + 1)^2 \\ &= 9k^2 + 6k + 1 \\ &= 3(3k^2 + 2k) + 1 \end{aligned}$$

In this case  $n^2$  is not a multiple of 3, since  $3k^2 + 2k$  is an integer.

Let  $n = 3k + 2$

$$\begin{aligned} n^2 &= (3k + 2)^2 \\ &= 9k^2 + 12k + 4 \\ &= 3(3k^2 + 4k + 1) + 1 \end{aligned}$$

In this case  $n^2$  is also not a multiple of 3, since  $3k^2 + 4k + 1$  is an integer.

This contradicts the assumption that  $n^2$  is a multiple of 3.

Therefore if  $n^2$  is a multiple of 3,  $n$  is a multiple of 3.

- b Assumption:  $\sqrt{3}$  is a rational number.

Then  $\sqrt{3} = \frac{a}{b}$  for some integers  $a$  and  $b$ .

Further assume that this fraction is in its simplest terms: there are no common factors between  $a$  and  $b$ .

$$\text{So } 3 = \frac{a^2}{b^2} \text{ or } a^2 = 3b^2$$

Therefore  $a^2$  must be a multiple of 3.

We know from part a that this means  $a$  must also be a multiple of 3.

Write  $a = 3c$ , which means  $a^2 = (3c)^2 = 9c^2$ .

Now  $9c^2 = 3b^2$ , or  $3c^2 = b^2$ .

Therefore  $b^2$  must be a multiple of 3, which means  $b$  is also a multiple of 3.

If  $a$  and  $b$  are both multiples of 3, this contradicts the statement that there are no common factors between  $a$  and  $b$ .

Therefore,  $\sqrt{3}$  is an irrational number.

- 8 Assumption: there is an integer solution to the equation  $x^2 - y^2 = 2$ .

Remember that  $x^2 - y^2 = (x - y)(x + y) = 2$ .

To make a product of 2 using integers, the possible pairs are: (2, 1), (1, 2), (-2, -1) and (-1, -2). Since  $(-x)^2 = x^2$ , we may assume that  $x$  is positive, and similarly for  $y$ . So we only need to consider the possibilities (2, 1) and (1, 2).

Consider each possibility in turn:

$$x - y = 2 \text{ and } x + y = 1 \Rightarrow x = \frac{3}{2}, y = -\frac{1}{2}$$

$$x - y = 1 \text{ and } x + y = 2 \Rightarrow x = \frac{3}{2}, y = \frac{1}{2}$$

This contradicts the statement that there is an integer solution to the equation  $x^2 - y^2 = 2$ .

Therefore the original statement must be true:

There are no integer solutions to the equation  $x^2 - y^2 = 2$ .

9 Assumption:  $\sqrt[3]{2}$  is a rational number

Then  $\sqrt[3]{2} = \frac{a}{b}$  for some integers  $a$  and  $b$ .

Further assume that this fraction is in its simplest terms:  
there are no common factors between  $a$  and  $b$ .

If  $\sqrt[3]{2} = \frac{a}{b}$  then  $2 = \frac{a^3}{b^3}$  or  $2b^3 = a^3$ .

This means that  $a^3$  is even, since  $b^3$  is an integer, and so  $a$  must also be even.

If  $a$  is even,  $a = 2n$ .

So  $a^3 = 2b^3$  becomes  $(2n)^3 = 2b^3$  which means  $8n^3 = 2b^3$  or  $4n^3 = b^3$  or  $2(2n^3) = b^3$ .

This means that  $b^3$  must be even, since  $2n^3$  is an integer, so  $b$  is also even.

If  $a$  and  $b$  are both even, they will have a common factor of 2.

This contradicts the statement that  $a$  and  $b$  have no common factors.

We can conclude the original statement is true:  $\sqrt[3]{2}$  is an irrational number.

10 a The number  $\frac{a-b}{b}$  could be negative.

e.g. If  $n = \frac{1}{2}$ ,  $n - 1$  is non-positive.

b Assumption: There is a least positive rational number,  $n$ .

$n = \frac{a}{b}$  where  $a$  and  $b$  are integers.

Let  $m = \frac{a}{2b}$ . Since  $a$  and  $b$  are integers,  $m$  is rational and  $m < n$ .

This contradicts the statement that  $n$  is the least positive rational number.

Therefore, there is no least positive rational number.