

Exercise 3B

1 a $x = 2 \sin t - 1$

$$\text{So } \sin t = \frac{x+1}{2} \quad (1)$$

$$y = 5 \cos t + 4$$

$$\cos t = \frac{y-4}{5} \quad (2)$$

Substitute (1) and (2) into

$$\sin^2 t + \cos^2 t \equiv 1:$$

$$\left(\frac{x+1}{2}\right)^2 + \left(\frac{y-4}{5}\right)^2 = 1$$

$$\frac{(x+1)^2}{4} + \frac{(y-4)^2}{25} = 1$$

$$25(x+1)^2 + 4(y-4)^2 = 100$$

b $y = \sin 2t$

$$= 2 \sin t \cos t$$

So, since $x = \cos t$,

$$y = 2x \sin t \quad (1)$$

$$\sin^2 t + \cos^2 t \equiv 1$$

$$\sin^2 t \equiv 1 - \cos^2 t = 1 - x^2$$

$$\sin t = \sqrt{1-x^2} \quad (2)$$

Substitute (2) into (1):

$$y = 2x\sqrt{1-x^2}$$

$$\text{or } y^2 = 4x^2(1-x^2)$$

c $y = 2 \cos 2t$

$$= 2(2 \cos^2 t - 1)$$

So, since $x = \cos t$,

$$y = 2(2x^2 - 1)$$

$$y = 4x^2 - 2$$

d $y = \tan 2t$

$$\text{So } y = \frac{2 \tan t}{1 - \tan^2 t} \quad (1)$$

$$\sin^2 t + \cos^2 t \equiv 1$$

$$\cos^2 t \equiv 1 - \sin^2 t = 1 - x^2$$

$$\cos t = \sqrt{1-x^2} \quad (2)$$

Substitute (2) and $x = \sin t$ into (1):

$$\begin{aligned} y &= \frac{2 \frac{\sin t}{\cos t}}{1 - \frac{\sin^2 t}{\cos^2 t}} = \frac{\frac{2x}{\sqrt{1-x^2}}}{1 - \frac{x^2}{1-x^2}} = \frac{\frac{2x}{\sqrt{1-x^2}}}{\frac{1-2x^2}{1-x^2}} \\ &= \frac{2x(1-x^2)}{(1-2x^2)\sqrt{1-x^2}} \end{aligned}$$

$$\text{Hence } y = \frac{2x\sqrt{1-x^2}}{1-2x^2}$$

e $x = \cos t + 2$

$$\cos t = x - 2 \quad (1)$$

$$y = 4 \sec t = \frac{4}{\cos t}$$

$$\cos t = \frac{4}{y} \quad (2)$$

Substitute (1) into (2):

$$x-2 = \frac{4}{y}$$

$$y = \frac{4}{x-2}$$

f $x = 3 \cot t$

$$\cot t = \frac{x}{3} \quad (1)$$

$$\operatorname{cosec} t = y \quad (2)$$

Substitute (1) and (2) into

$$1 + \cot^2 t \equiv \operatorname{cosec}^2 t :$$

$$1 + \left(\frac{x}{3}\right)^2 = y^2$$

$$y^2 = 1 + \frac{x^2}{9}$$

2 a $x = \sin t - 5$
 $\Rightarrow \sin t = x + 5$ (1)
 $y = \cos t + 2$
 $\Rightarrow \cos t = y - 2$ (2)
 Substitute (1) and (2) into
 $\sin^2 t + \cos^2 t \equiv 1$:
 $(x+5)^2 + (y-2)^2 = 1$

- b** This is a circle with centre $(-5, 2)$ and radius 1
- c** One full revolution around the circle is obtained for an interval of t corresponding to one period of both parametric equations $y = \cos t + 2$ and $x = \sin t - 5$.
 So $0 \leq t \leq 2\pi$ is a suitable domain.

3 $x = 4 \sin t + 3$
 $4 \sin t = x - 3$
 $\therefore \sin t = \frac{x-3}{4}$ (1)

$$y = 4 \cos t - 1$$

$$4 \cos t = y + 1$$

$$\therefore \cos t = \frac{y+1}{4}$$
 (2)

Substitute (1) and (2) into
 $\sin^2 t + \cos^2 t = 1$:

$$\left(\frac{x-3}{4}\right)^2 + \left(\frac{y+1}{4}\right)^2 = 1$$


$$\frac{(x-3)^2}{4^2} + \frac{(y+1)^2}{4^2} = 1$$

$$\frac{(x-3)^2}{16} + \frac{(y+1)^2}{16} = 1$$

$$(x-3)^2 + (y+1)^2 = 16$$

So the radius of the circle is 4 and the centre is $(3, -1)$.

4 $x = \cos t - 2$
 $\Rightarrow \cos t = x + 2$ (1)
 $y = \sin t + 3$
 $\Rightarrow \sin t = y - 3$ (2)
 Substitute (1) and (2) into $\sin^2 t + \cos^2 t \equiv 1$:
 $(x+2)^2 + (y-3)^2 = 1$
 This is a circle with centre $(-2, 3)$ and radius 1:



5 a $y = \sin\left(t + \frac{\pi}{4}\right)$
 $= \sin t \cos \frac{\pi}{4} + \cos t \sin \frac{\pi}{4}$
 $= \frac{\sqrt{2}}{2} \sin t + \frac{\sqrt{2}}{2} \cos t$
 $y = \frac{\sqrt{2}}{2} x + \frac{\sqrt{2}}{2} \cos t$ (1)
 (since $x = \sin t$)

$$\begin{aligned} & \sin^2 t + \cos^2 t \equiv 1 \\ & \cos^2 t \equiv 1 - \sin^2 t = 1 - x^2 \\ & \therefore \cos t = \sqrt{1-x^2} \end{aligned}$$
 (2)

Substitute (2) into (1):

$$\begin{aligned} & y = \frac{\sqrt{2}}{2} x + \frac{\sqrt{2}}{2} \sqrt{1-x^2} \\ & \text{or } y = \frac{\sqrt{2}}{2} x + \frac{\sqrt{2(1-x^2)}}{2} \end{aligned}$$

5 b $x = 3 \cos t$

$$\Rightarrow \cos t = \frac{x}{3}$$

$$y = 2 \cos\left(t + \frac{\pi}{6}\right)$$

$$= 2 \cos t \cos \frac{\pi}{6} - 2 \sin t \sin \frac{\pi}{6}$$

$$= 2 \cos t \times \frac{\sqrt{3}}{2} - 2 \sin t \times \frac{1}{2}$$

$$= \sqrt{3} \cos t - \sin t$$

$$\text{So } y = \frac{\sqrt{3}}{3} x - \sin t \quad (1)$$

$$\sin^2 t + \cos^2 t \equiv 1$$

$$\sin^2 t \equiv 1 - \cos^2 t = 1 - \left(\frac{x}{3}\right)^2$$

$$\sin t = \sqrt{1 - \left(\frac{x}{3}\right)^2} \quad (2)$$

Substitute (2) into (1):

$$\begin{aligned} y &= \frac{\sqrt{3}}{3} x - \sqrt{1 - \left(\frac{x}{3}\right)^2} \\ &= \frac{\sqrt{3}}{3} x - \sqrt{\frac{9-x^2}{9}} \\ \therefore y &= \frac{\sqrt{3}}{3} x - \frac{\sqrt{9-x^2}}{3} \end{aligned}$$

c $y = 3 \sin(t + \pi)$

$$= 3 \sin t \cos \pi + 3 \cos t \sin \pi$$

$$= 3 \sin t \times (-1) + 3 \cos t \times 0$$

$$= -3 \sin t$$

Since $x = \sin t$,

$$y = -3x$$

6 a $x = 8 \cos t$

$$\cos t = \frac{x}{8}$$

$$\text{So } y = \frac{1}{4} \sec^2 t = \frac{1}{4 \cos^2 t}$$

$$= \frac{1}{4 \left(\frac{x}{8}\right)^2} = \frac{1}{4} \times \frac{64}{x^2} = \frac{16}{x^2}$$

Therefore a Cartesian equation for C is


$$y = \frac{16}{x^2}$$

b For $-\frac{\pi}{2} \leq t \leq \frac{\pi}{2}$ the range of the parametric equation $x = 8 \cos t$ is $0 \leq x \leq 8$, so the domain of $y = f(x)$ is $0 \leq x \leq 8$.

The range of the parametric equation

$y = \frac{1}{4} \sec^2 t$ is $y \geq \frac{1}{4}$, so the range of

$y = f(x)$ is $y \geq \frac{1}{4}$



7 $x = 3 \cot^2 2t$

$$\cot^2 2t = \frac{x}{3}$$

$$\frac{x}{3} = \frac{\cos^2 2t}{\sin^2 2t} = \frac{1 - \sin^2 2t}{\sin^2 2t} = \frac{1}{\sin^2 2t} - 1$$

$$\frac{x}{3} + 1 = \frac{1}{\sin^2 2t}$$

$$\frac{x+3}{3} = \frac{1}{\sin^2 2t}$$

$$\sin^2 2t = \frac{3}{x+3}$$

$$\therefore y = 3 \sin^2 2t = 3 \times \frac{3}{x+3} = \frac{9}{x+3}$$

For $0 < t \leq \frac{\pi}{4}$ the range of the parametric

function $x = 3 \cot^2 2t$ is $x \geq 0$, so the domain of $f(x)$ is $x \geq 0$.

8 a

$$\begin{aligned}x &= \frac{1}{3} \sin t \\ \Rightarrow \sin t &= 3x \\ y &= \sin 3t = \sin(t+2t) \\ &= \sin t \cos 2t + \cos t \sin 2t \\ &= \sin t(1-2\sin^2 t) + \cos t(2\sin t \cos t) \\ &= \sin t(1-2\sin^2 t) + 2\sin t(1-\sin^2 t) \\ &= 3x(1-2\times 9x^2) + 6x(1-9x^2) \\ &= 3x - 54x^3 + 6x - 54x^3 \\ &= 9x - 108x^3 \\ &= 9x(1-12x^2)\end{aligned}$$

So the Cartesian equation of the curve is $y = 9x(1-12x^2)$, which is in the form $y = ax(1-bx^2)$ with $a = 9$ and $b = 12$.

- b** For $0 < t < \frac{\pi}{2}$ the range of the parametric function $x = \frac{1}{3} \sin t$ is $0 < x < \frac{1}{3}$ so the domain of $y = f(x)$ is $0 < x < \frac{1}{3}$
- For $0 < t < \frac{\pi}{2}$ the range of the parametric function $y = \sin 3t$ is $-1 < y < 1$ so the range of $y = f(x)$ is $-1 < y < 1$.

9

$$\begin{aligned}x &= 2 \cos t \Rightarrow \cos t = \frac{x}{2} \\ y &= \sin\left(t - \frac{\pi}{6}\right) \\ &= \sin t \cos \frac{\pi}{6} - \cos t \sin \frac{\pi}{6} \\ &= \frac{\sqrt{3}}{2} \sin t - \frac{1}{2} \cos t \\ \therefore y &= \frac{\sqrt{3}}{2} \sin t - \frac{1}{4} x \quad (1) \\ \sin^2 t + \cos^2 t &\equiv 1 \\ \sin^2 t &\equiv 1 - \cos^2 t = 1 - \frac{x^2}{4} \\ \therefore \sin t &= \sqrt{1 - \frac{x^2}{4}} \quad (2)\end{aligned}$$

Substitute (2) into (1):

$$\begin{aligned}y &= \frac{\sqrt{3}}{2} \sqrt{1 - \frac{x^2}{4}} - \frac{1}{4} x \\ &= \frac{1}{2} \sqrt{\frac{12-3x^2}{4}} - \frac{1}{4} x\end{aligned}$$

So the Cartesian equation is

$$y = \frac{1}{4} \left(\sqrt{12-3x^2} - x \right)$$

For $0 < t < \pi$, the range of the parametric function $x = 2 \cos t$ is $-2 < x < 2$, so the domain of $y = f(x)$ is $-2 < x < 2$.

10 a $y = 5 \sin t$

$$\text{So } \sin t = \frac{y}{5}$$

$$\sin^2 t = \frac{y^2}{25}$$

$$x = \tan^2 t + 5$$

$$\tan^2 t = x - 5$$

$$\frac{\sin^2 t}{\cos^2 t} = x - 5$$

$$\frac{\sin^2 t}{1 - \sin^2 t} = x - 5$$

$$\frac{1}{x-5} = \frac{1}{\sin^2 t} - 1$$

$$\frac{1}{x-5} + 1 = \frac{1}{\left(\frac{y^2}{25}\right)}$$

$$\frac{x-4}{x-5} = \frac{25}{y^2}$$

$$\therefore y^2 = 25 \left(\frac{x-5}{x-4} \right) = 25 \left(1 - \frac{1}{x-4} \right)$$

b For $0 < t < \frac{\pi}{2}$, the range of the parametric

function $x = \tan^2 t + 5$ is $x > 5$,

so the domain of the curve is $x > 5$.

The range of the parametric function

$y = 5 \sin t$ is $0 < y < 5$,

so the range of the curve is $0 < y < 5$.

11 $y = 3 \sin(t - \pi)$

$$= 3 \sin t \cos \pi - 3 \cos t \sin \pi$$

$$= 3 \sin t \times (-1) - 3 \cos t \times 0$$

$$= -3 \sin t$$

$$\text{So } \sin t = -\frac{y}{3}$$

$$\sin^2 t + \cos^2 t \equiv 1$$

$$\cos^2 t \equiv 1 - \sin^2 t$$

$$\cos t = \sqrt{1 - \sin^2 t} = \sqrt{1 - \frac{y^2}{9}}$$

$$\therefore x = \tan t = \frac{\sin t}{\cos t} = \frac{-\frac{y}{3}}{\sqrt{1 - \frac{y^2}{9}}} = \frac{-y}{\sqrt{9 - y^2}}$$

Therefore $x = -\frac{y}{\sqrt{9 - y^2}}$ is a Cartesian equation for C.

Challenge

$$\begin{aligned}x &= \frac{1}{2} \cos 2t \\&= \frac{1}{2}(2 \cos^2 t - 1) \\&\therefore \frac{2x+1}{2} = \cos^2 t \\&\cos t = \sqrt{\frac{2x+1}{2}}\end{aligned}$$

But also

$$\begin{aligned}x &= \frac{1}{2} \cos 2t \\&= \frac{1}{2}(1 - 2 \sin^2 t) \\&\therefore \sin^2 t = \frac{1-2x}{2} \\&\sin t = \sqrt{\frac{1-2x}{2}}\end{aligned}$$

Therefore

$$\begin{aligned}y &= \sin\left(t + \frac{\pi}{6}\right) \\&= \sin t \cos \frac{\pi}{6} + \cos t \sin \frac{\pi}{6} \\&= \frac{\sqrt{3}}{2} \sin t + \frac{1}{2} \cos t \\&= \frac{\sqrt{3}}{2} \sqrt{\frac{1-2x}{2}} + \frac{1}{2} \sqrt{\frac{2x+1}{2}} \\&= \sqrt{\frac{3-6x}{8}} + \sqrt{\frac{2x+1}{8}}\end{aligned}$$

So a Cartesian equation for the curve is

$$y = \sqrt{\frac{3-6x}{8}} + \sqrt{\frac{2x+1}{8}}$$