

Exercise 5C

$$1 \quad A = \frac{1}{4}\pi r^2$$

$$\frac{dA}{dr} = \frac{1}{2}\pi r$$

$$\begin{aligned} \frac{dA}{dt} &= \frac{dA}{dr} \times \frac{dr}{dt} \\ &= \frac{1}{2}\pi r \times 6 = 3\pi r \end{aligned}$$

When $r = 2$,

$$\frac{dA}{dt} = 3\pi \times 2 = 6\pi$$

$$2 \quad y = xe^x$$

$$\frac{dy}{dx} = xe^x + e^x = (x+1)e^x$$

$$\begin{aligned} \frac{dy}{dt} &= \frac{dy}{dx} \times \frac{dx}{dt} \\ &= (x+1)e^x \times 5 = 5(x+1)e^x \end{aligned}$$

When $x = 2$,

$$\frac{dy}{dt} = 5(2+1)e^2 = 15e^2$$

$$3 \quad r = 1 + 3\cos\theta$$

$$\frac{dr}{d\theta} = -3\sin\theta$$

$$\begin{aligned} \frac{dr}{dt} &= \frac{dr}{d\theta} \times \frac{d\theta}{dt} \\ &= -3\sin\theta \times 3 = -9\sin\theta \end{aligned}$$

When $\theta = \frac{\pi}{6}$,

$$\frac{dr}{dt} = -9\sin\frac{\pi}{6} = -\frac{9}{2}$$

$$4 \quad V = \frac{1}{3}\pi r^3$$

$$\frac{dV}{dr} = \pi r^2 \Rightarrow \frac{dr}{dV} = \frac{1}{\pi r^2}$$

$$\begin{aligned} \frac{dr}{dt} &= \frac{dr}{dV} \times \frac{dV}{dt} \\ &= \frac{1}{\pi r^2} \times 8 = \frac{8}{\pi r^2} \end{aligned}$$

When $r = 3$,

$$\frac{dr}{dt} = \frac{8}{\pi \times 3^2} = \frac{8}{9\pi}$$

5 Let P be the size of the population and let t be time. Then the rate of growth of the population is $\frac{dP}{dt}$

$$\frac{dP}{dt} \propto P$$

i.e. $\frac{dP}{dt} = kP$ where k is a positive constant.

6 The gradient of the curve is $\frac{dy}{dx}$

$$\therefore \frac{dy}{dx} \propto xy \text{ (product of } x\text{- and } y\text{-coordinates)}$$

$$\text{i.e. } \frac{dy}{dx} = kxy,$$

where k is the constant of proportion.

When $x = 4$, $y = 2$ and $\frac{dy}{dx} = \frac{1}{2}$.

Substituting into $\frac{dy}{dx} = kxy$ gives

$$\frac{1}{2} = k \times 4 \times 2$$

$$k = \frac{1}{16}$$

$$\therefore \frac{dy}{dx} = \frac{xy}{16}$$

7 The rate of increase of the volume of liquid in the container is $\frac{dV}{dt}$

$$\frac{dV}{dt} = \text{rate in} - \text{rate out}$$

$$= 30 - \frac{2}{15}V$$

Multiply both sides by -15 to give

$$-15 \frac{dV}{dt} = 2V - 450$$

8 The rate of change of the charge is $\frac{dQ}{dt}$

$$\frac{dQ}{dt} \propto Q$$

$$\text{i.e. } \frac{dQ}{dt} = -kQ$$

where k is a positive constant.
(The negative sign is required as the body is losing charge.)

9 The rate of increase of x is $\frac{dx}{dt}$

$$\therefore \frac{dx}{dt} \propto \frac{1}{x^2} \quad (\text{inverse proportion})$$

$$\text{i.e. } \frac{dx}{dt} = \frac{k}{x^2}$$

where k is the constant of proportion.

10 a Let r be the radius of the circle and let t be time. Then $\frac{dr}{dt} = 0.4 \text{ cm s}^{-1}$.

$$C = 2\pi r$$

$$\frac{dC}{dr} = 2\pi$$

$$\frac{dC}{dt} = \frac{dC}{dr} \times \frac{dr}{dt}$$

$$= 2\pi \times 0.4 = 0.8\pi \text{ cm s}^{-1}$$

This means that the circumference is increasing at a constant rate of $0.8\pi \text{ cm}$ per second.

b Let A be the radius of the circle; then

$$A = \pi r^2$$

$$\frac{dA}{dr} = 2\pi r$$

$$\frac{dA}{dt} = \frac{dA}{dr} \times \frac{dr}{dt}$$

$$= 2\pi r \times 0.4 = 0.8\pi r$$

When $r = 10$,

$$\frac{dA}{dt} = 0.8\pi \times 10 = 8\pi \text{ cm}^2 \text{ s}^{-1}$$

$$\text{c } \frac{dA}{dt} = 0.8\pi r$$

$$\text{When } \frac{dA}{dt} = 20,$$

$$0.8\pi r = 20$$

$$r = \frac{20}{0.8\pi} = \frac{25}{\pi} \text{ cm}$$

11 a Let l be the side length of the cube and let V be its volume.

$$\text{Then } V = l^3 \text{ and } \frac{dV}{dt} = -4.5$$

$$\frac{dV}{dl} = 3l^2 \text{ so } \frac{dl}{dV} = \frac{1}{3l^2}$$

$$\frac{dl}{dt} = \frac{dl}{dV} \times \frac{dV}{dt}$$

$$= \frac{1}{3l^2} \times (-4.5) = -\frac{3}{2l^2}$$

$$\text{When } V = 100, l = \sqrt[3]{100}$$

$$\frac{dl}{dt} = -\frac{3}{2(\sqrt[3]{100})^2} = 0.070 \text{ cm s}^{-1} \quad (2 \text{ s.f.})$$

$$\text{b } \frac{dl}{dt} = -\frac{3}{2l^2}$$

2 mm is 0.2 cm

$$\text{When } \frac{dl}{dt} = -0.2,$$

$$-\frac{3}{2l^2} = -0.2$$

$$2l^2 = 15$$

$$l = \sqrt{7.5}$$

$$\therefore V = l^3 = (\sqrt{7.5})^3 = 20.5 \text{ cm}^3 \quad (3 \text{ s.f.})$$

12 The rate of change of the volume of fluid in the tank is $\frac{dV}{dt}$

$$\frac{dV}{dt} \propto \sqrt{V}$$

$$\text{i.e. } \frac{dV}{dt} = -K\sqrt{V}$$

where K is a positive constant.

(The negative sign is present because fluid is flowing *out* of the tank, so the volume left in the tank is *decreasing*.)

Let A be the constant cross-section; then $V = Ah$ (where h is the depth)

$$\therefore \frac{dV}{dh} = A$$

Use the chain rule to find $\frac{dh}{dt}$:

$$\begin{aligned} \frac{dV}{dt} &= \frac{dV}{dh} \times \frac{dh}{dt} \\ \therefore \frac{dh}{dt} &= \frac{dV}{dt} \div \frac{dV}{dh} \\ &= \frac{-K\sqrt{V}}{A} = \frac{-K\sqrt{Ah}}{A} \\ &= \left(\frac{-K}{\sqrt{A}} \right) \sqrt{h} = -k\sqrt{h} \end{aligned}$$

where $k = \frac{K}{\sqrt{A}}$ is a positive constant.

13 a Let l be the length of one side of the cube. Surface area of cube $A = 6l^2$.

$$\text{So } l = \sqrt{\frac{A}{6}}$$

$$\text{Volume of cube } V = l^3 = \left(\sqrt{\frac{A}{6}} \right)^3 = \left(\frac{A}{6} \right)^{\frac{3}{2}}$$

$$\text{b } \frac{dV}{dA} = \frac{1}{6} \times \frac{3}{2} \left(\frac{A}{6} \right)^{\frac{1}{2}} = \frac{1}{4} \left(\frac{A}{6} \right)^{\frac{1}{2}}$$

c Rate of expansion of surface area is $\frac{dA}{dt}$

$$\text{Given } \frac{dA}{dt} = 2$$

Need $\frac{dV}{dt}$ so use the chain rule:


$$\begin{aligned} \frac{dV}{dt} &= \frac{dV}{dA} \times \frac{dA}{dt} \\ &= \frac{1}{4} \left(\frac{A}{6} \right)^{\frac{1}{2}} \times 2 = \frac{1}{2} \left(\frac{A}{6} \right)^{\frac{1}{2}} \end{aligned}$$

From $A = 6l^2$ and $V = l^3$,

$$A = 6 \left(\sqrt[3]{V} \right)^2 = 6V^{\frac{2}{3}}$$

$$\therefore \frac{dV}{dt} = \frac{1}{2} \left(\frac{6V^{\frac{2}{3}}}{6} \right)^{\frac{1}{2}} = \frac{1}{2} V^{\frac{1}{3}}$$

14



Let V be the volume of salt in the funnel at time t .

$$V = \frac{1}{3} \pi r^2 h$$

$$\tan 30^\circ = \frac{r}{h} \Rightarrow r = h \tan 30^\circ = \frac{h}{\sqrt{3}}$$

$$\therefore V = \frac{1}{3} \pi \left(\frac{h^2}{3} \right) h = \frac{1}{9} \pi h^3$$

$$\frac{dV}{dh} = \frac{1}{3} \pi h^2 \text{ and hence } \frac{dh}{dV} = \frac{3}{\pi h^2}$$

Given that $\frac{dV}{dt} = -6$

$$\frac{dh}{dt} = \frac{dh}{dV} \times \frac{dV}{dt} = \frac{3}{\pi h^2} \times (-6) = -\frac{18}{\pi h^2}$$

So the rate of change of the height, h , is inversely proportional to h^2 and is given by

the differential equation $\frac{dh}{dt} = -\frac{18}{\pi h^2}$