

Core 4 Vectors Questions

- 7 The quadrilateral $ABCD$ has vertices $A(2, 1, 3)$, $B(6, 5, 3)$, $C(6, 1, -1)$ and $D(2, -3, -1)$.

The line l_1 has vector equation $\mathbf{r} = \begin{bmatrix} 6 \\ 1 \\ -1 \end{bmatrix} + \lambda \begin{bmatrix} 1 \\ 1 \\ 0 \end{bmatrix}$.

- (a) (i) Find the vector \overrightarrow{AB} . (2 marks)
- (ii) Show that the line AB is parallel to l_1 . (1 mark)
- (iii) Verify that D lies on l_1 . (2 marks)
- (b) The line l_2 passes through $D(2, -3, -1)$ and $M(4, 1, 1)$.
- (i) Find the vector equation of l_2 . (2 marks)
- (ii) Find the angle between l_2 and AC . (3 marks)
-

- 6 The points A and B have coordinates $(2, 4, 1)$ and $(3, 2, -1)$ respectively. The point C is such that $\overrightarrow{OC} = 2\overrightarrow{OB}$, where O is the origin.

- (a) Find the vectors:
- (i) \overrightarrow{OC} ; (1 mark)
- (ii) \overrightarrow{AB} . (2 marks)
- (b) (i) Show that the distance between the points A and C is 5. (2 marks)
- (ii) Find the size of angle BAC , giving your answer to the nearest degree. (4 marks)
- (c) The point $P(\alpha, \beta, \gamma)$ is such that BP is perpendicular to AC .
- Show that $4\alpha - 3\gamma = 15$. (3 marks)
-

6 The points A , B and C have coordinates $(3, -2, 4)$, $(5, 4, 0)$ and $(11, 6, -4)$ respectively.

(a) (i) Find the vector \overrightarrow{BA} . (2 marks)

(ii) Show that the size of angle ABC is $\cos^{-1}\left(-\frac{5}{7}\right)$. (5 marks)

(b) The line l has equation $\mathbf{r} = \begin{bmatrix} 8 \\ -3 \\ 2 \end{bmatrix} + \lambda \begin{bmatrix} 1 \\ 3 \\ -2 \end{bmatrix}$.

(i) Verify that C lies on l . (2 marks)

(ii) Show that AB is parallel to l . (1 mark)

(c) The quadrilateral $ABCD$ is a parallelogram. Find the coordinates of D . (3 marks)

7 The lines l_1 and l_2 have equations $\mathbf{r} = \begin{bmatrix} 8 \\ 6 \\ -9 \end{bmatrix} + \lambda \begin{bmatrix} 3 \\ -3 \\ -1 \end{bmatrix}$ and $\mathbf{r} = \begin{bmatrix} -4 \\ 0 \\ 11 \end{bmatrix} + \mu \begin{bmatrix} 1 \\ 2 \\ -3 \end{bmatrix}$ respectively.

(a) Show that l_1 and l_2 are perpendicular. (2 marks)

(b) Show that l_1 and l_2 intersect and find the coordinates of the point of intersection, P . (5 marks)

(c) The point $A(-4, 0, 11)$ lies on l_2 . The point B on l_1 is such that $AP = BP$.

Find the length of AB . (4 marks)

Core 4 Vectors Answers

7(a)(i)	$\overline{AB} = \begin{bmatrix} 6 \\ 5 \\ 3 \end{bmatrix} - \begin{bmatrix} 2 \\ 1 \\ 3 \end{bmatrix} = \begin{bmatrix} 4 \\ 4 \\ 0 \end{bmatrix}$	M1		Penalise use of co-ordinates at first occurrence only
		A1	2	
(ii)	$\begin{bmatrix} 4 \\ 4 \\ 0 \end{bmatrix} = 4 \begin{bmatrix} 1 \\ 1 \\ 0 \end{bmatrix} \Rightarrow \text{parallel}$	E1	1	Needs comment "same direction" Or "same gradient" (Or by scalar product)
(iii)	$\begin{bmatrix} 2 \\ -3 \\ -1 \end{bmatrix} = \begin{bmatrix} 6 \\ 1 \\ -1 \end{bmatrix} + \lambda \begin{bmatrix} 1 \\ 1 \\ 0 \end{bmatrix}$ <p>is satisfied by $\lambda = -4$</p>	M1		
		A1	2	$\lambda = -4$ satisfies 2 equations
(b)(i)	l_2 has equation $r = \begin{bmatrix} 4 \\ 1 \\ 1 \end{bmatrix} + \lambda \left[\begin{bmatrix} 4 \\ 1 \\ 1 \end{bmatrix} - \begin{bmatrix} 2 \\ -3 \\ -1 \end{bmatrix} \right] = \begin{bmatrix} 4 \\ 1 \\ 1 \end{bmatrix} + \lambda \begin{bmatrix} 2 \\ 4 \\ 2 \end{bmatrix}$	M1A1	2	Or $r = \begin{bmatrix} 2 \\ -3 \\ -1 \end{bmatrix} + t \begin{bmatrix} 2 \\ 4 \\ 2 \end{bmatrix}$ M1 calculate and use direction vector A1 all correct
(ii)	$\begin{bmatrix} 1 \\ 2 \\ 1 \end{bmatrix} \cdot \begin{bmatrix} 4 \\ 0 \\ -4 \end{bmatrix} = 4 - 4 = 0$ <p>$\Rightarrow 90^\circ$ (or perpendicular)</p>	M1A1		Clear attempt to use directions of AC and l_2 in scalar product
		A1F	3	Accept a correct ft value of $\cos \theta$
Total			10	

6(a)(i)	$\overline{OC} = 2 \begin{bmatrix} 3 \\ 2 \\ -1 \end{bmatrix} = \begin{bmatrix} 6 \\ 4 \\ -2 \end{bmatrix}$	B1	1	(Penalise coordinates once only)
(ii)	$\overline{AB} = \begin{bmatrix} 3 \\ 2 \\ -1 \end{bmatrix} - \begin{bmatrix} 2 \\ 4 \\ 1 \end{bmatrix} = \begin{bmatrix} 1 \\ -2 \\ -2 \end{bmatrix}$	M1 A1	2	$\overline{OA} - \overline{OB}$ or $\overline{OB} - \overline{OA}$ or 2/3 correct cpts. A0 for line AB
(b)(i)	$AC^2 = (6-2)^2 + (4-4)^2 + (-1-2)^2 = 25$ $AC = 5$	M1 A1	2	Components of AC AG
(ii)	$\overline{AB} \bullet \overline{AC} = \begin{bmatrix} 1 \\ -2 \\ -2 \end{bmatrix} \bullet \begin{bmatrix} 4 \\ 0 \\ -3 \end{bmatrix} = 4 + 6 = 10$ $3 \times 5 \times \cos \theta = 10$ $\theta = 48.189 \approx 48^\circ$	M1 A1F M1 A1	4	Clear attempt to use \overline{AB} and \overline{AC} ft \overline{AB} from a(ii) and/or \overline{AC} from b(i) Use of $ a b \cos \theta = \mathbf{a} \cdot \mathbf{b}$ with one correct and $\mathbf{a} \cdot \mathbf{b}$ evaluated CAO (AWRT)
(c)	<p>Alternative: use of cos rule Find 3rd side + use cos rule</p> $\overline{BP} = \begin{bmatrix} \alpha - 3 \\ \beta - 2 \\ \gamma - -1 \end{bmatrix}$ $\begin{bmatrix} 4 \\ 0 \\ -3 \end{bmatrix} \bullet \overline{BP} = 0$ $4\alpha - 3\gamma - 15 = 0$	(M2) (A1F) (A1) B1 M1 A1	3	ft on previously found vectors CAO (AWRT) Their \overline{BP} AG convincingly obtained
Total			12	

6(a)(i)	$\overline{BA} = \begin{bmatrix} 3 \\ -2 \\ 4 \end{bmatrix} - \begin{bmatrix} 5 \\ 4 \\ 0 \end{bmatrix} = \begin{bmatrix} -2 \\ -6 \\ 4 \end{bmatrix}$	M1A1	2	Attempt $\pm \overline{BA}$ ($OA - OB$ or $OB - OA$)
(ii)	$\overline{BC} = \begin{bmatrix} 6 \\ 2 \\ -4 \end{bmatrix}$	B1		Allow \overline{CB} ; or $\begin{bmatrix} -6 \\ -2 \\ 4 \end{bmatrix} = \overline{BC}$ or $\overline{CB} = \begin{bmatrix} 6 \\ 2 \\ -4 \end{bmatrix}$ May not see explicitly
	$ \overline{BA} = \left(\sqrt{(-2)^2 + (-6)^2 + (4)^2} \right) = \sqrt{56}$	B1F		Calculate modulus of \overline{BA} or \overline{BC} ; for finding modulus of one of vectors they have used
	$\overline{BA} \cdot \overline{BC} = \begin{bmatrix} -2 \\ -6 \\ 4 \end{bmatrix} \cdot \begin{bmatrix} 6 \\ 2 \\ -4 \end{bmatrix} = -12 - 12 - 16$	M1		Attempt at $\overline{BA} \cdot \overline{BC}$ with numerical answer; or $\overline{AB} \cdot \overline{CB}$
		A1		for -40 , or correct if done with multiples of vectors

$\cos ABC = \frac{-40}{\sqrt{56}\sqrt{56}} = -\frac{5}{7}$	A1	5	AG (convincingly obtained) Cosine rule: M1 attempt to find 3 sides A1 lengths of sides M1 cosine rule A1F correct A1 rearrange to get $\cos ABC = \frac{-5}{7}$ (ft on length of sides)
--	----	---	--

6 (cont) (b)(i)	$\begin{bmatrix} 8 \\ -3 \\ 2 \end{bmatrix} + \lambda \begin{bmatrix} 1 \\ 3 \\ -2 \end{bmatrix} = \begin{bmatrix} 11 \\ 6 \\ -4 \end{bmatrix} \quad (\lambda = 3)$	M1A1	2	$\lambda = 3$ verified in three equations M1 for $\begin{cases} 11 = 8 + \lambda \\ 6 = -3 + 3\lambda \\ -4 = 2 - 2\lambda \end{cases}$
				A1 for $\lambda = 3$ shown for all three equations
				$\lambda \begin{bmatrix} 1 \\ 3 \\ -2 \end{bmatrix} = \begin{bmatrix} 11 \\ 6 \\ -4 \end{bmatrix} - \begin{bmatrix} 8 \\ -3 \\ 2 \end{bmatrix} \therefore \lambda = 3 \quad \text{M1A1}$
				SC: $\lambda = 3$ written and nothing else: SC1
(ii)	$\begin{bmatrix} 2 \\ 6 \\ -4 \end{bmatrix} = 2 \begin{bmatrix} 1 \\ 3 \\ -2 \end{bmatrix}$ <p>\therefore same direction or same gradient or parallel</p>	E1	1	

(c)	$\overline{OD} = \overline{OC} + \overline{BA}$	B1		PI; \overline{OD} = correct vector expression which may involve \overline{AD}
	$= \begin{bmatrix} 11 \\ 6 \\ -4 \end{bmatrix} + \begin{bmatrix} -2 \\ -6 \\ 4 \end{bmatrix} = \begin{bmatrix} 9 \\ 0 \\ 0 \end{bmatrix}$ D is $(9, 0, 0)$	M1A1	3	M1 for substituting into vector expression for \overline{OD} NMS 3/3
Total			13	

7(a)	$\begin{bmatrix} 3 \\ -3 \\ -1 \end{bmatrix} \cdot \begin{bmatrix} 1 \\ 2 \\ -3 \end{bmatrix} = 3 - 6 + 3 = 0$ $= 0 \Rightarrow$ perpendicular	M1		attempt at sp, 3 terms, added
		A1	2	$= 0 \Rightarrow$ perpendicular seen (or $\cos \theta = 0 \Rightarrow \theta = 90^\circ$)
				Allow $\frac{3}{-6}$ but not $\begin{bmatrix} 3 \\ -6 \\ 3 \end{bmatrix} = 0$
(b)	$8 + 3\lambda = -4 + \mu$ $6 - 3\lambda = 2\mu$ $-9 - \lambda = 11 - 3\mu$ $\lambda = -2, \mu = 6$ verify third equation intersect at $(2, 12, -7)$ Alt (for last two marks) substitute λ into l_1 and μ into l_2	M1		set up any two equations
		m1 A1 m1		solve for λ and μ substitute λ, μ in third equation
		A1	5	CAO
		(m1)		
7(c)	intersect at $(2, 12, -7)$, condone $\begin{pmatrix} 2 \\ 12 \\ -7 \end{pmatrix}$	(A1)		$(2, 12, -7)$ found from both lines Note: working for (b) done in (a): award marks in (b)
	$\overline{AP} = \begin{pmatrix} 6 \\ 12 \\ -18 \end{pmatrix}$ $AP^2 = 504$ $AB^2 = 2AP^2$ $AB = 12\sqrt{7}$	M1		$\overline{AP} = \pm \left\{ \text{their } \overline{OP} - \begin{pmatrix} -4 \\ 0 \\ 11 \end{pmatrix} \right\}$
		A1F		fit on P
		M1		Calculate AB^2
		A1	4	OE accept 31.7 or better
Total			11	