

Edexcel Maths C4

Topic Questions from Papers

Vectors

7. The line l_1 has vector equation

$$\mathbf{r} = \begin{pmatrix} 3 \\ 1 \\ 2 \end{pmatrix} + \lambda \begin{pmatrix} 1 \\ -1 \\ 4 \end{pmatrix}$$

and the line l_2 has vector equation

$$\mathbf{r} = \begin{pmatrix} 0 \\ 4 \\ -2 \end{pmatrix} + \mu \begin{pmatrix} 1 \\ -1 \\ 0 \end{pmatrix},$$

where λ and μ are parameters.

The lines l_1 and l_2 intersect at the point B and the acute angle between l_1 and l_2 is θ .

(a) Find the coordinates of B . (4)

(b) Find the value of $\cos \theta$, giving your answer as a simplified fraction. (4)

The point A , which lies on l_1 , has position vector $\mathbf{a} = 3\mathbf{i} + \mathbf{j} + 2\mathbf{k}$.

The point C , which lies on l_2 , has position vector $\mathbf{c} = 5\mathbf{i} - \mathbf{j} - 2\mathbf{k}$.

The point D is such that $ABCD$ is a parallelogram.

(c) Show that $|\vec{AB}| = |\vec{BC}|$. (3)

(d) Find the position vector of the point D . (2)



6. The line l_1 has vector equation

$$\mathbf{r} = 8\mathbf{i} + 12\mathbf{j} + 14\mathbf{k} + \lambda(\mathbf{i} + \mathbf{j} - \mathbf{k}),$$

where λ is a parameter.

The point A has coordinates $(4, 8, a)$, where a is a constant. The point B has coordinates $(b, 13, 13)$, where b is a constant. Points A and B lie on the line l_1 .

(a) Find the values of a and b .

(3)

Given that the point O is the origin, and that the point P lies on l_1 such that OP is perpendicular to l_1 ,

(b) find the coordinates of P .

(5)

(c) Hence find the distance OP , giving your answer as a simplified surd.

(2)



5. The point A , with coordinates $(0, a, b)$ lies on the line l_1 , which has equation

$$\mathbf{r} = 6\mathbf{i} + 19\mathbf{j} - \mathbf{k} + \lambda(\mathbf{i} + 4\mathbf{j} - 2\mathbf{k}).$$

(a) Find the values of a and b .

(3)

The point P lies on l_1 and is such that OP is perpendicular to l_1 , where O is the origin.

(b) Find the position vector of point P .

(6)

Given that B has coordinates $(5, 15, 1)$,

(c) show that the points A, P and B are collinear and find the ratio $AP:PB$.

(4)



Leave
blank

Question 5 continued

Lined writing area for the answer to Question 5.



Leave
blank

Question 6 continued

Handwritten response area consisting of approximately 30 horizontal lines.



N 2 6 2 8 2 A 0 1 3 2 4

6. With respect to a fixed origin O , the lines l_1 and l_2 are given by the equations

$$l_1: \mathbf{r} = (-9\mathbf{i} + 10\mathbf{k}) + \lambda(2\mathbf{i} + \mathbf{j} - \mathbf{k})$$

$$l_2: \mathbf{r} = (3\mathbf{i} + \mathbf{j} + 17\mathbf{k}) + \mu(3\mathbf{i} - \mathbf{j} + 5\mathbf{k})$$

where λ and μ are scalar parameters.

(a) Show that l_1 and l_2 meet and find the position vector of their point of intersection. **(6)**

(b) Show that l_1 and l_2 are perpendicular to each other. **(2)**

The point A has position vector $5\mathbf{i} + 7\mathbf{j} + 3\mathbf{k}$.

(c) Show that A lies on l_1 . **(1)**

The point B is the image of A after reflection in the line l_2 .

(d) Find the position vector of B . **(3)**



Leave
blank

Question 6 continued

Lined area for writing answers, consisting of numerous horizontal lines.



4. With respect to a fixed origin O the lines l_1 and l_2 are given by the equations

$$l_1: \mathbf{r} = \begin{pmatrix} 11 \\ 2 \\ 17 \end{pmatrix} + \lambda \begin{pmatrix} -2 \\ 1 \\ -4 \end{pmatrix} \qquad l_2: \mathbf{r} = \begin{pmatrix} -5 \\ 11 \\ p \end{pmatrix} + \mu \begin{pmatrix} q \\ 2 \\ 2 \end{pmatrix}$$

where λ and μ are parameters and p and q are constants. Given that l_1 and l_2 are perpendicular,

(a) show that $q = -3$. (2)

Given further that l_1 and l_2 intersect, find

(b) the value of p , (6)

(c) the coordinates of the point of intersection. (2)

The point A lies on l_1 and has position vector $\begin{pmatrix} 9 \\ 3 \\ 13 \end{pmatrix}$. The point C lies on l_2 .

Given that a circle, with centre C , cuts the line l_1 at the points A and B ,

(d) find the position vector of B . (3)



7. Relative to a fixed origin O , the point A has position vector $(8\mathbf{i} + 13\mathbf{j} - 2\mathbf{k})$, the point B has position vector $(10\mathbf{i} + 14\mathbf{j} - 4\mathbf{k})$, and the point C has position vector $(9\mathbf{i} + 9\mathbf{j} + 6\mathbf{k})$.

The line l passes through the points A and B .

- (a) Find a vector equation for the line l . (3)

- (b) Find $|\vec{CB}|$. (2)

- (c) Find the size of the acute angle between the line segment CB and the line l , giving your answer in degrees to 1 decimal place. (3)

- (d) Find the shortest distance from the point C to the line l . (3)

The point X lies on l . Given that the vector \vec{CX} is perpendicular to l ,

- (e) find the area of the triangle CXB , giving your answer to 3 significant figures. (3)



4. The line l_1 has vector equation

$$\mathbf{r} = \begin{pmatrix} -6 \\ 4 \\ -1 \end{pmatrix} + \lambda \begin{pmatrix} 4 \\ -1 \\ 3 \end{pmatrix}$$

and the line l_2 has vector equation

$$\mathbf{r} = \begin{pmatrix} -6 \\ 4 \\ -1 \end{pmatrix} + \mu \begin{pmatrix} 3 \\ -4 \\ 1 \end{pmatrix}$$

where λ and μ are parameters.

The lines l_1 and l_2 intersect at the point A and the acute angle between l_1 and l_2 is θ .

(a) Write down the coordinates of A . (1)

(b) Find the value of $\cos \theta$. (3)

The point X lies on l_1 where $\lambda = 4$.

(c) Find the coordinates of X . (1)

(d) Find the vector \overrightarrow{AX} . (2)

(e) Hence, or otherwise, show that $|\overrightarrow{AX}| = 4\sqrt{26}$. (2)

The point Y lies on l_2 . Given that the vector \overrightarrow{YX} is perpendicular to l_1 ,

(f) find the length of AY , giving your answer to 3 significant figures. (3)



7. The line l_1 has equation $\mathbf{r} = \begin{pmatrix} 2 \\ 3 \\ -4 \end{pmatrix} + \lambda \begin{pmatrix} 1 \\ 2 \\ 1 \end{pmatrix}$, where λ is a scalar parameter.

The line l_2 has equation $\mathbf{r} = \begin{pmatrix} 0 \\ 9 \\ -3 \end{pmatrix} + \mu \begin{pmatrix} 5 \\ 0 \\ 2 \end{pmatrix}$, where μ is a scalar parameter.

Given that l_1 and l_2 meet at the point C , find

(a) the coordinates of C .

(3)

The point A is the point on l_1 where $\lambda = 0$ and the point B is the point on l_2 where $\mu = -1$.

(b) Find the size of the angle ACB . Give your answer in degrees to 2 decimal places.

(4)

(c) Hence, or otherwise, find the area of the triangle ABC .

(5)



4. Relative to a fixed origin O , the point A has position vector $\mathbf{i} - 3\mathbf{j} + 2\mathbf{k}$ and the point B has position vector $-2\mathbf{i} + 2\mathbf{j} - \mathbf{k}$. The points A and B lie on a straight line l .

(a) Find \vec{AB} . (2)

(b) Find a vector equation of l . (2)

The point C has position vector $2\mathbf{i} + p\mathbf{j} - 4\mathbf{k}$ with respect to O , where p is a constant. Given that AC is perpendicular to l , find

(c) the value of p , (4)

(d) the distance AC . (2)



6. With respect to a fixed origin O , the lines l_1 and l_2 are given by the equations

$$l_1: \mathbf{r} = \begin{pmatrix} 6 \\ -3 \\ -2 \end{pmatrix} + \lambda \begin{pmatrix} -1 \\ 2 \\ 3 \end{pmatrix}, \quad l_2: \mathbf{r} = \begin{pmatrix} -5 \\ 15 \\ 3 \end{pmatrix} + \mu \begin{pmatrix} 2 \\ -3 \\ 1 \end{pmatrix},$$

where λ and μ are scalar parameters.

(a) Show that l_1 and l_2 meet and find the position vector of their point of intersection A . (6)

(b) Find, to the nearest 0.1° , the acute angle between l_1 and l_2 . (3)

The point B has position vector $\begin{pmatrix} 5 \\ -1 \\ 1 \end{pmatrix}$.

(c) Show that B lies on l_1 . (1)

(d) Find the shortest distance from B to the line l_2 , giving your answer to 3 significant figures. (4)



7. Relative to a fixed origin O , the point A has position vector $(2\mathbf{i} - \mathbf{j} + 5\mathbf{k})$, the point B has position vector $(5\mathbf{i} + 2\mathbf{j} + 10\mathbf{k})$, and the point D has position vector $(-\mathbf{i} + \mathbf{j} + 4\mathbf{k})$.

The line l passes through the points A and B .

- (a) Find the vector \overrightarrow{AB} . (2)

- (b) Find a vector equation for the line l . (2)

- (c) Show that the size of the angle BAD is 109° , to the nearest degree. (4)

The points A , B and D , together with a point C , are the vertices of the parallelogram $ABCD$, where $\overrightarrow{AB} = \overrightarrow{DC}$.

- (d) Find the position vector of C . (2)

- (e) Find the area of the parallelogram $ABCD$, giving your answer to 3 significant figures. (3)

- (f) Find the shortest distance from the point D to the line l , giving your answer to 3 significant figures. (2)



Leave blank

8. Relative to a fixed origin O , the point A has position vector $(10\mathbf{i} + 2\mathbf{j} + 3\mathbf{k})$, and the point B has position vector $(8\mathbf{i} + 3\mathbf{j} + 4\mathbf{k})$.

The line l passes through the points A and B .

- (a) Find the vector \vec{AB} . **(2)**

- (b) Find a vector equation for the line l . **(2)**

The point C has position vector $(3\mathbf{i} + 12\mathbf{j} + 3\mathbf{k})$.

The point P lies on l . Given that the vector \vec{CP} is perpendicular to l ,

- (c) find the position vector of the point P . **(6)**



7. With respect to a fixed origin O , the lines l_1 and l_2 are given by the equations

$$l_1 : \mathbf{r} = (9\mathbf{i} + 13\mathbf{j} - 3\mathbf{k}) + \lambda(\mathbf{i} + 4\mathbf{j} - 2\mathbf{k})$$

$$l_2 : \mathbf{r} = (2\mathbf{i} - \mathbf{j} + \mathbf{k}) + \mu(2\mathbf{i} + \mathbf{j} + \mathbf{k})$$

where λ and μ are scalar parameters.

(a) Given that l_1 and l_2 meet, find the position vector of their point of intersection. **(5)**

(b) Find the acute angle between l_1 and l_2 , giving your answer in degrees to 1 decimal place. **(3)**

Given that the point A has position vector $4\mathbf{i} + 16\mathbf{j} - 3\mathbf{k}$ and that the point P lies on l_1 such that AP is perpendicular to l_1 ,

(c) find the exact coordinates of P . **(6)**



Leave blank

6. Relative to a fixed origin O , the point A has position vector $21\mathbf{i} - 17\mathbf{j} + 6\mathbf{k}$ and the point B has position vector $25\mathbf{i} - 14\mathbf{j} + 18\mathbf{k}$.

The line l has vector equation

$$\mathbf{r} = \begin{pmatrix} a \\ b \\ 10 \end{pmatrix} + \lambda \begin{pmatrix} 6 \\ c \\ -1 \end{pmatrix}$$

where a , b and c are constants and λ is a parameter.

Given that the point A lies on the line l ,

- (a) find the value of a . (3)

Given also that the vector \vec{AB} is perpendicular to l ,

- (b) find the values of b and c , (5)

- (c) find the distance AB . (2)

The image of the point B after reflection in the line l is the point B' .

- (d) Find the position vector of the point B' . (2)



8. With respect to a fixed origin O , the line l has equation

$$\mathbf{r} = \begin{pmatrix} 13 \\ 8 \\ 1 \end{pmatrix} + \lambda \begin{pmatrix} 2 \\ 2 \\ -1 \end{pmatrix}, \text{ where } \lambda \text{ is a scalar parameter.}$$

The point A lies on l and has coordinates $(3, -2, 6)$.

The point P has position vector $(-p\mathbf{i} + 2p\mathbf{k})$ relative to O , where p is a constant.

Given that vector \vec{PA} is perpendicular to l ,

(a) find the value of p . (4)

Given also that B is a point on l such that $\angle BPA = 45^\circ$,

(b) find the coordinates of the two possible positions of B . (5)



Core Mathematics C4

Candidates sitting C4 may also require those formulae listed under Core Mathematics C1, C2 and C3.

Integration (+ constant)

$f(x)$	$\int f(x) \, dx$
$\sec^2 kx$	$\frac{1}{k} \tan kx$
$\tan x$	$\ln \sec x $
$\cot x$	$\ln \sin x $
$\operatorname{cosec} x$	$-\ln \operatorname{cosec} x + \cot x , \quad \ln \tan(\frac{1}{2}x) $
$\sec x$	$\ln \sec x + \tan x , \quad \ln \tan(\frac{1}{2}x + \frac{1}{4}\pi) $

$$\int u \frac{dv}{dx} dx = uv - \int v \frac{du}{dx} dx$$

Core Mathematics C3

Candidates sitting C3 may also require those formulae listed under Core Mathematics C1 and C2.

Logarithms and exponentials

$$e^{x \ln a} = a^x$$

Trigonometric identities

$$\sin(A \pm B) = \sin A \cos B \pm \cos A \sin B$$

$$\cos(A \pm B) = \cos A \cos B \mp \sin A \sin B$$

$$\tan(A \pm B) = \frac{\tan A \pm \tan B}{1 \mp \tan A \tan B} \quad (A \pm B \neq (k + \frac{1}{2})\pi)$$

$$\sin A + \sin B = 2 \sin \frac{A+B}{2} \cos \frac{A-B}{2}$$

$$\sin A - \sin B = 2 \cos \frac{A+B}{2} \sin \frac{A-B}{2}$$

$$\cos A + \cos B = 2 \cos \frac{A+B}{2} \cos \frac{A-B}{2}$$

$$\cos A - \cos B = -2 \sin \frac{A+B}{2} \sin \frac{A-B}{2}$$

Differentiation

f(x)	f'(x)
$\tan kx$	$k \sec^2 kx$
$\sec x$	$\sec x \tan x$
$\cot x$	$-\operatorname{cosec}^2 x$
$\operatorname{cosec} x$	$-\operatorname{cosec} x \cot x$
$\frac{f(x)}{g(x)}$	$\frac{f'(x)g(x) - f(x)g'(x)}{(g(x))^2}$

Core Mathematics C2

Candidates sitting C2 may also require those formulae listed under Core Mathematics C1.

Cosine rule

$$a^2 = b^2 + c^2 - 2bc \cos A$$

Binomial series

$$(a+b)^n = a^n + \binom{n}{1} a^{n-1}b + \binom{n}{2} a^{n-2}b^2 + \dots + \binom{n}{r} a^{n-r}b^r + \dots + b^n \quad (n \in \mathbb{N})$$

$$\text{where } \binom{n}{r} = {}^n C_r = \frac{n!}{r!(n-r)!}$$

$$(1+x)^n = 1 + nx + \frac{n(n-1)}{1 \times 2} x^2 + \dots + \frac{n(n-1)\dots(n-r+1)}{1 \times 2 \times \dots \times r} x^r + \dots \quad (|x| < 1, n \in \mathbb{R})$$

Logarithms and exponentials

$$\log_a x = \frac{\log_b x}{\log_b a}$$

Geometric series

$$u_n = ar^{n-1}$$

$$S_n = \frac{a(1-r^n)}{1-r}$$

$$S_\infty = \frac{a}{1-r} \text{ for } |r| < 1$$

Numerical integration

The trapezium rule: $\int_a^b y \, dx \approx \frac{1}{2} h \{ (y_0 + y_n) + 2(y_1 + y_2 + \dots + y_{n-1}) \}$, where $h = \frac{b-a}{n}$

Core Mathematics C1

Mensuration

$$\text{Surface area of sphere} = 4\pi r^2$$

$$\text{Area of curved surface of cone} = \pi r \times \text{slant height}$$

Arithmetic series

$$u_n = a + (n - 1)d$$

$$S_n = \frac{1}{2}n(a + l) = \frac{1}{2}n[2a + (n - 1)d]$$