

$$\begin{aligned} 8 \quad \mathbf{a} &= \int_0^1 (4x + 2e^x) \, dx \\ &= [2x^2 + 2e^x]_0^1 \\ &= (2 + 2e) - (0 + 2) = 2e \end{aligned}$$

$$\begin{aligned} \mathbf{c} &= \int_{-3}^{-1} (4 - \frac{1}{x}) \, dx \\ &= [4x - \ln|x|]_{-3}^{-1} \\ &= (-4 - 0) - (-12 - \ln 3) = 8 + \ln 3 \end{aligned}$$

$$\begin{aligned} \mathbf{e} &= \int_{\frac{1}{2}}^2 (e^x + \frac{5}{x}) \, dx \\ &= [e^x + 5 \ln|x|]_{\frac{1}{2}}^2 \\ &= (e^2 + 5 \ln 2) - (e^{\frac{1}{2}} + 5 \ln \frac{1}{2}) \\ &= e^2 - e^{\frac{1}{2}} + 10 \ln 2 \end{aligned}$$

$$\begin{aligned} 9 \quad \mathbf{a} \quad 9 - \frac{7}{x} - 2x &= 0 \\ 2x^2 - 9x + 7 &= 0 \\ (2x - 7)(x - 1) &= 0 \\ x &= 1, \frac{7}{2} \\ \therefore (1, 0) \text{ and } (\frac{7}{2}, 0) \end{aligned}$$

$$\begin{aligned} \mathbf{b} &= \int_1^{\frac{7}{2}} (9 - \frac{7}{x} - 2x) \, dx \\ &= [9x - 7 \ln|x| - x^2]_1^{\frac{7}{2}} \\ &= (\frac{63}{2} - 7 \ln \frac{7}{2} - \frac{49}{4}) - (9 - 0 - 1) \\ &= 11\frac{1}{4} - 7 \ln \frac{7}{2} \end{aligned}$$

$$\begin{aligned} 11 \quad \mathbf{a} \quad x = 3 \quad \therefore y &= e^3 \\ \frac{dy}{dx} &= e^x \quad \therefore \text{grad} = e^3 \\ \therefore y - e^3 &= e^3(x - 3) \quad [y = e^3(x - 2)] \end{aligned}$$

$$\mathbf{b} \text{ at } Q, y = 0 \quad \therefore x = 2$$

$$\text{at } R, x = 0 \quad \therefore y = -2e^3$$

$$\therefore Q(2, 0), R(0, -2e^3)$$

$$\begin{aligned} \mathbf{c} \text{ area under curve, } 0 \leq x \leq 3 \\ &= \int_0^3 e^x \, dx = [e^x]_0^3 = e^3 - 1 \end{aligned}$$

area of triangle under PQ

$$= \frac{1}{2} \times 1 \times e^3 = \frac{1}{2}e^3$$

area of triangle above QR

$$= \frac{1}{2} \times 2 \times 2e^3 = 2e^3$$


shaded area

$$= (e^3 - 1) - \frac{1}{2}e^3 + 2e^3 = \frac{5}{2}e^3 - 1$$

$$\begin{aligned} \mathbf{b} &= \int_2^4 (1 + \frac{3}{x}) \, dx \\ &= [x + 3 \ln|x|]_2^4 \\ &= (4 + 3 \ln 4) - (2 + 3 \ln 2) = 2 + 3 \ln 2 \end{aligned}$$

$$\begin{aligned} \mathbf{d} &= \int_0^{\ln 2} (2 - \frac{1}{2}e^x) \, dx \\ &= [2x - \frac{1}{2}e^x]_0^{\ln 2} \\ &= (2 \ln 2 - 1) - (0 - \frac{1}{2}) = 2 \ln 2 - \frac{1}{2} \end{aligned}$$

$$\begin{aligned} \mathbf{f} &= \int_2^3 (x^2 - \frac{2}{x}) \, dx \\ &= [\frac{1}{3}x^3 - 2 \ln|x|]_2^3 \\ &= (9 - 2 \ln 3) - (\frac{8}{3} - 2 \ln 2) \\ &= \frac{19}{3} - 2 \ln \frac{3}{2} \end{aligned}$$



$$\begin{aligned} \mathbf{b} &= -\int_0^{\ln a} (e^x - a) \, dx = -[e^x - ax]_0^{\ln a} \\ &= -[(a - a \ln a) - (1 - 0)] = 1 - a + a \ln a \end{aligned}$$

$$\mathbf{c} \quad 1 - a + a \ln a = 1 + a$$

$$a \ln a = 2a, \quad \ln a = 2, \quad a = e^2$$

$$\begin{aligned} 12 \quad \mathbf{a} \quad (\frac{3}{\sqrt{x}} - 4)^2 &= 0 \\ \sqrt{x} &= \frac{3}{4} \end{aligned}$$

$$x = \frac{9}{16} \quad \therefore (\frac{9}{16}, 0)$$

$$\begin{aligned} \mathbf{b} &= \int_{\frac{9}{16}}^1 (\frac{3}{\sqrt{x}} - 4)^2 \, dx \\ &= \int_{\frac{9}{16}}^1 (9x^{-1} - 24x^{-\frac{1}{2}} + 16) \, dx \\ &= [9 \ln|x| - 48x^{\frac{1}{2}} + 16x]_{\frac{9}{16}}^1 \\ &= (0 - 48 + 16) - (9 \ln \frac{9}{16} - 36 + 9) \\ &= -5 - 9 \ln \frac{9}{16} \approx 0.178 \end{aligned}$$