

C4 VECTORS

Worksheet E

- 1 Relative to a fixed origin, the line l has vector equation

$$\mathbf{r} = \mathbf{i} - 4\mathbf{j} + p\mathbf{k} + \lambda(2\mathbf{i} + q\mathbf{j} - 3\mathbf{k}),$$

where λ is a scalar parameter.

Given that l passes through the point with position vector $(7\mathbf{i} - \mathbf{j} - \mathbf{k})$,

- a find the values of the constants p and q , (3)

- b find, in degrees, the acute angle l makes with the line with equation

$$\mathbf{r} = 3\mathbf{i} + 4\mathbf{j} - 3\mathbf{k} + \mu(-4\mathbf{i} + 5\mathbf{j} - 2\mathbf{k}). \quad (4)$$

- 2 The points A and B have position vectors $\begin{pmatrix} 1 \\ 6 \\ 4 \end{pmatrix}$ and $\begin{pmatrix} 5 \\ 0 \\ -6 \end{pmatrix}$ respectively, relative to a fixed origin.

- a Find, in vector form, an equation of the line l which passes through A and B . (2)

The line m has equation

$$\mathbf{r} = \begin{pmatrix} 5 \\ -5 \\ 3 \end{pmatrix} + t \begin{pmatrix} 1 \\ -4 \\ 2 \end{pmatrix}.$$

Given that lines l and m intersect at the point C ,

- b find the position vector of C , (5)

- c show that C is the mid-point of AB . (2)

- 3 Relative to a fixed origin, the points P and Q have position vectors $(5\mathbf{i} - 2\mathbf{j} + 2\mathbf{k})$ and $(3\mathbf{i} + \mathbf{j})$ respectively.

- a Find, in vector form, an equation of the line L_1 which passes through P and Q . (2)

The line L_2 has equation

$$\mathbf{r} = 4\mathbf{i} + 6\mathbf{j} - \mathbf{k} + \mu(5\mathbf{i} - \mathbf{j} + 3\mathbf{k}).$$

- b Show that lines L_1 and L_2 intersect and find the position vector of their point of intersection. (6)

- c Find, in degrees to 1 decimal place, the acute angle between lines L_1 and L_2 . (4)

- 4 Relative to a fixed origin, the lines l_1 and l_2 have vector equations as follows:

$$l_1: \quad \mathbf{r} = 5\mathbf{i} + \mathbf{k} + \lambda(2\mathbf{i} - \mathbf{j} + 2\mathbf{k}),$$

$$l_2: \quad \mathbf{r} = 7\mathbf{i} - 3\mathbf{j} + 7\mathbf{k} + \mu(-\mathbf{i} + \mathbf{j} - 2\mathbf{k}),$$

where λ and μ are scalar parameters.

- a Show that lines l_1 and l_2 intersect and find the position vector of their point of intersection. (6)

The points A and C lie on l_1 and the points B and D lie on l_2 .

Given that $ABCD$ is a parallelogram and that A has position vector $(9\mathbf{i} - 2\mathbf{j} + 5\mathbf{k})$,

- b find the position vector of C . (3)

Given also that the area of parallelogram $ABCD$ is 54,

- c find the distance of the point B from the line l_1 . (4)

C4 VECTORS

Worksheet E continued

- 5 Relative to a fixed origin, the points A and B have position vectors $(4\mathbf{i} + 2\mathbf{j} - 4\mathbf{k})$ and $(2\mathbf{i} - \mathbf{j} + 2\mathbf{k})$ respectively.

a Find, in vector form, an equation of the line l_1 which passes through A and B . (2)

The line l_2 passes through the point C with position vector $(4\mathbf{i} - 7\mathbf{j} - \mathbf{k})$ and is parallel to the vector $(6\mathbf{j} - 2\mathbf{k})$.

b Write down, in vector form, an equation of the line l_2 . (1)

c Show that A lies on l_2 . (2)

d Find, in degrees, the acute angle between lines l_1 and l_2 . (4)

- 6 The points A and B have position vectors $\begin{pmatrix} 5 \\ -1 \\ -10 \end{pmatrix}$ and $\begin{pmatrix} 4 \\ 1 \\ -8 \end{pmatrix}$ respectively, relative to a fixed origin O .

a Find, in vector form, an equation of the line l which passes through A and B . (2)

The line l intersects the y -axis at the point C .

b Find the coordinates of C . (2)

The point D on the line l is such that OD is perpendicular to l .

c Find the coordinates of D . (5)

d Find the area of triangle OCD , giving your answer in the form $k\sqrt{5}$. (3)

- 7 Relative to a fixed origin, the line l_1 has the equation

$$\mathbf{r} = \begin{pmatrix} 1 \\ -6 \\ -2 \end{pmatrix} + s \begin{pmatrix} 0 \\ 4 \\ -1 \end{pmatrix}.$$

a Show that the point P with coordinates $(1, 6, -5)$ lies on l_1 . (1)

The line l_2 has the equation

$$\mathbf{r} = \begin{pmatrix} 4 \\ -4 \\ -1 \end{pmatrix} + t \begin{pmatrix} 3 \\ -2 \\ 2 \end{pmatrix},$$

and intersects l_1 at the point Q .

b Find the position vector of Q . (3)

The point R lies on l_2 such that $PQ = QR$.

c Find the two possible position vectors of the point R . (5)

- 8 Relative to a fixed origin, the points A and B have position vectors $(4\mathbf{i} + 5\mathbf{j} + 6\mathbf{k})$ and $(4\mathbf{i} + 6\mathbf{j} + 2\mathbf{k})$ respectively.

a Find, in vector form, an equation of the line l_1 which passes through A and B . (2)

The line l_2 has equation

$$\mathbf{r} = \mathbf{i} + 5\mathbf{j} - 3\mathbf{k} + \mu(\mathbf{i} + \mathbf{j} - \mathbf{k}).$$

b Show that l_1 and l_2 intersect and find the position vector of their point of intersection. (4)

c Find the acute angle between lines l_1 and l_2 . (3)

d Show that the point on l_2 closest to A has position vector $(-\mathbf{i} + 3\mathbf{j} - \mathbf{k})$. (5)