

Edexcel GCE

Further Pure Mathematics FP1

Advanced/Advanced Subsidiary

Monday 10 June 2013 – Morning

Time: 1 hour 30 minutes

Withdrawn

1.
$$M = \begin{pmatrix} a & 1 \\ 1 & 2 - a \end{pmatrix}, \text{ where } a \text{ is a constant.}$$

- (a) Find $\det M$ in terms of a . (2)

A triangle T is transformed to T' by the matrix M .

Given that the area of T' is 0,

- (b) find the value of a . (3)

2.
$$f(z) = z^3 + 5z^2 + 11z + 15$$

Given that $z = 2i - 1$ is a solution of the equation $f(z) = 0$, use algebra to solve $f(z) = 0$ completely. (5)

3.
$$z_1 = \frac{1}{2}(1 + i\sqrt{3}), \quad z_2 = -\sqrt{3} + i$$

- (a) Express z_1 and z_2 in the form $r(\cos \theta + i \sin \theta)$ giving exact values of r and θ . (4)

- (b) Find $|z_1 z_2|$. (2)

- (c) Show and label z_1 and z_2 on a single Argand diagram. (2)

4. The hyperbola H has equation

$$xy = 3$$

The point $Q(1, 3)$ is on H .

- (a) Find the equation of the normal to H at Q in the form $y = ax + b$, where a and b are constants. (5)

The normal at Q intersects H again at the point R .

- (b) Find the coordinates of R . (5)

5. Prove, by induction, that $3^{2n} + 7$ is divisible by 8 for all positive integers n . (6)

6. A curve C is in the form of a parabola with equation $y^2 = 4x$.

$P(p^2, 2p)$ and $Q(q^2, 2q)$ are points on C where $p > q$.

- (a) Find an equation of the tangent to C at P . (5)

- (b) The tangent at P and the tangent at Q are perpendicular and intersect at the point $R(-1, 2)$.

- (i) Find the exact value of p and the exact value of q . (4)

- (ii) Find the area of the triangle PQR . (4)

7. (a) Use the standard results for $\sum_{r=1}^n r^2$ and $\sum_{r=1}^n r^3$ to show that

$$\sum_{r=1}^n r^2(r-1) = \frac{n(n+1)(3n+2)(n-1)}{12}$$

- for all positive integers n . (5)

- (b) Hence find the sum of the series

$$10^2 \times 9 + 11^2 \times 10 + 12^2 \times 11 + \dots + 50^2 \times 49$$
(3)

8. $f(x) = x^3 - 2x - 3$

- (a) Show that $f(x) = 0$ has a root, α , in the interval $[1, 2]$. (3)

- (b) Starting with the interval $[1, 2]$, use interval bisection twice to find an interval of width 0.25 which contains α . (3)

- (c) Using $x_0 = 1.8$ as a first approximation to α , apply the Newton-Raphson procedure once to $f(x)$ to find a second approximation to α , giving your answer to 3 significant figures. (5)

9. With reference to a fixed origin O and coordinate axes Ox and Oy , a transformation from $\mathbb{R}^2 \rightarrow \mathbb{R}^2$ is represented by the matrix A where

$$A = \begin{pmatrix} 3 & 1 \\ 1 & -2 \end{pmatrix}$$

- (a) Find A^2 . (2)
- (b) Show that the matrix A is non-singular. (2)
- (c) Find A^{-1} . (2)

The transformation represented by matrix A maps the point P onto the point Q .

Given that Q has coordinates $(k - 1, 2 - k)$, where k is a constant,

- (d) show that P lies on the line with equation $y = 4x - 1$ (3)