

Exercise 2A

1 a $3x^2 + 7x - 4 = 0$
 $a = 3, b = 7, c = -4$
 $\alpha + \beta = -\frac{b}{a} = -\frac{7}{3}$

b $\alpha\beta = \frac{c}{a} = -\frac{4}{3}$

c $\frac{1}{\alpha} + \frac{1}{\beta} = \frac{\alpha + \beta}{\alpha\beta}$
 $= \left(\frac{-\frac{7}{3}}{-\frac{4}{3}} \right)$
 $= \frac{7}{4}$

d $\alpha^2 + \beta^2 = (\alpha + \beta)^2 - 2\alpha\beta$
 $= \left(-\frac{7}{3}\right)^2 - 2\left(-\frac{4}{3}\right)$
 $= \frac{49}{9} + \frac{8}{3}$
 $= \frac{73}{9}$

2 a $7x^2 - 3x + 1 = 0$
 $a = 7, b = -3, c = 1$
 $\alpha + \beta = -\frac{b}{a} = \frac{3}{7}$

b $\alpha\beta = \frac{c}{a} = \frac{1}{7}$

c $\frac{1}{\alpha} + \frac{1}{\beta} = \frac{\alpha + \beta}{\alpha\beta}$
 $= \frac{\frac{3}{7}}{\frac{1}{7}}$
 $= 3$

d $\alpha^2 + \beta^2 = (\alpha + \beta)^2 - 2\alpha\beta$
 $= \left(\frac{3}{7}\right)^2 - 2\left(\frac{1}{7}\right)$
 $= \frac{9}{49} - \frac{2}{7}$
 $= \frac{9}{49} - \frac{14}{49}$
 $= -\frac{5}{49}$

3 a $6x^2 - 9x + 2 = 0$
 $a = 6, b = -9, c = 2$
 $\alpha + \beta = -\frac{b}{a} = -\frac{-9}{6} = \frac{3}{2}$

b $\alpha^2 \times \beta^2 = (\alpha\beta)^2 = \left(\frac{c}{a}\right)^2 = \left(\frac{2}{6}\right)^2 = \frac{1}{9}$

c $\frac{1}{\alpha} + \frac{1}{\beta} = \frac{\alpha + \beta}{\alpha\beta} = \frac{\frac{3}{2}}{\frac{1}{3}} = \frac{9}{2}$

d $(\alpha + \beta)^3 = \alpha^3 + 3\alpha^2\beta + 3\alpha\beta^2 + \beta^3$
So $\alpha^3 + \beta^3 = (\alpha + \beta)^3 - 3\alpha\beta(\alpha + \beta)$
 $= \left(\frac{3}{2}\right)^3 - 3\left(\frac{1}{3}\right)\left(\frac{3}{2}\right)$
 $= \frac{27}{8} - \frac{3}{2}$
 $= \frac{27}{8} - \frac{12}{8}$
 $= \frac{15}{8}$

4 $ax^2 + bx + c = 0$
 $\alpha = 2 \text{ and } \beta = -3$
 $\alpha + \beta = 2 + (-3) = -1 = -\frac{b}{a}$
 $\alpha\beta = (2)(-3) = -6 = \frac{c}{a}$
 $ax^2 + bx + c = 0 \Rightarrow x^2 + \frac{b}{a}x + \frac{c}{a} = 0$
 $-\frac{b}{a} = -1 \Rightarrow \frac{b}{a} = 1$

$\frac{c}{a} = -6$
So equation is $x^2 + x - 6 = 0$,
with $a = 1, b = 1$ and $c = -6$

5 $ax^2 + bx + c = 0$
 $\alpha = -\frac{1}{2} \text{ and } \beta = -\frac{1}{3}$
 $\alpha + \beta = -\frac{1}{2} - \frac{1}{3} = -\frac{5}{6} = -\frac{b}{a}$
 $\alpha\beta = \left(-\frac{1}{2}\right)\left(-\frac{1}{3}\right) = \frac{1}{6} = \frac{c}{a}$
 $ax^2 + bx + c = 0 \Rightarrow x^2 + \frac{b}{a}x + \frac{c}{a} = 0$
 $-\frac{b}{a} = -\frac{5}{6}, \text{ so } \frac{b}{a} = \frac{5}{6}$
 $\frac{c}{a} = \frac{1}{6}$

Further Pure Maths 1**Solution Bank**

- 5** So equation is $x^2 + \frac{5}{6}x + \frac{1}{6} = 0$
 or $6x^2 + 5x + 1 = 0$. So
 $a = 6, b = 5$ and $c = 1$.

6 $ax^2 + bx + c = 0$
 $\alpha = \frac{-1+i}{2}$ and $\beta = \frac{-1-i}{2}$
 $\alpha + \beta = -1 = -\frac{b}{a}$
 $\alpha\beta = \left(\frac{-1+i}{2}\right)\left(\frac{-1-i}{2}\right) = \frac{1-i^2}{2} = \frac{2}{4} = \frac{1}{2} = \frac{c}{a}$
 $ax^2 + bx + c = 0 \Rightarrow x^2 + \frac{b}{a}x + \frac{c}{a} = 0$
 $-\frac{b}{a} = -1 \Rightarrow \frac{b}{a} = 1$
 $\frac{c}{a} = \frac{1}{2}$

So equation is $x^2 + x + \frac{1}{2} = 0$
 or $2x^2 + 2x + 1 = 0$. So
 $a = 2, b = 2$ and $c = 1$.

- 7 a** The other root, β , is the complex conjugate. So $\beta = -1 + 4i$

b $\alpha = -1 - 4i$ and $\beta = -1 + 4i$
 $\alpha + \beta = -2 = -\frac{b}{a}$
 $\alpha\beta = (-1 - 4i)(-1 + 4i) = 1 - 16i^2 = 17 = \frac{c}{a}$
 $ax^2 + bx + c = 0 \Rightarrow x^2 + \frac{b}{a}x + \frac{c}{a} = 0$
 $-\frac{b}{a} = -2 \Rightarrow \frac{b}{a} = 2$

$$\frac{c}{a} = 17$$

So equation is $x^2 + 2x + 17 = 0$,
 with $a = 1, b = 2$ and $c = 17$.

- 8** The sum of the roots is 4,
 so $\alpha + \beta = 4$

Also $-\frac{b}{a} = 4$, but $-\frac{b}{a} = \frac{-(k-3)}{k} = \frac{3-k}{k}$

So $\frac{3-k}{k} = 4$

$3-k = 4k$

$3 = 5k$

$k = \frac{3}{5}$

- 9** The roots are α and $-\alpha$.

$$\alpha + -\alpha = 0 = -\frac{b}{a}$$

If $-\frac{b}{a} = 0$, then $b = 0$

$$b = 0 \Rightarrow 16 + n = 0 \Rightarrow n = -16$$

- 10** The roots are reciprocals, so the roots are α and $\frac{1}{\alpha}$.

$$\alpha\left(\frac{1}{\alpha}\right) = 1 = \frac{c}{a}$$

But $\frac{c}{a} = \frac{k}{6}$

So $\frac{k}{6} = 1 \Rightarrow k = 6$

- 11** The roots are k and $2k$.

$$k + 2k = 3k = -\frac{b}{a}$$

But $-\frac{b}{a} = -\frac{4}{m}$, so $3k = -\frac{4}{m}$

$$(k)(2k) = 2k^2 = \frac{c}{a}$$

But $\frac{c}{a} = \frac{4m}{m} = 4$, so $2k^2 = 4$

If $2k^2 = 4, k^2 = 2$ and $k = \pm\sqrt{2}$.

$3k = -\frac{4}{m}$, so $m = -\frac{4}{3k}$

If $k = \sqrt{2}, m = -\frac{4}{3\sqrt{2}} = -\frac{2\sqrt{2}}{3}$

If $k = -\sqrt{2}, m = \frac{4}{3\sqrt{2}} = \frac{2\sqrt{2}}{3}$