



**General Certificate of Education (A-level)
June 2013**

Mathematics

MFP2

(Specification 6360)

Further Pure 2

Final

Mark Scheme

Mark schemes are prepared by the Principal Examiner and considered, together with the relevant questions, by a panel of subject teachers. This mark scheme includes any amendments made at the standardisation events which all examiners participate in and is the scheme which was used by them in this examination. The standardisation process ensures that the mark scheme covers the students' responses to questions and that every examiner understands and applies it in the same correct way. As preparation for standardisation each examiner analyses a number of students' scripts: alternative answers not already covered by the mark scheme are discussed and legislated for. If, after the standardisation process, examiners encounter unusual answers which have not been raised they are required to refer these to the Principal Examiner.

It must be stressed that a mark scheme is a working document, in many cases further developed and expanded on the basis of students' reactions to a particular paper. Assumptions about future mark schemes on the basis of one year's document should be avoided; whilst the guiding principles of assessment remain constant, details will change, depending on the content of a particular examination paper.

Further copies of this Mark Scheme are available from: aqa.org.uk

Copyright © 2013 AQA and its licensors. All rights reserved.

Copyright

AQA retains the copyright on all its publications. However, registered schools/colleges for AQA are permitted to copy material from this booklet for their own internal use, with the following important exception: AQA cannot give permission to schools/colleges to photocopy any material that is acknowledged to a third party even for internal use within the centre.

Set and published by the Assessment and Qualifications Alliance.

Key to mark scheme abbreviations

M	mark is for method
m or dM	mark is dependent on one or more M marks and is for method
A	mark is dependent on M or m marks and is for accuracy
B	mark is independent of M or m marks and is for method and accuracy
E	mark is for explanation
✓ or ft or F	follow through from previous incorrect result
CAO	correct answer only
CSO	correct solution only
AWFW	anything which falls within
AWRT	anything which rounds to
ACF	any correct form
AG	answer given
SC	special case
OE	or equivalent
A2,1	2 or 1 (or 0) accuracy marks
-x EE	deduct x marks for each error
NMS	no method shown
PI	possibly implied
SCA	substantially correct approach
c	candidate
sf	significant figure(s)
dp	decimal place(s)

No Method Shown


Where the question specifically requires a particular method to be used, we must usually see evidence of use of this method for any marks to be awarded.



Where the answer can be reasonably obtained without showing working and it is very unlikely that the correct answer can be obtained by using an incorrect method, we must award **full marks**. However, the obvious penalty to candidates showing no working is that incorrect answers, however close, earn **no marks**.

Where a question asks the candidate to state or write down a result, no method need be shown for full marks.

Where the permitted calculator has functions which reasonably allow the solution of the question directly, the correct answer without working earns **full marks**, unless it is given to less than the degree of accuracy accepted in the mark scheme, when it gains **no marks**.

Otherwise we require evidence of a correct method for any marks to be awarded.

Q	Solution	Marks	Total	Comments
1(a)	 <p data-bbox="240 517 735 651">Circle Centre at $6i$ Radius 3 & cutting positive Im axis twice</p>	M1 A1 A1	3	freehand circle 6 marked on Im axis as centre radius of 3 clearly indicated with circle in position shown
(b)(i)	(Max $ z $ is) 9	B1	1	
(ii)	Tangent from O to circle	M1		FT their circle position
	Angle of $\frac{\pi}{6}$ or $\frac{\pi}{3}$ <i>correctly</i> marked	A1		PI ; condone degrees for first A1
	(Max $\arg z$ is) $\frac{2\pi}{3}$	A1cso	3	exactly this
	Total		7	

Q	Solution	Marks	Total	Comments
<p>2(a)(i)</p> <p>sinh x graph</p>  <p>cosh x graph</p>  <p>Gradient of $\sinh x > 0$ at origin and cosh x minimum at $(0,1)$</p>	<p>M1</p> <p>M1</p> <p>A1</p>	<p>3</p>	<p>shape - curve through O, in 1st and 3rd quadrants</p> <p>shape - curve all above x-axis</p>	
<p>(ii)</p>	<p>cosh $x = 0$ has no solutions</p> <p>and $\sinh x = -k$ has one solution (hence equation has exactly one solution)</p>	<p>E1</p>	<p>1</p>	<p>or cosh $x > 0$ etc (since $y = -k$ cuts $y = \sinh x$ exactly once)</p>
<p>(b)</p>	<p>$\frac{dy}{dx} = 6\cosh x + 2\cosh x \sinh x$</p> <p>(2)cosh $x(3 + \sinh x) = 0$</p> <p>therefore C has only one stationary point</p> <p>$\Rightarrow \sinh x = -3$</p> <p>$\cosh^2 x = 10$</p> <p>$y (= -18 + 10) = -8$</p>	<p>M1</p> <p>A1</p> <p>E1\checkmark</p> <p>m1</p> <p>A1</p>	<p>5</p>	<p>one term correct all correct - may have $6\cosh x + \sinh 2x$</p> <p>putting = 0, factorising and concluding statement (may be later)</p> <p>finding $\sinh x$ from "their" equation</p> <p>answer must be integer so do not accept calculator approximation rounded to -8</p>
	Total		9	

Q	Solution	Marks	Total	Comments
3	$n = 1, \frac{3+1}{3-1} = \frac{4}{2} = 2$ <p>$(u_1 = 2$ so formula is) true when $n = 1$</p> <p>Assume formula is true for $n = k$ (*)</p> $(u_{k+1} =) \frac{5 \frac{3k+1}{3k-1} - 3}{3 \frac{3k+1}{3k-1} - 1}$ $(u_{k+1} =) \frac{5(3k+1) - 3(3k-1)}{3(3k+1) - (3k-1)}$ $u_{k+1} = \frac{3k+4}{3k+2} \text{ or } u_{k+1} = \frac{3(k+1)+1}{3(k+1)-1}$ <p>Hence formula is true for $n = k+1$ (**)</p> <p>must have lines (*) & (**) and “Result true for $n = 1$ therefore true for $n = 2, n = 3$ etc by induction.”</p>	<p>B1</p> <p>M1</p> <p>m1</p> <p>A1</p> <p>A1cso</p> <p>E1</p>	6	<p>be convinced they have used $u_n = \frac{3n+1}{3n-1}$</p> <p>clear attempt at RHS of this formula</p> <p>clear attempt to remove “double fraction”</p> $\frac{6k+8}{6k+4}$ <p>must have “$u_{k+1} =$ “ on at least this line</p> <p>must also have earned previous 5 marks before E1 is scored</p>
Total			6	
4(a)	$f(r) - f(r-1) =$ $r^2(2r^2 - 1) - (r-1)^2(2(r-1)^2 - 1)$ $= 2r^4 - r^2 - (r^2 - 2r + 1)(2r^2 - 4r + 1)$ $= 2r^4 - r^2 - (2r^4 - 8r^3 + 11r^2 - 6r + 1)$ $= 8r^3 - 12r^2 + 6r - 1$ $= (2r - 1)^3$	<p>M1</p> <p>A1</p> <p>A1cso</p>	3	<p>condone one slip here</p> <p>attempt to multiply out “their” $f(r-1)$</p> <p>$f(r)$ & $f(r-1)$ expanded correctly</p> <p>condone correct unsimplified</p> <p>AG</p>
(b)	<p>Attempt to use method of differences</p> $f(2n) - f(n)$ $f(2n) - f(n) = 4n^2(8n^2 - 1) - n^2(2n^2 - 1)$ $= 30n^4 - 3n^2$ $= 3n^2(10n^2 - 1)$	<p>M1</p> <p>m1</p> <p>A1</p> <p>A1cso</p>	4	$(2n)^2 \{2(2n^2) - 1\} - n^2(2n^2 - 1)$ <p>AG be convinced</p>
Total			7	

Q	Solution	Marks	Total	Comments
5(a)(i)	$(\alpha\beta\gamma =) -37 + 36i$	B1	1	
(ii)	$(\beta\gamma =) (-2 + 3i)(1 + 2i) = -2 + 3i - 4i - 6$ $(-8 - i) \alpha = -37 + 36i$ $\Rightarrow (8 + i) \alpha = 37 - 36i$	M1 A1cso	2	correct unsimplified but must simplify i^2 AG be convinced
(iii)	$\Rightarrow \alpha = \frac{37 - 36i}{8 + i} \times \frac{8 - i}{8 - i}$ $= \frac{296 - 37i - 288i - 36}{65}$ $= \frac{260 - 325i}{65}$ $= 4 - 5i$	M1 A1 A1cao	3	correct unsimplified Alternative $(8 + i)(m + ni) = 37 - 36i$ $8m - n = 37; m + 8n = -36$ M1 <i>either</i> $m = 4$ <i>or</i> $n = -5$ A1 $\alpha = 4 - 5i$ A1
(b)	$\alpha + \beta + \gamma = -p$ $-2 + 3i + 1 + 2i + 4 - 5i = 3$ $(\Rightarrow p =) -3$	B1	1	
(c)	$\alpha\beta + \beta\gamma + \gamma\alpha = q$ $(7 + 22i) + (-8 - i) + (14 + 3i) = q$ $q = 13 + 24i$	M1 A1cao	2	$q = \sum \alpha\beta$ and attempt to evaluate three products FT "their" α
Total			9	

Q	Solution	Marks	Total	Comments
6(a)	$(5 \cosh x - 3 \sinh x)$ $= \frac{5}{2}(e^x + e^{-x}) - \frac{3}{2}(e^x - e^{-x})$ $= e^x + 4e^{-x}$ $\frac{1}{5 \cosh x - 3 \sinh x} = \frac{e^x}{4 + e^{2x}}$	<p>M1</p> <p>A1</p> <p>A1cso</p>	<p>3</p>	<p>cosh x and sinh x correct in terms of e^x</p> <p>may be seen as denominator</p> <p>** must have left hand-side ; $m = 4$</p>
(b)	$u = e^x \Rightarrow du = e^x dx$ $\Rightarrow \int \frac{1}{4+u^2} (du)$ $= \frac{1}{2} \tan^{-1} \frac{u}{2}$ $x = 0 \Rightarrow u = 1 \quad x = \ln 2 \Rightarrow u = 2$ $\frac{1}{2} \tan^{-1} 1 - \frac{1}{2} \tan^{-1} \frac{1}{2}$ $= \frac{\pi}{8} - \frac{1}{2} \tan^{-1} \frac{1}{2}$	<p>M1</p> <p>A1✓</p> <p>A1✓</p> <p>A1✓</p> <p>A1cso</p>	<p>5</p>	<p>or $\frac{du}{dx} = e^x$</p> <p>FT “their” m from part(a) $\Rightarrow \int \frac{1}{m+u^2} du$</p> <p>FT “their” $\frac{1}{\sqrt{m}} \tan^{-1} \frac{u}{\sqrt{m}}$</p> <p>FT “their” $\frac{1}{\sqrt{m}} \left(\tan^{-1} \frac{2}{\sqrt{m}} - \tan^{-1} \frac{1}{\sqrt{m}} \right)$</p> <p>AG</p>
	Total		8	

Q	Solution	Marks	Total	Comments
7(a)(i)	$\frac{d}{du}(2u\sqrt{1+4u^2}) = \frac{8u^2}{\sqrt{1+4u^2}} + 2\sqrt{1+4u^2}$	M1	4	M1 for clear use of product rule (condone one error in one term) correct unsimplified be convinced – must see this line OE
	$\frac{d}{du}(\sinh^{-1} 2u) = \frac{2}{\sqrt{1+4u^2}}$	A1		
	$\frac{8u^2 + 2}{\sqrt{1+4u^2}} = \frac{2(1+4u^2)}{\sqrt{1+4u^2}} = 2\sqrt{1+4u^2}$	B1		
	$\frac{d}{du}(2u\sqrt{1+4u^2} + 4\sinh^{-1} 2u) = 4\sqrt{1+4u^2}$	A1cso		
(ii)	$\frac{1}{\text{“their” } k} [2u\sqrt{1+4u^2} + \sinh^{-1} 2u]_0^1$	M1	2	anti differentiation FT “their” k or even use of k
	$= \frac{\sqrt{5}}{2} + \frac{1}{4} \sinh^{-1} 2$	A1✓		
(b)(i)	$y = \frac{1}{2} \cos 4x$ and $\frac{dy}{dx} = A \sin 4x$			$\frac{dy}{dx} = -2 \sin 4x$
	substituted into $\int K y \left(1 + \left(\frac{dy}{dx}\right)^2\right) (dx)$	M1		clear attempt to use formula for CSA
	$(S =) \int_0^{\frac{\pi}{8}} 2\pi \times \frac{1}{2} \cos 4x \sqrt{1+4 \sin^2 4x} dx$ = printed answer (combining $2 \times \frac{1}{2}$)	A1cso	2	AG $\frac{dy}{dx} = -2 \sin 4x$ and $2 \times \frac{1}{2}$ and dx must be seen to award A1cso
	(ii) $u = \sin 4x \Rightarrow du = 4 \cos 4x dx$	M1		condone $du = B \cos 4x dx$ for M1
	$(S =) \frac{\pi}{4} \int_0^1 \sqrt{1+4u^2} (du)$	A1		condone limits seen later
		m1		use of their result from (a)(ii) correctly FT “their” B
	$(S =) \frac{\pi\sqrt{5}}{8} + \frac{\pi}{16} \sinh^{-1} 2$	A1cso	4	OE
Total			12	

Q	Solution	Marks	Total	Comments	
8(a)(i)	$\cos 4\theta + i \sin 4\theta = (\cos \theta + i \sin \theta)^4$ $\cos^4 \theta + 4i \cos^3 \theta \sin \theta + 6i^2 \cos^2 \theta \sin^2 \theta + 4i^3 \cos \theta \sin^3 \theta + i^4 \sin^4 \theta$	M1	5	De Moivre & attempt to expand RHS	
	Equating "their" real parts $\cos 4\theta = \cos^4 \theta - 6 \cos^2 \theta \sin^2 \theta + \sin^4 \theta$ $\sin 4\theta = 4 \cos^3 \theta \sin \theta - 4 \cos \theta \sin^3 \theta$	A1 m1 A1 B1		any correct expansion or imaginary parts AG be convinced correct	
(ii)	$\tan 4\theta = \frac{\text{"their expression for " } \sin 4\theta}{\text{"their expression for " } \cos 4\theta}$	M1		3	AG be convinced
	Division by $\cos^4 \theta$ $\tan 4\theta = \frac{4 \tan \theta - 4 \tan^3 \theta}{1 - 6 \tan^2 \theta + \tan^4 \theta}$	m1 A1			
(b)	$(\tan 4\theta = 1 \Rightarrow) \quad 1 = \frac{4t - 4t^3}{1 - 6t^2 + t^4}$	M1		4	when $\theta = \frac{\pi}{16}$
	$1 - 6t^2 + t^4 = 4t - 4t^3$	A1	AG be convinced		
	$\Rightarrow t^4 + 4t^3 - 6t^2 - 4t + 1 = 0$ $\theta = \frac{\pi}{16} \text{ satisfies } \tan 4\theta = 1$	E1	both statements required		
	$\Rightarrow \tan \frac{\pi}{16} \text{ is root of quartic equation}$		or equivalent tan expressions		
(c)	$\sum \alpha = -4 \quad \text{and} \quad \sum \alpha\beta = -6$	B1	5	watch for minus signs	
	$\sum \alpha^2 = (\sum \alpha)^2 - 2 \sum \alpha\beta$ $= (16 + 12) = 28$	M1 A1cso		correct formula	
	$\tan \frac{9\pi}{16} = -\tan \frac{7\pi}{16}, \quad \tan \frac{13\pi}{16} = -\tan \frac{3\pi}{16}$ $\tan^2 \frac{\pi}{16} + \tan^2 \frac{3\pi}{16} + \tan^2 \frac{5\pi}{16} + \tan^2 \frac{7\pi}{16} = 28$	B1 A1cso		explicitly seen AG must earn previous 4 marks	
	Total		17		
	TOTAL		75		