


1. (a)

Shape, vertex on x -axis

B1

At least $2a$ seen on positive x -axis

B1 2

- (b) Attempting to solve $-(x - 2a) = 2x + a$ anywhere
 Completely correct method
 [e.g. solving $-(x - 2a) > 2x + a$;
 if finding two “solutions” needs to be evidence for giving
 “correct” result]
 $x < 1/3 a$

M1
dep M1A1 3
[5]

2. I.F. = $e^{\int 2 \cot 2x dx}$; = $\sin 2x$

M1 A1

Multiplying **throughout** by IF.

M1(*)

 $Y \times (\text{IF}) = \text{integral of candidate's RHS}$

M1

$$= \int 2 \sin^2 x \cos x \, dx \text{ or } \int -\left(\frac{\cos 3x - \cos x}{2}\right) dx$$

M1

[This M gained when in position to complete integration, dep on M(*)]

$$= \frac{2}{3} \sin^3 x (+C) \text{ or } -\frac{1}{6} \sin 3x + \frac{1}{2} \sin x + c$$


A1

$$y = \frac{2 \sin^3 x}{3 \sin 2x} + \frac{C}{\sin 2x} \text{ or } -\frac{\sin 3x}{6 \sin 2x} + \frac{\sin x}{2 \sin 2x} + \frac{c}{\sin 2x} \text{ or equiv.}$$

A1ft

[7]

3. (a) $\frac{dy}{dx} = x \frac{dv}{dx} + v$, $\frac{d^2y}{dx^2} = x \frac{d^2v}{dx^2} + 2 \frac{dv}{dx}$ M1A1
- [M1 for diff. product, A1 both correct]*
- $\therefore x^2 \left(x \frac{d^2v}{dx^2} + 2 \frac{dv}{dx} \right) - 2x \left(x \frac{dv}{dx} + v \right) + (2 + 9x^2)vx = x^5$ M1
- $x^3 \frac{d^2v}{dx^2} + 2x^2 \frac{dv}{dx} - 2x^2 \frac{dv}{dx} - 2vx + 2vx + 9vx^3 = x^5$ A1
- $[x^3 \frac{d^2v}{dx^2} + 9vx^3 = x^5]$
- Given result: $\frac{d^2v}{dx^2} + 9v = x^2$ A1 5
- cs0*
- (b) CF: $v = A \sin 3x + B \cos 3x$ (may just write it down) M1A1
 Appropriate form for P1: $v = \lambda x^2 + \mu$ (or $ax^2 + bx + c$) M1
 Complete method to find λ and μ (or a, b, c) M1
- $v = A \sin 3x + B \cos 3x + \frac{1}{9}x^2 - \frac{2}{81}$ M1A1ft6
- [f.t. only on wrong CF]*
- (c) $\therefore y = Ax \sin 3x + Bx \cos 3x + \frac{1}{9}x^3 - \frac{2}{81}x$ B1ft 1
- [f.t. for $y = x$ (candidate's CF + PI), providing two arbitrary constants]*
- [12]**
4. (a) For C: Using polar/ Cartesian relationships to form Cartesian equation M1
 so $x^2 + y^2 = 6x$ A1
 [Equation in any form: e.g. $(x - 3)^2 + y^2 = 9$ from sketch.
 or $\sqrt{x^2 + y^2} = \frac{6x}{\sqrt{x^2 + y^2}}$]
- For D: $r \cos \left(\frac{\pi}{3} - \theta \right) = 3$ and attempt to expand M1
- $\frac{x}{2} + \frac{\sqrt{3}y}{2} = 3$ (any form) M1A1 5



“Circle”, symmetric in initial line passing through pole
 Straight line
 Both passing through (6, 0)

B1
 B1
 B1 3

(c) Polars: Meet where $6\cos\theta\cos(\frac{\pi}{3}-\theta) = 3$

M1

$$\sqrt{3}\sin\theta\cos\theta = \sin^2\theta$$

M1

$$\sin\theta = 0 \text{ or } \tan\theta = \sqrt{3} \quad [\theta = 0 \text{ or } \frac{\pi}{3}]$$

M1

Points are (6, 0) and $(3, \frac{\pi}{3})$

B1, A1 5

[13]

Alternatives (only more common):

(a) Equation of D :

Finding two points on line

M1

Using correctly in Cartesian equation for straight line

M1

Correct Cartesian equation

A1

(c) Cartesian: Eliminate x or y to form quadratic in one variable

M1

$$[2x^2 - 15x + 18 = 0, 4y^2 - 6\sqrt{3}y = 0]$$

Solve to find values of x or y

M1

Substitute to find values of other variable

$$\left[x = \frac{3}{2} \text{ or } 6; \quad y = 0 \text{ or } \frac{3\sqrt{3}}{2} \right]$$

B1A1

Points must be (6, 0) and $(3, \frac{\pi}{3})$

B1A1

5. $\frac{dy}{dx} + \frac{2}{1+x}y = \frac{1}{x(x+1)}$ M1

Attempt $y' = Py = Q$ form


I.F. = $e^{\int \frac{2}{1+x} dx} = e^{2\ln(1+x)} = (1+x)^2$ M1, A1

$\therefore y(1+x)^2 = \int \left(\frac{x+1}{x}\right) dx$ OR $\frac{d}{dx}(y(1+x)^2) = \frac{x+1}{x}$ M1 (ft I.F.)

i.e. $(y(1+x)^2)' = x + \ln x + (C)$ M1 A1

$y = \frac{x + \ln x + C}{(1+x)^2}$ A1 c.a.o.

[7]



shape – Symmetric about y-axis B1

shape – Vertex on positive x-axis B1

$-z, z$ B1

$\frac{1}{2}$ B1 4

(b) $x^2 - 4 = 2x - 1$ M1

$x^2 - 2x - 3 = 0 \Rightarrow x = 3, -1$ A1

$x^2 - 4 = -(2x - 1)$ M1

$x^2 + 2x - 5 = 0, \Rightarrow x = \frac{-2 \pm \sqrt{4 + 20}}{2}$ A1,

$x = -1 \pm \sqrt{6}$ A1 5

correct 3 term quadratic = 0


(c) $x < -1 - \sqrt{6}; -1 < x < \sqrt{6} - 1, x > 3$ ($\sqrt{\text{surds}}$) B1ft; B1ft; B1

Accept 3sf.

[12]

7. (a) $2m^2 + 5m + 2 = 0$ M1
- Attempt aux eqn $\rightarrow m =$*
- $\Rightarrow m = -\frac{1}{2}, -2$
- $\therefore x_{CF} = Ae^{-2t} + Be^{-\frac{1}{2}t}$ A1
- C.F.*
- Particular Integral: $x = pt + q$ B1
- P.I.*
- $\dot{x} = p, \ddot{x} = 0$ and sub. M1
- $\Rightarrow 5p + 2q + 2pt = 2t + q \rightarrow p = 1, q = 2$ A1
- General solution $x = \underline{Ae^{-2t} + Be^{-\frac{1}{2}t} + t + 2}$ A1ft (ft ms, p.q)
-
- (b) $x = 3, t = 0 \Rightarrow 3 = A + B + 2$ (or $A + B = 1$) M1
- $\dot{x} = -2Ae^{-2t} - \frac{1}{2}Be^{-\frac{1}{2}t} + 1$ M1
- Attempt \dot{x}*
- $\dot{x} = -1, t = 0 \Rightarrow -1 = -2A - \frac{1}{2}B + 1$ (or $4A + B = 4$) A1
- 2 correct eqns*
- Solving $\rightarrow A = 1, B = 0$ and $x = \underline{e^{-2t} + t + 2}$ A1 4
-
- (c) $\dot{x} = -2e^{-2t} + 1 = 0$ M1
- $\dot{x} = 0$
- $\Rightarrow t = \frac{1}{2} \ln 2$ A1
- $\ddot{x} = 4e^{-2t} > 0 (\forall t) \therefore \text{min}$ M1
- Min $x = e^{-\ln 2} + \frac{1}{2} \ln 2 + 2$
- $= \frac{1}{2} + \frac{1}{2} \ln 2 + 2$
- $\equiv \frac{1}{2} (5 + \ln 2) (*)$ A1 c.s.o.
- [14]**

8. (a)



$$4a(1 + \cos \theta) = \frac{3a}{\cos \theta} \quad \text{or} \quad r = 4a \left(1 + \frac{3a}{r} \right) \quad \text{M1}$$

$$4\cos^2 \theta + 4\cos \theta - 3 = 0 \quad \text{or} \quad r^2 - 4ar - 12a^2 = 0 \quad \text{A1}$$

$$(2\cos \theta - 1)(2\cos \theta + 3) = 0 \quad \text{or} \quad (r - 6a)(r + 2a) = 0 \quad \text{M1}$$

$$\cos \theta = \frac{1}{2}, \quad \left(\theta = \frac{\pi}{3} \right) \quad \text{or} \quad r = 6a \quad \text{A1}$$

Note $ON = 3a$

$$PQ = 2 \times ON \tan \frac{\pi}{6} = 6\sqrt{3}a (*) \quad \text{cso M1 A1}$$

$$\text{or } PQ = 2 \times \sqrt{[(6a)^2 - (3a)^2]} = 2\sqrt{(27a^2)} = 6\sqrt{3}a (*) \quad \text{cso}$$

or any complete equivalent

$$(b) \quad 2 \times \frac{1}{2} \int_0^{\pi/3} r^2 d\theta = \dots \int \dots 16a^2 (1 + \cos \theta)^2 d\theta \quad \text{M1}$$

$$\int r^2 d\theta$$

$$= \dots \int \dots \left(1 + 2\cos \theta + \frac{1}{2} + \frac{1}{2} \cos 2\theta \right) d\theta \quad \text{M1}$$

$$\cos^2 \theta \rightarrow \cos 2\theta$$

$$= \dots \left[\frac{3}{2}\theta + 2\sin \theta + \frac{1}{4}\sin 2\theta \right] \quad \text{A1}$$

$$= 16a^2 \left[\frac{\pi}{2} + \sqrt{3} + \frac{\sqrt{3}}{8} \right] (= 2a^2[4\pi + 9\sqrt{3}] \approx 56.3a^2) \quad \text{M1 A1}$$


use of their $\frac{\pi}{3}$ for M1

$$\text{Area of } \triangle POQ = \frac{1}{2} 6\sqrt{3} a \times 3a \text{ or } 9a^2 \sqrt{3} \quad \text{B1}$$

$$R = a^2(8\pi + 9\sqrt{3}) \quad \text{cao A1 7}$$

[13]

9. (a)



Circle
Correct circle.
(centre (0, 3), radius 3)

M1
A1 2

(b) Drawing correct **half**-line passing as shown

B1

Find either x or y coord of A .

M1A1

$$z = -\frac{3\sqrt{2}}{2} + \left(3 + \frac{3\sqrt{2}}{2}\right)i$$

A1 4

[Algebraic approach, i.e. using $y = 3 - x$ and equation of circle will only gain M1A1, unless the second solution is ruled out, when B1 can be given by implication, and final A1, if correct]

(c) $|z - 3i| = 3 \rightarrow \left| \frac{2i}{\omega} - 3i \right| = 3$

M1

$$\Rightarrow \frac{|2i - 3i\omega|}{|\omega|} = 3$$

A1

$$\Rightarrow |\omega - \frac{2}{3}| = |\omega|$$

M1A1

Line with equation $u = \frac{1}{3}$ ($x = \frac{1}{3}$)

A1 5
[11]

Some alternatives:

(i) $\omega = \frac{2i}{x + iy} = \frac{2i(x - iy)}{x^2 + y^2} \Rightarrow u = \frac{2y}{x^2 + y^2}, v = \frac{2x}{x^2 + y^2}$

M1A1

$$\text{As } x^2 + y^2 - 6y = 0, u = \frac{1}{3}$$

M1, A1A1

(ii) $\omega = \frac{2i}{3\cos\theta + 3i(1 + \sin\theta)} = \frac{2i\{\cos\theta - i(1 + \sin\theta)\}}{3\{\cos^2\theta + (1 + \sin\theta)^2\}}$

M1A1

$$= \frac{2}{3} \frac{(1 + \sin\theta) + i\cos\theta}{2 + 2\sin\theta}, = \frac{1}{3} + i \frac{\cos\theta}{1 + \sin\theta},$$

M1A1

So locus is line $u = \frac{1}{3}$

A1

10. (a) $z^n = e^{in\theta} = (\cos n\theta + i \sin n\theta)$, $z^{-n} = e^{-in\theta} = \overline{\cos n\theta - i \sin n\theta}$ M1
 Completion (needs to be convincing) $z^n - \overline{z^n} = 2i \sin n\theta$ (*) AG A1 2
- (b) $\left(z - \frac{1}{z}\right)^5 = z^5 - 5z^3 + 10z - \frac{10}{z} + \frac{5}{z^3} - \frac{1}{z^5}$ M1A1
 $= \left(z^5 - \frac{1}{z^5}\right) - 5\left(z^3 - \frac{1}{z^3}\right) + 10\left(z - \frac{1}{z}\right)$
 $(2i \sin \theta)^5 = 32i \sin^5 \theta = 2i \sin 5\theta - 10i \sin 3\theta + 20i \sin \theta$ M1A1
 $\Rightarrow \sin^5 \theta = \frac{1}{16} (\sin 5\theta - 5 \sin 3\theta + 10 \sin \theta)$ (*) AG A1 5
- (c) Finding $\sin^5 \theta = \frac{1}{4} \sin \theta$ M1
 $\theta = 0, \pi$ (both) B1
 $(\sin^4 \theta = \frac{1}{4}) \Rightarrow \sin \theta = \pm \frac{1}{\sqrt{2}}$ M1
 $\theta = \frac{\pi}{4}, \frac{3\pi}{4}, \frac{5\pi}{4}, \frac{7\pi}{4}$ A1;A1 5
 [12]
11. (a) $\left(\frac{d^2 y}{dx^2}\right)_0 = \frac{1}{4}$ B1 1
- (b) $\left(\frac{dy}{dx}\right)_0 \approx \frac{y_1 - y_{-1}}{2h} \Rightarrow \frac{1}{2} \approx \frac{y_1 - y_{-1}}{0.2} \Rightarrow y_1 - y_{-1} \approx 0.1$ M1A1
 $\left(\frac{d^2 y}{dx^2}\right)_0 \approx \frac{y_1 - 2y_0 + y_{-1}}{h^2} \Rightarrow \frac{1}{4} \approx \frac{y_1 - 2 + y_{-1}}{0.01}$ M1
 $\Rightarrow y_1 + y_{-1} \approx 2.0025$ A1
 Adding to give $y_1 \approx 1.05125$ M1A1 6
- (c) Diff: $4(1+x^2) \frac{d^3 y}{dx^3} + 8x \frac{d^2 y}{dx^2} + 4x \frac{d^2 y}{dx^2} + 4 \frac{dy}{dx} = \frac{dy}{dx}$ M1A1
 Substituting appropriate vales $\Rightarrow 4 \left(\frac{d^3 y}{dx^3}\right)_0 = -\frac{3}{2} \Rightarrow \left(\frac{d^3 y}{dx^3}\right)_0 = -\frac{3}{8}$ M1A1 4
- (d) $y = y_0 + y_0'x + \frac{y_0''}{2!}x^2 + \frac{y_0'''}{3!}x^3 + \dots = 1 + \frac{1}{2}x + \frac{1}{8}x^2 - \frac{1}{16}x^3 + \dots$ M1A1ft2
- (e) 1.05119 A1 1
 [14]