

OXFORD CAMBRIDGE AND RSA EXAMINATIONS

**Advanced Subsidiary General Certificate of Education
Advanced General Certificate of Education**

MEI STRUCTURED MATHEMATICS

4756

Further Methods for Advanced Mathematics (FP2)

Tuesday

6 JUNE 2006

Afternoon

1 hour 30 minutes

Additional materials:

8 page answer booklet

Graph paper

MEI Examination Formulae and Tables (MF2)

TIME 1 hour 30 minutes

INSTRUCTIONS TO CANDIDATES

- Write your name, centre number and candidate number in the spaces provided on the answer booklet.
- Answer **all** the questions in Section A and **one** question from section B.
- You are permitted to use a graphical calculator in this paper.
- Final answers should be given to a degree of accuracy appropriate to the context.

INFORMATION FOR CANDIDATES

- The number of marks is given in brackets [] at the end of each question or part question.
- You are advised that an answer may receive **no marks** unless you show sufficient detail of the working to indicate that a correct method is being used.
- The total number of marks for this paper is 72.

This question paper consists of 4 printed pages.

2

Section A (54 marks)

Answer all the questions

- 1 (a) A curve has polar equation $r = a(\sqrt{2} + 2\cos\theta)$ for $-\frac{3}{4}\pi \leq \theta \leq \frac{3}{4}\pi$, where a is a positive constant.

(i) Sketch the curve. [2]

(ii) Find, in an exact form, the area of the region enclosed by the curve. [7]

- (b) (i) Find the Maclaurin series for the function $f(x) = \tan\left(\frac{1}{4}\pi + x\right)$, up to the term in x^2 . [6]

(ii) Use the Maclaurin series to show that, when h is small,

$$\int_{-h}^h x^2 \tan\left(\frac{1}{4}\pi + x\right) dx \approx \frac{2}{3}h^3 + \frac{4}{5}h^5. \quad [3]$$

- 2 (a) (i) Given that $z = \cos\theta + j\sin\theta$, express $z^n + \frac{1}{z^n}$ and $z^n - \frac{1}{z^n}$ in simplified trigonometric form. [2]

(ii) By considering $\left(z - \frac{1}{z}\right)^4 \left(z + \frac{1}{z}\right)^2$, find A , B , C and D such that

$$\sin^4\theta \cos^2\theta = A \cos 6\theta + B \cos 4\theta + C \cos 2\theta + D. \quad [6]$$

- (b) (i) Find the modulus and argument of $4 + 4j$. [2]

(ii) Find the fifth roots of $4 + 4j$ in the form $re^{j\theta}$, where $r > 0$ and $-\pi < \theta \leq \pi$.

Illustrate these fifth roots on an Argand diagram. [6]

(iii) Find integers p and q such that $(p + jq)^5 = 4 + 4j$. [2]

3

- 3 (i) Find the inverse of the matrix $\begin{pmatrix} 4 & 1 & k \\ 3 & 2 & 5 \\ 8 & 5 & 13 \end{pmatrix}$, where $k \neq 5$. [6]

- (ii) Solve the simultaneous equations

$$\begin{aligned} 4x + y + 7z &= 12 \\ 3x + 2y + 5z &= m \\ 8x + 5y + 13z &= 0 \end{aligned}$$

giving x , y and z in terms of m . [5]

- (iii) Find the value of p for which the simultaneous equations

$$\begin{aligned} 4x + y + 5z &= 12 \\ 3x + 2y + 5z &= p \\ 8x + 5y + 13z &= 0 \end{aligned}$$

have solutions, and find the general solution in this case. [7]

Section B (18 marks)

Answer one question

Option 1: Hyperbolic functions

- 4 (i) Starting from the definitions of $\sinh x$ and $\cosh x$ in terms of exponentials, prove that

$$1 + 2 \sinh^2 x = \cosh 2x. \quad [3]$$

- (ii) Solve the equation

$$2 \cosh 2x + \sinh x = 5,$$

giving the answers in an exact logarithmic form. [6]

- (iii) Show that $\int_0^{\ln 3} \sinh^2 x \, dx = \frac{10}{9} - \frac{1}{2} \ln 3$. [5]

- (iv) Find the exact value of $\int_3^5 \sqrt{x^2 - 9} \, dx$. [4]

[Question 5 is printed overleaf.]

Option 2: Investigation of curves

This question requires the use of a graphical calculator.

5 A curve has parametric equations

$$x = \theta - k \sin \theta, \quad y = 1 - \cos \theta,$$

where k is a positive constant.

- (i) For the case $k = 1$, use your graphical calculator to sketch the curve. Describe its main features. [4]
- (ii) Sketch the curve for a value of k between 0 and 1. Describe briefly how the main features differ from those for the case $k = 1$. [3]
- (iii) For the case $k = 2$:
- (A) sketch the curve; [2]
- (B) find $\frac{dy}{dx}$ in terms of θ ; [2]
- (C) show that the width of each loop, measured parallel to the x -axis, is
- $$2\sqrt{3} - \frac{2\pi}{3}. \quad [5]$$
- (iv) Use your calculator to find, correct to one decimal place, the value of k for which successive loops just touch each other. [2]