

Thursday 21 June 2012 – Afternoon

A2 GCE MATHEMATICS (MEI)

4756 Further Methods for Advanced Mathematics (FP2)

QUESTION PAPER

Candidates answer on the Printed Answer Book.

OCR supplied materials:

- Printed Answer Book 4756
- MEI Examination Formulae and Tables (MF2)

Other materials required:

- Scientific or graphical calculator

Duration: 1 hour 30 minutes



INSTRUCTIONS TO CANDIDATES

These instructions are the same on the Printed Answer Book and the Question Paper.

- The Question Paper will be found in the centre of the Printed Answer Book.
- Write your name, centre number and candidate number in the spaces provided on the Printed Answer Book. Please write clearly and in capital letters.
- **Write your answer to each question in the space provided in the Printed Answer Book.** Additional paper may be used if necessary but you must clearly show your candidate number, centre number and question number(s).
- Use black ink. HB pencil may be used for graphs and diagrams only.
- Read each question carefully. Make sure you know what you have to do before starting your answer.
- Answer **all** the questions in Section A and **one** question from Section B.
- Do **not** write in the bar codes.
- You are permitted to use a scientific or graphical calculator in this paper.
- Final answers should be given to a degree of accuracy appropriate to the context.

INFORMATION FOR CANDIDATES

This information is the same on the Printed Answer Book and the Question Paper.

- The number of marks is given in brackets [] at the end of each question or part question on the Question Paper.
- You are advised that an answer may receive **no marks** unless you show sufficient detail of the working to indicate that a correct method is being used.
- The total number of marks for this paper is **72**.
- The Printed Answer Book consists of **16** pages. The Question Paper consists of **4** pages. Any blank pages are indicated.

INSTRUCTION TO EXAMS OFFICER/INVIGILATOR

- Do not send this Question Paper for marking; it should be retained in the centre or recycled. Please contact OCR Copyright should you wish to re-use this document.

Section A (54 marks)

Answer all the questions

- 1 (a) (i) Differentiate the equation $\sin y = x$ with respect to x , and hence show that the derivative of $\arcsin x$ is $\frac{1}{\sqrt{1-x^2}}$. [4]

- (ii) Evaluate the following integrals, giving your answers in exact form.

$$(A) \int_{-1}^1 \frac{1}{\sqrt{2-x^2}} dx \quad [3]$$

$$(B) \int_{-\frac{1}{2}}^{\frac{1}{2}} \frac{1}{\sqrt{1-2x^2}} dx \quad [4]$$

- (b) A curve has polar equation $r = \tan \theta$, $0 \leq \theta < \frac{1}{2}\pi$. The points on the curve have cartesian coordinates (x, y) . A sketch of the curve is given in Fig. 1.




Fig. 1

Show that $x = \sin \theta$ and that $r^2 = \frac{x^2}{1-x^2}$.

Hence show that the cartesian equation of the curve is

$$y = \frac{x^2}{\sqrt{1-x^2}}.$$

Give the cartesian equation of the asymptote of the curve. [7]

3

- 2 (a) (i) Given that $z = \cos \theta + j \sin \theta$, express $z^n + \frac{1}{z^n}$ and $z^n - \frac{1}{z^n}$ in simplified trigonometric form. [2]

- (ii) Beginning with an expression for $\left(z + \frac{1}{z}\right)^4$, find the constants A, B, C in the identity

$$\cos^4 \theta \equiv A + B \cos 2\theta + C \cos 4\theta. \quad [4]$$

- (iii) Use the identity in part (ii) to obtain an expression for $\cos 4\theta$ as a polynomial in $\cos \theta$. [2]

- (b) (i) Given that $z = 4e^{j\pi/3}$ and that $w^2 = z$, write down the possible values of w in the form $re^{j\theta}$, where $r > 0$. Show z and the possible values of w in an Argand diagram. [5]

- (ii) Find the least positive integer n for which z^n is real.

Show that there is no positive integer n for which z^n is imaginary.

For each possible value of w , find the value of w^3 in the form $a + jb$ where a and b are real. [5]

- 3 (i) Find the value of a for which the matrix

$$\mathbf{M} = \begin{pmatrix} 1 & 2 & 3 \\ -1 & a & 4 \\ 3 & -2 & 2 \end{pmatrix}$$

does not have an inverse.

Assuming that a does not have this value, find the inverse of \mathbf{M} in terms of a . [7]

- (ii) Hence solve the following system of equations.

$$\begin{aligned} x + 2y + 3z &= 1 \\ -x &+ 4z = -2 \\ 3x - 2y + 2z &= 1 \end{aligned} \quad [4]$$

- (iii) Find the value of b for which the following system of equations has a solution.

$$\begin{aligned} x + 2y + 3z &= 1 \\ -x + 6y + 4z &= -2 \\ 3x - 2y + 2z &= b \end{aligned}$$

Find the general solution in this case and describe the solution geometrically. [7]

Section B (18 marks)**Answer one question***Option 1: Hyperbolic functions*

- 4 (i) Prove, from definitions involving exponential functions, that

$$\cosh 2u = 2 \sinh^2 u + 1. \quad [3]$$

- (ii) Prove that, if $y \geq 0$ and $\cosh y = u$, then $y = \ln(u + \sqrt{u^2 - 1})$. [4]

- (iii) Using the substitution $2x = \cosh u$, show that

$$\int \sqrt{4x^2 - 1} dx = ax\sqrt{4x^2 - 1} - b \operatorname{arcosh} 2x + c,$$

where a and b are constants to be determined and c is an arbitrary constant. [7]

- (iv) Find $\int_{\frac{1}{2}}^1 \sqrt{4x^2 - 1} dx$, expressing your answer in an exact form involving logarithms. [4]

Option 2: Investigation of curves

This question requires the use of a graphical calculator.

- 5 This question concerns curves with polar equation $r = \sec \theta + a$, where a is a constant.

- (i) State the set of values of θ between 0 and 2π for which r is undefined. [2]

For the rest of the question you should assume that θ takes all values between 0 and 2π for which r is defined.

- (ii) Use your graphical calculator to obtain a sketch of the curve in the case $a = 0$. Confirm the shape of the curve by writing the equation in cartesian form. [3]

- (iii) Sketch the curve in the case $a = 1$.

Now consider the curve in the case $a = -1$. What do you notice?

By considering both curves for $0 < \theta < \pi$ and $\pi < \theta < 2\pi$ separately, describe the relationship between the cases $a = 1$ and $a = -1$. [6]

- (iv) What feature does the curve exhibit for values of a greater than 1?

Sketch a typical case. [3]

- (v) Show that a cartesian equation of the curve $r = \sec \theta + a$ is $(x^2 + y^2)(x - 1)^2 = a^2 x^2$. [4]

Copyright Information

OCR is committed to seeking permission to reproduce all third-party content that it uses in its assessment materials. OCR has attempted to identify and contact all copyright holders whose work is used in this paper. To avoid the issue of disclosure of answer-related information to candidates, all copyright acknowledgements are reproduced in the OCR Copyright Acknowledgements Booklet. This is produced for each series of examinations and is freely available to download from our public website (www.ocr.org.uk) after the live examination series.

If OCR has unwittingly failed to correctly acknowledge or clear any third-party content in this assessment material, OCR will be happy to correct its mistake at the earliest possible opportunity.

For queries or further information please contact the Copyright Team, First Floor, 9 Hills Road, Cambridge CB2 1GE.

OCR is part of the Cambridge Assessment Group; Cambridge Assessment is the brand name of University of Cambridge Local Examinations Syndicate (UCLES), which is itself a department of the University of Cambridge.