



ADVANCED GCE
MATHEMATICS
 Further Pure Mathematics 2

4726

Candidates answer on the Answer Booklet

OCR Supplied Materials:

- 8 page Answer Booklet
- List of Formulae (MF1)

Other Materials Required:

- Scientific or graphical calculator

Thursday 27 May 2010
Morning

Duration: 1 hour 30 minutes



INSTRUCTIONS TO CANDIDATES

- Write your name clearly in capital letters, your Centre Number and Candidate Number in the spaces provided on the Answer Booklet.
- Use black ink. Pencil may be used for graphs and diagrams only.
- Read each question carefully and make sure that you know what you have to do before starting your answer.
- Answer **all** the questions.
- Do **not** write in the bar codes.
- Give non-exact numerical answers correct to 3 significant figures unless a different degree of accuracy is specified in the question or is clearly appropriate.
- You are permitted to use a graphical calculator in this paper.

INFORMATION FOR CANDIDATES

- The number of marks is given in brackets [] at the end of each question or part question.
- **You are reminded of the need for clear presentation in your answers.**
- The total number of marks for this paper is **72**.
- This document consists of **4** pages. Any blank pages are indicated.


1 It is given that $f(x) = \tan^{-1} 2x$ and $g(x) = p \tan^{-1} x$, where p is a constant. Find the value of p for which $f'(\frac{1}{2}) = g'(\frac{1}{2})$. [4]

2 Given that the first three terms of the Maclaurin series for $(1 + \sin x)e^{2x}$ are identical to the first three terms of the binomial series for $(1 + ax)^n$, find the values of the constants a and n . (You may use appropriate results given in the List of Formulae (MF1).) [6]

3 Use the substitution $t = \tan \frac{1}{2}x$ to show that

$$\int_0^{\frac{1}{3}\pi} \frac{1}{1 - \sin x} dx = 1 + \sqrt{3}. \quad [6]$$

4



The diagram shows the curve with equation

$$y = \frac{ax + b}{x + c},$$

where a , b and c are constants.

(i) Given that the asymptotes of the curve are $x = -1$ and $y = -2$ and that the curve passes through $(3, 0)$, find the values of a , b and c . [3]

(ii) Sketch the curve with equation

$$y^2 = \frac{ax + b}{x + c},$$

for the values of a , b and c found in part (i). State the coordinates of any points where the curve crosses the axes, and give the equations of any asymptotes. [4]

5 It is given that, for $n \geq 0$,

$$I_n = \int_0^{\frac{1}{2}} (1 - 2x)^n e^x dx.$$

(i) Prove that, for $n \geq 1$,

$$I_n = 2nI_{n-1} - 1. \quad [4]$$


(ii) Find the exact value of I_3 . [4]

6 (i) Show that $\frac{d}{dx}(\sinh^{-1} x) = \frac{1}{\sqrt{x^2 + 1}}$. [2]

(ii) Given that $y = \cosh(a \sinh^{-1} x)$, where a is a constant, show that

$$(x^2 + 1) \frac{d^2y}{dx^2} + x \frac{dy}{dx} - a^2y = 0. \quad [5]$$

7



The line $y = x$ and the curve $y = 2 \ln(3x - 2)$ meet where $x = \alpha$ and $x = \beta$, as shown in the diagram.

(i) Use the iteration $x_{n+1} = 2 \ln(3x_n - 2)$, with initial value $x_1 = 5.25$, to find the value of β correct to 2 decimal places. Show all your working. [2]

(ii) With the help of a ‘staircase’ diagram, explain why this iteration will not converge to α , whatever value of x_1 (other than α) is used. [3]

(iii) Show that the equation $x = 2 \ln(3x - 2)$ can be rewritten as $x = \frac{1}{3}(e^{\frac{1}{2}x} + 2)$. Use the Newton-Raphson method, with $f(x) = \frac{1}{3}(e^{\frac{1}{2}x} + 2) - x$ and $x_1 = 1.2$, to find α correct to 2 decimal places. Show all your working. [4]

(iv) Given that $x_1 = \ln 36$, explain why the Newton-Raphson method would not converge to a root of $f(x) = 0$. [2]

[Questions 8 and 9 are printed overleaf.]


- 8 (i) Using the definition of $\cosh x$ in terms of e^x and e^{-x} , show that

$$4 \cosh^3 x - 3 \cosh x \equiv \cosh 3x. \quad [4]$$

- (ii) Use the substitution $u = \cosh x$ to find, in terms of $5^{\frac{1}{3}}$, the real root of the equation

$$20u^3 - 15u - 13 = 0. \quad [6]$$

9



The diagram shows the curve with equation $y = \sqrt{2x + 1}$ between the points $A(-\frac{1}{2}, 0)$ and $B(4, 3)$.

- (i) Find the area of the region bounded by the curve, the x -axis and the line $x = 4$. Hence find the area of the region bounded by the curve and the lines OA and OB , where O is the origin. [4]
- (ii) Show that the curve between B and A can be expressed in polar coordinates as

$$r = \frac{1}{1 - \cos \theta}, \quad \text{where } \tan^{-1}\left(\frac{3}{4}\right) \leq \theta \leq \pi. \quad [5]$$

- (iii) Deduce from parts (i) and (ii) that $\int_{\tan^{-1}(\frac{3}{4})}^{\pi} \operatorname{cosec}^4\left(\frac{1}{2}\theta\right) d\theta = 24$. [4]



Copyright Information

OCR is committed to seeking permission to reproduce all third-party content that it uses in its assessment materials. OCR has attempted to identify and contact all copyright holders whose work is used in this paper. To avoid the issue of disclosure of answer-related information to candidates, all copyright acknowledgements are reproduced in the OCR Copyright Acknowledgements Booklet. This is produced for each series of examinations, is given to all schools that receive assessment material and is freely available to download from our public website (www.ocr.org.uk) after the live examination series.

If OCR has unwittingly failed to correctly acknowledge or clear any third-party content in this assessment material, OCR will be happy to correct its mistake at the earliest possible opportunity. For queries or further information please contact the Copyright Team, First Floor, 9 Hills Road, Cambridge CB2 1GE.

OCR is part of the Cambridge Assessment Group; Cambridge Assessment is the brand name of University of Cambridge Local Examinations Syndicate (UCLES), which is itself a department of the University of Cambridge.