

June 2009
6669 Further Pure Mathematics FP3 (new)
Mark Scheme

Question Number	Scheme	Marks
Q1	$\frac{7}{\cosh x} - \frac{\sinh x}{\cosh x} = 5 \Rightarrow \frac{14}{e^x + e^{-x}} - \frac{(e^x - e^{-x})}{e^x + e^{-x}} = 5$ $\therefore 14 - (e^x - e^{-x}) = 5(e^x + e^{-x}) \Rightarrow 6e^x - 14 + 4e^{-x} = 0$ $\therefore 3e^{2x} - 7e^x + 2 = 0 \Rightarrow (3e^x - 1)(e^x - 2) = 0$ $\therefore e^x = \frac{1}{3} \text{ or } 2$ $x = \ln\left(\frac{1}{3}\right) \text{ or } \ln 2$	M1 A1 M1 A1 B1ft [5]
Alternative (i)	Write $7 - \sinh x = 5 \cosh x$, then use exponential substitution $7 - \frac{1}{2}(e^x - e^{-x}) = \frac{5}{2}(e^x + e^{-x})$ Then proceed as method above.	M1
Alternative (ii)	$(7 \operatorname{sech} x - 5)^2 = \tanh^2 x = 1 - \operatorname{sech}^2 x$ $50 \operatorname{sech}^2 x - 70 \operatorname{sech} x + 24 = 0$ $2(5 \operatorname{sech} x - 3)(5 \operatorname{sech} x - 4) = 0$ $\operatorname{sech} x = \frac{3}{5} \text{ or } \operatorname{sech} x = \frac{4}{5}$ $x = \ln\left(\frac{1}{3}\right) \text{ or } \ln 2$	M1 A1 M1 A1 B1ft
Q2 (a)	$\mathbf{b} \times \mathbf{c} = 0\mathbf{i} + 5\mathbf{j} + 5\mathbf{k}$	M1 A1 A1 (3)
(b)	$\mathbf{a} \cdot (\mathbf{b} \times \mathbf{c}) = 0 + 5 = 5$	M1 A1 ft (2)
(c)	Area of triangle $OBC = \frac{1}{2} 5\mathbf{j} + 5\mathbf{k} = \frac{5}{2} \sqrt{2}$	M1 A1 (2)
(d)	Volume of tetrahedron = $\frac{1}{6} \times 5 = \frac{5}{6}$	B1 ft (1) [8]

Question Number	Scheme	Marks
<p>Q3 (a)</p> $\begin{vmatrix} 6-\lambda & 1 & -1 \\ 0 & 7-\lambda & 0 \\ 3 & -1 & 2-\lambda \end{vmatrix} = 0 \quad \therefore (6-\lambda)(7-\lambda)(2-\lambda) + 3(7-\lambda) = 0$ <p>$(7-\lambda) = 0$ verifies $\lambda = 7$ is an eigenvalue (can be seen anywhere)</p> <p>$\therefore (7-\lambda)\{12-8\lambda+\lambda^2+3\} = 0 \quad \therefore (7-\lambda)\{\lambda^2-8\lambda+15\} = 0$</p> <p>$\therefore (7-\lambda)(\lambda-5)(\lambda-3) = 0$ and 3 and 5 are the other two eigenvalues</p> <p>(b)</p> $\text{Set } \begin{pmatrix} 6 & 1 & -1 \\ 0 & 7 & 0 \\ 3 & -1 & 2 \end{pmatrix} \begin{pmatrix} x \\ y \\ z \end{pmatrix} = 7 \begin{pmatrix} x \\ y \\ z \end{pmatrix} \quad \text{or} \quad \begin{pmatrix} -1 & 1 & -1 \\ 0 & 0 & 0 \\ 3 & -1 & -5 \end{pmatrix} \begin{pmatrix} x \\ y \\ z \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \\ 0 \end{pmatrix}$ <p>Solve $-x + y - z = 0$ and $3x - y - 5z = 0$ to obtain $x = 3z$ or $y = 4z$ and a second equation which can contain 3 variables</p> <p>Obtain eigenvector as $3\mathbf{i} + 4\mathbf{j} + \mathbf{k}$ (or multiple)</p>		<p>M1</p> <p>M1</p> <p>A1</p> <p>M1 A1</p> <p>(5)</p> <p>M1</p> <p>M1 A1</p> <p>A1</p> <p>(4)</p> <p>[9]</p>

Question Number	Scheme	Marks
Q4 (a)	$\frac{dy}{dx} = \frac{1}{2} x^{-\frac{1}{2}} \times \frac{1}{\sqrt{1+(\sqrt{x})^2}}$ $\therefore \frac{dy}{dx} = \frac{\frac{1}{2} x^{-\frac{1}{2}}}{\sqrt{1+x}} \quad \left(= \frac{1}{2\sqrt{x(1+x)}} \right)$	B1, M1 A1 (3)
(b)	$\therefore \int_{\frac{1}{4}}^4 \frac{1}{\sqrt{x(x+1)}} dx = [2\operatorname{ar sinh} \sqrt{x}]_{\frac{1}{4}}^4$ $= [2\operatorname{ar sinh} 2 - 2\operatorname{ar sinh}(\frac{1}{2})]$ $= [2\ln(2 + \sqrt{5})] - [2\ln(\frac{1}{2} + \sqrt{\frac{5}{4}})]$ $2\ln \frac{(2 + \sqrt{5})}{(\frac{1}{2} + \sqrt{\frac{5}{4}})} = 2\ln \frac{2(2 + \sqrt{5})}{(1 + \sqrt{5})} = 2\ln \frac{2(\sqrt{5} + 2)(\sqrt{5} - 1)}{(\sqrt{5} + 1)(\sqrt{5} - 1)} = 2\ln \frac{(3 + \sqrt{5})}{2}$ $= \ln \frac{(3 + \sqrt{5})(3 + \sqrt{5})}{4} = \ln \frac{14 + 6\sqrt{5}}{4} = \ln \left(\frac{7}{2} + \frac{3\sqrt{5}}{2} \right)$	M1 M1 M1 M1 A1 A1 (6) [9]
Alternative (i) for part (a)	<p>Use $\sinh y = \sqrt{x}$ and state $\cosh y \frac{dy}{dx} = \frac{1}{2} x^{-\frac{1}{2}}$</p> $\therefore \frac{dy}{dx} = \frac{\frac{1}{2} x^{-\frac{1}{2}}}{\sqrt{1 + \sinh^2 y}} = \frac{\frac{1}{2} x^{-\frac{1}{2}}}{\sqrt{1 + (\sqrt{x})^2}}$ $\therefore \frac{dy}{dx} = \frac{\frac{1}{2} x^{-\frac{1}{2}}}{\sqrt{1+x}} \quad \left(= \frac{1}{2\sqrt{x(1+x)}} \right)$	B1 M1 A1 (3)
Alternative (i) for part (b) Alternative (ii) for part (b)	<p>Use $x = \tan^2 \theta$, $\frac{dx}{d\theta} = 2 \tan \theta \sec^2 \theta$ to give $2 \int \sec \theta d\theta = [2 \ln(\sec \theta + \tan \theta)]$</p> $= [2 \ln(\sec \theta + \tan \theta)]_{\tan \theta = \frac{1}{2}}^{\tan \theta = 2}$ <p>i.e. use of limits</p> <p>then proceed as before from line 3 of scheme</p> <p>Use $\int \frac{1}{\sqrt{[(x + \frac{1}{2})^2 - \frac{1}{4}]}} dx = \operatorname{arcosh} \frac{x + \frac{1}{2}}{\frac{1}{2}}$</p> $= [\operatorname{arcosh} 9 - \operatorname{arcosh}(\frac{3}{2})]$ $= [\ln(9 + \sqrt{80})] - [\ln(\frac{3}{2} + \frac{1}{2}\sqrt{5})]$ $= \ln \frac{(9 + \sqrt{80})}{(\frac{3}{2} + \frac{1}{2}\sqrt{5})} = \ln \frac{2(9 + \sqrt{80})(3 - \sqrt{5})}{(3 + \sqrt{5})(3 - \sqrt{5})}$ $= \ln \frac{2(9 + 4\sqrt{5})(3 - \sqrt{5})}{(3 + \sqrt{5})(3 - \sqrt{5})} = \ln \left(\frac{7}{2} + \frac{3\sqrt{5}}{2} \right)$	M1 M1 M1 M1 M1 A1 A1 (6) [9]

Question Number	Scheme	Marks
<p>Q5 (a)</p> <p>(b)</p> <p>(c)</p>	<p>$-(25-x^2)^{\frac{1}{2}} + c$</p> <p>$I_n = \int x^{n-1} \cdot \frac{x}{\sqrt{(25-x^2)}} dx = -x^{n-1} \sqrt{25-x^2} + \int (n-1)x^{n-2} \sqrt{(25-x^2)} dx$</p> <p>$I_n = \left[-x^{n-1} \sqrt{25-x^2} \right]_0^5 + \int_0^5 \frac{(n-1)x^{n-2}(25-x^2)}{\sqrt{(25-x^2)}} dx$</p> <p>$I_n = 0 + 25(n-1) I_{n-2} - (n-1) I_n$</p> <p>$\therefore nI_n = 25(n-1)I_{n-2}$ and so $I_n = \frac{25(n-1)}{n} I_{n-2}$ *</p> <p>$I_0 = \int_0^5 \frac{1}{\sqrt{(25-x^2)}} dx = \left[\arcsin\left(\frac{x}{5}\right) \right]_0^5 = \frac{\pi}{2}$</p> <p>$I_4 = \frac{25 \times 3}{4} \times \frac{25 \times 1}{2} I_0 = \frac{1875}{16} \pi$</p>	<p>M1A1 (2)</p> <p>M1 A1ft</p> <p>M1</p> <p>M1</p> <p>A1 (5)</p> <p>M1 A1</p> <p>M1 A1 (4)</p> <p>[11]</p>
<p>Alternative for (b)</p>	<p>Using substitution $x = 5\sin\theta$</p> <p>$I_n = 5^n \int_0^{\frac{\pi}{2}} \sin^n \theta d\theta = \left[-5^n \sin^{n-1} \theta \cos \theta \right]_0^{\frac{\pi}{2}} + 5^n (n-1) \int_0^{\frac{\pi}{2}} \sin^{n-2} \theta \cos^2 \theta d\theta$</p> <p>$= \left[-5^n \sin^{n-1} \theta \cos \theta \right]_0^{\frac{\pi}{2}} + 5^n (n-1) \int_0^{\frac{\pi}{2}} \sin^{n-2} \theta (1 - \sin^2 \theta) d\theta$</p> <p>$I_n = 0 + 25(n-1) I_{n-2} - (n-1) I_n$</p> <p>$\therefore nI_n = 25(n-1)I_{n-2}$ and so $I_n = \frac{25(n-1)}{n} I_{n-2}$ *</p> <p>(need to see that $I_{n-2} = 5^{n-2} \int_0^{\frac{\pi}{2}} \sin^{n-2} \theta d\theta$ for final A1)</p>	<p>M1A1</p> <p>M1</p> <p>M1</p> <p>A1 (5)</p>

Question Number	Scheme	Marks
<p>Q6 (a)</p> <p>(b)</p> <p>(c)</p>	<p>$\frac{x^2}{a^2} - \frac{(mx+c)^2}{b^2} = 1$ and so $b^2x^2 - a^2(mx+c)^2 = a^2b^2$</p> <p>$\therefore (b^2 - a^2m^2)x^2 - 2a^2mcx - a^2(c^2 + b^2) = 0$</p> <p>Or $(a^2m^2 - b^2)x^2 + 2a^2mcx + a^2(c^2 + b^2) = 0$ *</p> <p>$(2a^2mc)^2 = 4(a^2m^2 - b^2) \times a^2(c^2 + b^2)$</p> <p>$4a^4m^2c^2 = -4a^2(b^2c^2 + b^4 - a^2m^2c^2 - a^2m^2b^2)$</p> <p>$c^2 = a^2m^2 - b^2$ or $a^2m^2 = b^2 + c^2$ *</p> <p>Substitute (1, 4) into $y = mx+c$ to give $4 = m + c$ and</p> <p>Substitute $a = 5$ and $b = 4$ into $c^2 = a^2m^2 - b^2$ to give $c^2 = 25m^2 - 16$</p> <p>Solve simultaneous equations to eliminate m or $c : (4 - m)^2 = 25m^2 - 16$</p> <p>To obtain $24m^2 + 8m - 32 = 0$</p> <p>Solve to obtain $8(3m + 4)(m - 1) = 0 \dots m = \dots$</p> <p>$m = 1$ or $-\frac{4}{3}$</p> <p>Substitute to get $c = 3$ or $\frac{16}{3}$</p> <p>Lines are $y = x + 3$ and $3y + 4x = 16$</p>	<p>M1</p> <p>A1 (2)</p> <p>M1</p> <p>A1 (2)</p> <p>B1 M1 A1 M1 A1 M1 A1</p> <p>(7) [11]</p>

Question Number	Scheme	Marks
Q7 (a)	<p>If the lines meet, $-1+3\lambda = -4+3\mu$ and $2+4\lambda = 2\mu$</p> <p>Solve to give $\lambda = 0$ ($\mu = 1$ but this need not be seen).</p> <p>Also $1-\lambda = \alpha$ and so $\alpha = 1$.</p> <p>(b) $\begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ -1 & 3 & 4 \\ 0 & 3 & 2 \end{vmatrix} = -6\mathbf{i}+2\mathbf{j}-3\mathbf{k}$ is perpendicular to both lines and hence to the plane</p> <p>The plane has equation $\mathbf{r}\cdot\mathbf{n}=\mathbf{a}\cdot\mathbf{n}$, which is $-6x + 2y - 3z = -14$, i.e. $-6x + 2y - 3z + 14 = 0$.</p>	<p>M1</p> <p>M1 A1</p> <p>B1</p> <p>(4)</p> <p>M1 A1</p> <p>M1</p> <p>A1 o.a.e.</p> <p>(4)</p>
OR (b)	<p>Alternative scheme</p> <p>Use $(1, -1, 2)$ and $(\alpha, -4, 0)$ in equation $ax+by+cz+d=0$</p> <p>And third point so three equations, and attempt to solve</p> <p>Obtain $6x - 2y + 3z =$ $(6x - 2y + 3z) - 14 = 0$</p>	<p>M1</p> <p>M1</p> <p>A1</p> <p>A1 o.a.e.</p> <p>(4)</p>
(c)	<p>$(\mathbf{a}_1 - \mathbf{a}_2) = \mathbf{i} - 3\mathbf{j} - 2\mathbf{k}$</p> <p>Use formula $\frac{(\mathbf{a}_1 - \mathbf{a}_2) \cdot \mathbf{n}}{ \mathbf{n} } = \frac{(\mathbf{i} - 3\mathbf{j} - 2\mathbf{k}) \cdot (-6\mathbf{i} + 2\mathbf{j} - 3\mathbf{k})}{\sqrt{(36+4+9)}} = \left(\frac{-6}{7}\right)$</p> <p>Distance is $\frac{6}{7}$</p>	<p>M1</p> <p>M1</p> <p>A1</p> <p>(3)</p> <p>[11]</p>

Question Number	Scheme	Marks
<p>Q8 (a)</p> <p>(b)</p>	<p>$\frac{dx}{d\theta} = -3\sin\theta, \frac{dy}{d\theta} = 5\cos\theta$</p> <p>so $S = 2\pi \int 5\sin\theta \sqrt{(-3\sin\theta)^2 + (5\cos\theta)^2} d\theta$</p> <p>$\therefore S = 2\pi \int 5\sin\theta \sqrt{9 - 9\cos^2\theta + 25\cos^2\theta} d\theta$</p> <p>Let $c = \cos\theta, \frac{dc}{d\theta} = -\sin\theta$, limits 0 and $\frac{\pi}{2}$ become 1 and 0</p> <p>So $S = k\pi \int_0^{\alpha} \sqrt{16c^2 + 9} dc$, where $k = 10$, and α is 1</p> <p>Let $c = \frac{3}{4}\sinh u$. Then $\frac{dc}{du} = \frac{3}{4}\cosh u$</p> <p>So $S = k\pi \int \sqrt{9\sinh^2 u + 9} \frac{3}{4}\cosh u du$</p> <p>$= k\pi \int \frac{9}{4}\cosh^2 u du = k\pi \int \frac{9}{8}(\cosh 2u + 1) du$</p> <p>$= k\pi \left[\frac{9}{16}\sinh 2u + \frac{9}{8}u \right]_0^d$</p> <p>$= \frac{45\pi}{4} \left[\frac{20}{9} + \ln 3 \right]$ i.e. <u>117</u></p>	<p>B1</p> <p>M1</p> <p>M1</p> <p>M1</p> <p>A1, A1 (6)</p> <p>M1</p> <p>A1</p> <p>M1</p> <p>A1</p> <p>B1</p> <p>(5)</p> <p>[11]</p>

June 2010
Further Pure Mathematics FP3 6669
Mark Scheme

Question Number	Scheme	Marks
1.	$\pm \frac{a}{e} = 8, \quad \pm ae = 2$ $\frac{a}{e} \times ae = a^2 = 16$ $a = 4$ $b^2 = a^2(1 - e^2) = a^2 - a^2e^2$ $\Rightarrow b^2 = 16 - 4 = 12$ $\Rightarrow b = \sqrt{12} = 2\sqrt{3}$	B1, B1 B1 M1 A1 (5) 5

Question Number	Scheme	Marks
2.	$x^2 + 4x + 13 = (x + 2)^2 + 9$ $\int \frac{1}{(x + 2)^2 + 9} dx = \frac{1}{3} \arctan\left(\frac{x + 2}{3}\right)$ $\left[\frac{1}{3} \arctan\left(\frac{x + 2}{3}\right)\right]_{-2}^1 = \frac{1}{3}(\arctan 1 - \arctan 0)$ $= \frac{\pi}{12}$	B1 M1 A1 M1 A1 (5) 5

Question Number	Scheme	Marks
<p>6(a)</p> <p>(b)</p> <p>(c)</p> <p>(d)</p>	$\begin{pmatrix} 1 & 0 & 3 \\ 0 & -2 & 1 \\ k & 0 & 1 \end{pmatrix} \begin{pmatrix} 6 \\ 1 \\ 6 \end{pmatrix} = \lambda \begin{pmatrix} 6 \\ 1 \\ 6 \end{pmatrix}$ $\begin{pmatrix} 24 \\ 4 \\ 6k+6 \end{pmatrix} = \begin{pmatrix} 6\lambda \\ \lambda \\ 6\lambda \end{pmatrix}$ <p>Uses the first or second row to obtain $\lambda = 4$</p> <p>Uses the third row and their $\lambda = 4$ to obtain $6k + 6 = 24 \Rightarrow k = 3$ *</p> $\begin{vmatrix} 1-\lambda & 0 & 3 \\ 0 & -2-\lambda & 1 \\ 3 & 0 & 1-\lambda \end{vmatrix} = 0$ $\Rightarrow (1-\lambda)((-2-\lambda)(1-\lambda)-0)-0(0(1-\lambda)-3)+3(0-3(-2-\lambda))=0$ $\Rightarrow (1-\lambda)(-2-\lambda)(1-\lambda)+9(2+\lambda)=(2+\lambda)(9-(1-\lambda)^2)=0$ $(\lambda^3 - 12\lambda - 16 = 0)$ $\Rightarrow (\lambda + 2)(\lambda^2 - 2\lambda - 8) = 0$ $\Rightarrow (\lambda + 2)(\lambda + 2)(\lambda - 4) = 0$ $\lambda = -2, 4$ <p>Parametric form of $l_1 : (t+2, -3t, 4t-1)$</p> $\begin{pmatrix} 1 & 0 & 3 \\ 0 & -2 & 1 \\ 3 & 0 & 1 \end{pmatrix} \begin{pmatrix} t+2 \\ -3t \\ 4t-1 \end{pmatrix} = \begin{pmatrix} 13t-1 \\ 10t-1 \\ 7t+5 \end{pmatrix}$ <p>Cartesian equations of $l_2 : \frac{x+1}{13} = \frac{y+1}{10} = \frac{z-5}{7}$</p>	<p>M1A1 (2)</p> <p>M1 A1 (2)</p> <p>M1 A1</p> <p>M1 A1 (4) M1</p> <p>M1 A1</p> <p>ddM1A1(5)</p> <p>13</p>


Question Number	Scheme	Marks
<p>7(a)</p> <p>(b)</p> <p>(c)</p>	$\begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ -4 & 1 & 0 \\ 6 & -2 & 1 \end{vmatrix} = \begin{pmatrix} 1 \\ 4 \\ 2 \end{pmatrix}$ $\begin{pmatrix} 1 \\ 4 \\ 2 \end{pmatrix} \cdot \begin{pmatrix} 3 \\ 0 \\ 1 \end{pmatrix} = 5$ $\mathbf{r} \cdot \begin{pmatrix} 1 \\ 4 \\ 2 \end{pmatrix} = 5$ <p>Equation of l is $\mathbf{r} = \begin{pmatrix} 6 \\ 13 \\ 5 \end{pmatrix} + t \begin{pmatrix} 1 \\ 4 \\ 2 \end{pmatrix}$</p> <p>At intersection $\begin{pmatrix} 6+t \\ 13+4t \\ 5+2t \end{pmatrix} \cdot \begin{pmatrix} 1 \\ 4 \\ 2 \end{pmatrix} = 5$</p> $\Rightarrow 6+t+4(13+4t)+2(5+2t)=5 \Rightarrow t=-3$ <p>N is $(3,1,-1)$ *</p> $\overrightarrow{PN} \cdot \overrightarrow{PR} = (-3\mathbf{i} - 12\mathbf{j} - 6\mathbf{k}) \cdot (-5\mathbf{i} - 13\mathbf{j} - 3\mathbf{k}) = 189$ $\sqrt{9+144+36}\sqrt{25+169+9} \cos NPR = 189$ $NX = NP \sin NPR = \sqrt{189} \sin NPR = 3.61$	<p>M1 A2(1,0)</p> <p>M1A1 (5)</p> <p>M1</p> <p>M1</p> <p>M1</p> <p>A1 (4)</p> <p>M1 A1ft</p> <p>A1</p> <p>M1A1 (5)</p> <p>14</p>

Question Number	Scheme	Marks
<p>8(a)</p> <p>(b)</p>	$\frac{dx}{dt} = 4 \sec t \tan t \quad \frac{dy}{dt} = 2 \sec^2 t$ $\frac{dy}{dx} = \frac{2 \sec^2 t}{4 \sec t \tan t} \quad \left(= \frac{1}{2 \sin t} \right)$ $y - 2 \tan t = \frac{1}{2 \sin t} (x - 4 \sec t)$ $2y \sin t - \frac{4 \sin^2 t}{\cos t} = x - \frac{4}{\cos t}$ $2y \sin t = x - \frac{4 - 4 \sin^2 t}{\cos t} = x - 4 \cos t \quad *$ <p>Gradient of l_2 is $-2 \sin t$</p> $y = -2x \sin t \quad (2)$ $2(-2x \sin t) \sin t = x - 4 \cos t \Rightarrow x = \frac{4 \cos t}{1 + 4 \sin^2 t} \quad (1)$ $y = \frac{-8 \sin t \cos t}{1 + 4 \sin^2 t}$ $(x^2 + y^2)^2 = \left(\frac{16 \cos^2 t}{(1 + 4 \sin^2 t)^2} + \frac{64 \sin^2 t \cos^2 t}{(1 + 4 \sin^2 t)^2} \right)^2$ $= \frac{256 \cos^4 t}{(1 + 4 \sin^2 t)^4} (1 + 4 \sin^2 t)^2 = \frac{256 \cos^4 t}{(1 + 4 \sin^2 t)^2}$ $16x^2 - 4y^2 = \frac{256 \cos^2 t}{(1 + 4 \sin^2 t)^2} - \frac{256 \sin^2 t \cos^2 t}{(1 + 4 \sin^2 t)^2} = \frac{256 \cos^4 t}{(1 + 4 \sin^2 t)^2}$	<p>B1 (both)</p> <p>M1</p> <p>M1 A1</p> <p>A1 (5)</p> <p>M1</p> <p>A1</p> <p>M1 A1</p> <p>M1 A1</p> <p>M1</p> <p>A1 (8)</p> <p>13</p>

June 2011
Further Pure Mathematics FP3 6669
Mark Scheme

Question Number	Scheme	Marks
1.	$\frac{dy}{dx} = 6x^2 \text{ and so surface area} = 2\pi \int 2x^3 \sqrt{1+(6x^2)^2} dx$ $= 4\pi \left[\frac{2}{3 \times 36 \times 4} (1+36x^4)^{\frac{3}{2}} \right]$ <p>Use limits 2 and 0 to give $\frac{4\pi}{216} [13860.016 - 1] = 806$ (to 3 sf)</p>	B1 M1 A1 DM1 A1 5
B1 1M1 1A1 2DM1 2A1	<p style="text-align: center;">Notes:</p> <p>Both bits CAO but condone lack of 2π</p> <p>Integrating $\int \left(y \sqrt{1 + \left(\text{their } \frac{dy}{dx} \right)^2} \right) dx$, getting $k(1+36x^4)^{\frac{3}{2}}$, condone lack of 2π</p> <p>If they use a substitution it must be a complete method.</p> <p>CAO</p> <p>Correct use of 2 and 0 as limits</p> <p>CAO</p>	
2. (a) (i) (ii)	$\frac{dy}{dx} = \frac{x}{\sqrt{1-x^2}} + \arcsin x$ $\text{At given value derivative} = \frac{1}{\sqrt{3}} + \frac{\pi}{6} = \frac{2\sqrt{3} + \pi}{6}$	M1 A1 B1 (2) (1)
(b)	$\frac{dy}{dx} = \frac{6e^{2x}}{1+9e^{4x}}$ $= \frac{6}{e^{-2x} + 9e^{2x}}$ $= \frac{3}{\frac{5}{2}(e^{2x} + e^{-2x}) + \frac{4}{2}(e^{2x} - e^{-2x})}$ $\therefore \frac{dy}{dx} = \frac{3}{5 \cosh 2x + 4 \sinh 2x} \quad *$	1M1 A1 2M1 3M1 A1 cso (5) 8
(a) M1 A1 B1	<p style="text-align: center;">Notes:</p> <p>Differentiating getting an arcsinx term and a $\frac{1}{\sqrt{1 \pm x^2}}$ term</p> <p>CAO</p> <p>CAO any correct form</p>	

Question Number	Scheme	Marks
<p>(b) 1M1 Integrating to get a ln or hyperbolic term 1A1 CAO 2DM1 Correctly using limits. 2A1 CAO</p>		
<p>4.</p> <p>(a)</p> $I_n = \left[\frac{x^3}{3} (\ln x)^n \right] - \int \frac{x^3}{3} \times \frac{n(\ln x)^{n-1}}{x} dx$ $= \left[\frac{x^3}{3} (\ln x)^n \right]_1^e - \int_1^e \frac{nx^2 (\ln x)^{n-1}}{3} dx$ $\therefore I_n = \frac{e^3}{3} - \frac{n}{3} I_{n-1} \quad *$		<p>M1 A1</p> <p>DM1</p> <p>A1cso</p> <p>(4)</p>
<p>(b)</p> $I_0 = \int_1^e x^2 dx = \left[\frac{x^3}{3} \right]_1^e = \frac{e^3}{3} - \frac{1}{3} \text{ or } I_1 = \frac{e^3}{3} - \frac{1}{3} \left(\frac{e^3}{3} - \frac{1}{3} \right) = \frac{2e^3}{9} + \frac{1}{9}$ $I_1 = \frac{e^3}{3} - \frac{1}{3} I_0, \quad I_2 = \frac{e^3}{3} - \frac{2}{3} I_1 \text{ and } I_3 = \frac{e^3}{3} - \frac{3}{3} I_2 \text{ so } I_3 = \frac{4e^3}{27} + \frac{2}{27}$ <p style="text-align: center;">Notes:</p> <p>(a)1M1 Using integration by parts, integrating x^2, differentiating $(\ln x)^n$ 1A1 CAO 2DM1 Correctly using limits 1 and e 2A1 CSO answer given</p> <p>(b)1M1 Evaluating I_0 or I_1 by an attempt to integrate something 1A1 CAO 2M1 Finding I_3 (also probably I_1 and I_2) If 'n's left in M0 2A1 I_3 CAO</p>		<p>M1 A1</p> <p>M1 A1</p> <p>(4)</p> <p>8</p>

Question Number	Scheme	Marks
<p>5. (a)</p>	 <p>Graph of $y = 3\sinh 2x$</p> <p>Shape of $-e^{2x}$ graph</p> <p>Asymptote: $y = 13$</p> <p>Value 10 on y axis and value 0.7 or $\frac{1}{2} \ln\left(\frac{13}{3}\right)$ on x axis</p>	<p>B1</p> <p>B1</p> <p>B1</p> <p>B1</p> <p>(4)</p>
<p>(b)</p>	<p>Use definition $\frac{3}{2}(e^{2x} - e^{-2x}) = 13 - 3e^{2x} \rightarrow 9e^{4x} - 26e^{2x} - 3 = 0$ to form quadratic</p> <p>$\therefore e^{2x} = -\frac{1}{9}$ or 3</p> <p>$\therefore x = \frac{1}{2} \ln(3)$</p>	<p>M1 A1</p> <p>DM1 A1</p> <p>B1</p> <p>(5)</p> <p>9</p>
<p>(a) 1B1</p> <p>2B1</p> <p>3B1</p> <p>4B1</p> <p>(b) 1M1</p> <p>1A1</p> <p>2DM1</p> <p>2A1</p> <p>B1</p>	<p style="text-align: center;">Notes:</p> <p>$y = 3\sinh 2x$ first and third quadrant.</p> <p>Shape of $y = -e^{2x}$ correct intersects on positive axes.</p> <p>Equation of asymptote, $y = 13$, given. Penalise 'extra' asymptotes here</p> <p>Intercepts correct both</p> <p>Getting a three terms quadratic in e^{2x}</p> <p>Correct three term quadratic</p> <p>Solving for e^{2x}</p> <p>CAO for e^{2x} condone omission of negative value.</p> <p>CAO one answer only</p>	

Question Number	Scheme	Marks
6.		
(a)	$\mathbf{n} = (2\mathbf{j} - \mathbf{k}) \times (3\mathbf{i} + 2\mathbf{j} + 2\mathbf{k}) = 6\mathbf{i} - 3\mathbf{j} - 6\mathbf{k}$ o.a.e. (e.g. $2\mathbf{i} - \mathbf{j} - 2\mathbf{k}$)	M1 A1 (2)
(b)	Line l has direction $2\mathbf{i} - 2\mathbf{j} - \mathbf{k}$ Angle between line l and normal is given by $(\cos \beta \text{ or } \sin \alpha) = \frac{4+2+2}{\sqrt{9}\sqrt{9}} = \frac{8}{9}$ $\alpha = 90 - \beta = 63$ degrees to nearest degree.	B1 M1 A1ft A1 awrt (4)
(c) Alt 1	Plane P has equation $\mathbf{r} \cdot (2\mathbf{i} - \mathbf{j} - 2\mathbf{k}) = 1$ Perpendicular distance is $\frac{1 - (-7)}{\sqrt{9}} = \frac{8}{3}$	M1 A1 M1 A1 (4)
(c) Alt 2	Parallel plane through A has equation $\mathbf{r} \cdot \frac{2\mathbf{i} - \mathbf{j} - 2\mathbf{k}}{3} = \frac{-7}{3}$ Plane P has equation $\mathbf{r} \cdot \frac{2\mathbf{i} - \mathbf{j} - 2\mathbf{k}}{3} = \frac{1}{3}$ So O lies between the two and perpendicular distance is $\frac{1}{3} + \frac{7}{3} = \frac{8}{3}$	M1 A1 M1 A1 (4)
(c) Alt 3	Distance A to $(3,1,2) = \sqrt{2^2 + 2^2 + 1^2} = 3$ Perpendicular distance is '3' $\sin \alpha = 3 \times \frac{8}{9} = \frac{8}{3}$	M1A1 M1A1 (4)
(c) Alt 4	Finding Cartesian equation of plane P: $2x - y - 2z - 1 = 0$ $d = \frac{ n_1\alpha + n_2\beta + n_3\gamma + d }{\sqrt{n_1^2 + n_2^2 + n_3^2}} = \frac{ 2(1) - 1(3) - 2(3) - 1 }{\sqrt{2^2 + 1^2 + 2^2}} = \frac{8}{3}$	M1 A1 M1A1 (4)
	Notes:	
(a) M1 A1	Cross product of the correct vectors CAO o.e.	
(b) B1 M1 1A1ft 2A1	CAO Angle between ' $2\mathbf{i} - \mathbf{j} - 2\mathbf{k}$ ' and $2\mathbf{i} - 2\mathbf{j} - \mathbf{k}$, formula of correct form 8/9ft CAO awrt	
(c) 1M1 1A1 2M1 2A1	Eqn of plane using $2\mathbf{i} - \mathbf{j} - 2\mathbf{k}$ or dist of A from O or finding length of AP Correct equation (must have =) or A to $(3,1,2) = 3$ Using correct method to find perpendicular distance CAO	

Question Number	Scheme	Marks
7. (a)	$\text{Det } \mathbf{M} = k(0 - 2) + 1(1 + 3) + 1(-2 - 0) = -2k + 4 - 2 = 2 - 2k$	M1 A1 (2)
(b)	$\mathbf{M}^T = \begin{pmatrix} k & 1 & 3 \\ -1 & 0 & -2 \\ 1 & -1 & 1 \end{pmatrix} \text{ so cofactors} = \begin{pmatrix} -2 & -1 & 1 \\ -4 & k-3 & k+1 \\ -2 & 2k-3 & 1 \end{pmatrix}$ <p>(-1 A mark for each term wrong)</p> $\mathbf{M}^{-1} = \frac{1}{2-2k} \begin{pmatrix} -2 & -1 & 1 \\ -4 & k-3 & k+1 \\ -2 & 2k-3 & 1 \end{pmatrix}$	M1 M1 A3 (5)
(c)	<p>Let (x, y, z) be on l_1. Equation of l_2 can be written as $\begin{pmatrix} x' \\ y' \\ z' \end{pmatrix} = \begin{pmatrix} 4 \\ 1 \\ 7 \end{pmatrix} + \lambda \begin{pmatrix} 4 \\ 1 \\ 3 \end{pmatrix}$.</p> <p>Use $\begin{pmatrix} x \\ y \\ z \end{pmatrix} = \mathbf{M}^{-1} \begin{pmatrix} x' \\ y' \\ z' \end{pmatrix}$ with $k = 2$. i.e. $\begin{pmatrix} x \\ y \\ z \end{pmatrix} = \frac{1}{-2} \begin{pmatrix} -2 & -1 & 1 \\ -4 & -1 & 3 \\ -2 & 1 & 1 \end{pmatrix} \begin{pmatrix} 4+4\lambda \\ 1+\lambda \\ 7+3\lambda \end{pmatrix}$</p> <p>$\therefore \begin{pmatrix} x \\ y \\ z \end{pmatrix} = \begin{pmatrix} 3\lambda+1 \\ 4\lambda-2 \\ 2\lambda \end{pmatrix}$</p> <p>and so $(\mathbf{r} - \mathbf{a}) \times \mathbf{b} = \mathbf{0}$ where $\mathbf{a} = \mathbf{i} - 2\mathbf{j}$ and $\mathbf{b} = 3\mathbf{i} + 4\mathbf{j} + 2\mathbf{k}$ or equivalent or $\mathbf{r} = \mathbf{a} + \lambda\mathbf{b}$ where $\mathbf{a} = \mathbf{i} - 2\mathbf{j}$ and $\mathbf{b} = 3\mathbf{i} + 4\mathbf{j} + 2\mathbf{k}$ or equivalent</p>	B1 M1 M1 A1 B1ft (5) 12
	<p style="text-align: center;">Notes:</p> <p>(a) M1 Finding determinant at least one component correct. A1 CAO</p> <p>(b) 1M1 Finding matrix of cofactors or its transpose 2M1 Finding inverse matrix, 1/(det) cofactors + transpose 1A1 At least seven terms correct (so at most 2 incorrect) condone missing det 2A1 At least eight terms correct (so at most 1 incorrect) condone missing det 3A1 All nine terms correct, condone missing det</p> <p>(c) 1B1 Equation of l_2 1M1 Using inverse transformation matrix correctly 2M1 Finding general point in terms of λ. A1 CAO for general point in terms of one parameter 2B1 ft for vector equation of their l_1</p>	

Question Number	Scheme	Marks
<p>8.</p> <p>(a)</p>	<p>Uses $\frac{dy}{dx} = \frac{\frac{dy}{d\theta}}{\frac{dx}{d\theta}} = \frac{b \cosh \theta}{a \sinh \theta}$ or $\frac{2x}{a^2} - \frac{2yy'}{b^2} = 0 \rightarrow y' = \frac{xb^2}{ya^2} = \frac{b \cosh \theta}{a \sinh \theta}$</p> <p>So $y - b \sinh \theta = \frac{b \cosh \theta}{a \sinh \theta} (x - a \cosh \theta)$</p> <p>$\therefore ab(\cosh^2 \theta - \sinh^2 \theta) = xb \cosh \theta - ya \sinh \theta$ and as $(\cosh^2 \theta - \sinh^2 \theta) = 1$</p> <p>$xb \cosh \theta - ya \sinh \theta = ab$ *</p>	<p>M1 A1</p> <p>M1</p> <p>A1cso</p> <p>(4)</p>
<p>(b)</p>	<p>P is the point $(\frac{a}{\cosh \theta}, 0)$</p>	<p>M1 A1</p> <p>(2)</p>
<p>(c)</p>	<p>l_2 has equation $x = a$ and meets l_1 at $Q(a, \frac{b(\cosh \theta - 1)}{\sinh \theta})$</p>	<p>M1 A1</p> <p>(2)</p>
<p>(d) Alt 1</p> <p>(d) Alt 2</p>	<p>The mid point of PQ is given by $X = \frac{a(\cosh \theta + 1)}{2 \cosh \theta}$, $Y = \frac{b(\cosh \theta - 1)}{2 \sinh \theta}$</p> <p>$4Y^2 + b^2 = b^2 \left(\frac{\cosh^2 \theta + 1 - 2 \cosh \theta + \sinh^2 \theta}{\sinh^2 \theta} \right)$</p> <p>$= b^2 \left(\frac{2 \cosh^2 \theta - 2 \cosh \theta}{\sinh^2 \theta} \right)$</p> <p>$X(4Y^2 + b^2) = ab^2 \left(\frac{(\cosh \theta + 1)(\cosh \theta - 1)2 \cosh \theta}{2 \cosh \theta \sinh^2 \theta} \right)$</p> <p>Simplify fraction by using $\cosh^2 \theta - \sinh^2 \theta = 1$ to give $x(4y^2 + b^2) = ab^2$ *</p> <p>First line of solution as before</p> <p>$4Y^2 + b^2 = b^2 (\coth^2 \theta + \operatorname{cosech}^2 \theta - 2 \coth \theta \operatorname{cosech} \theta + 1)$</p> <p>$= b^2 (2 \coth^2 \theta - 2 \coth \theta \operatorname{cosech} \theta)$</p> <p>$X(4Y^2 + b^2) = ab^2 (\coth \theta (\coth \theta - \operatorname{cosech} \theta) (1 + \operatorname{sech} \theta))$</p> <p>Simplify expansion by using $\coth^2 \theta - \operatorname{cosech}^2 \theta = 1$ to give $x(4y^2 + b^2) = ab^2$ *</p>	<p>1M1 A1ft</p> <p>2M1</p> <p>3M1</p> <p>4M1</p> <p>A1cso</p> <p>(6)</p> <p>1M1A1ft</p> <p>2M1</p> <p>3M1</p> <p>4M1</p> <p>A1cso</p> <p>(6)</p> <p>14</p>

Question Number	Scheme	Marks
8. (a)1M1 Finding gradient in terms of θ 1A1 CAO 2M1 Finding equation of tangent 2A1 CSO (answer given) look for $\pm(\cosh^2\theta - \sinh^2\theta)$ (b)M1 Putting $y = 0$ into their tangent A1ft P found, ft for their tangent o.e. (c) M1 Putting $x = a$ into their tangent. A1 CAO Q found o.e. (d) For Alt 1 and 2 1M1 Finding expressions, in terms of $\sinh \theta$ and $\cosh \theta$ but must be adding 1A1 Ft on their P and Q, 2M1 Finding $4y^2 + b^2$ 3M1 Simplified, factorised, maximum of 2 terms per bracket 4M1 Finding $x(4y^2 + b^2)$, completely factorised, maximum of 2 terms per bracket 2A1 CSO (d) For Alts 3, 4 and 5 1M1 Finding expressions, in terms of $\sinh \theta$ and $\cosh \theta$ but must be adding 1A1 Ft on their P and Q 2M1 Getting $\cosh \theta$ in terms of x 3M1 y or y^2 in terms of $\cosh \theta$ or $\sinh \theta$ in terms of x and y 4M1 Getting equation in terms of x and y only. No square roots. 2A1 CSO		

Question Number	Scheme	Marks
<p>8(d) Alt 3</p>	$X = \frac{a(\cosh \theta + 1)}{2 \cosh \theta}, \quad Y = \frac{b(\cosh \theta - 1)}{2 \sinh \theta}$ $\cosh \theta = \frac{a}{2x - a}$ $\sinh \theta = \frac{b(\cosh \theta - 1)}{2y} = \frac{b(a - x)}{(2x - a)y}$ $\left(\frac{a}{2x - a} \right)^2 - \left(\frac{b(a - x)}{(2x - a)y} \right)^2 = 1$ <p>Simplifies to give required equation $[y^2 4x(a - x) = b^2(a - x)^2, \quad x(4y^2 + b^2) = ab^2]$</p>	<p>As main scheme 1M1 A1ft</p> <p>cosh θ in terms of x 2M1</p> <p>sinh θ in terms of x and y 3M1</p> <p>Using $\cosh^2 \theta - \sinh^2 \theta = 1$ 4M1</p> <p>A1cso (6)</p>
<p>Alt 4</p>	$X = \frac{a(\cosh \theta + 1)}{2 \cosh \theta}, \quad Y = \frac{b(\cosh \theta - 1)}{2 \sinh \theta}$ $\cosh \theta = \frac{a}{2x - a}$ $y^2 = \frac{b^2(\cosh \theta - 1)^2}{4(\cosh^2 \theta - 1)} = \frac{b^2(\cosh \theta - 1)}{4(\cosh \theta + 1)}$ $y^2 = \frac{b^2 \left(\frac{2a - 2x}{2x - a} \right)^2}{4 \left(\frac{2x}{2x - a} \right)} \text{ o.e.}$ <p>Simplifies to give required equation</p>	<p>As main scheme 1M1 A1ft</p> <p>cosh θ in terms of x 2M1</p> <p>y^2 in terms of cosh θ only 3M1</p> <p>Forms equation in x and y only 4M1</p> <p>A1 cso (6)</p>
<p>Alt 5</p>	$X = \frac{a(\cosh \theta + 1)}{2 \cosh \theta}, \quad Y = \frac{b(\cosh \theta - 1)}{2 \sinh \theta}$ $\cosh \theta = \frac{a}{2x - a}$ $y = \left(\frac{b(\cosh \theta - 1)}{2 \sinh \theta} \right) = \left(\frac{b(\cosh \theta - 1)}{2 \sqrt{\cosh^2 \theta - 1}} \right)$ <p>Eliminate $\sqrt{\quad}$ and forms equation in x and y Simplifies to give required equation</p>	<p>As main scheme 1M1 A1ft</p> <p>cosh θ in terms of x 2M1</p> <p>y in terms of cosh θ only 3M1</p> <p>4M1 A1cso</p>

June 2012
6669 Further Pure Maths FP3
Mark Scheme

Question Number	Scheme	Marks
1. (a)	Uses formula to obtain $e = \frac{5}{4}$	M1A1
(b)	Uses ae formula	M1 (3)
	Uses other formula $\frac{a}{e}$	M1
	Obtains both Foci are $(\pm 5, 0)$ and Directrices are $x = \pm \frac{16}{5}$ (needs both method marks)	A1 cso (2) (5 marks)

Notes

a1M1: Uses $b^2 = a^2(e^2 - 1)$ to get $e > 1$

a1A1: cao

a2M1: Uses ae

b1M1: Uses $\frac{a}{e}$

b1A1: cso for both foci and both directrices. Must have both of the 2 previous M marks may be implicit.

Question Number	Scheme	Marks
2.	$\frac{dy}{dx} = \sinh 3x$ $\text{so } s = \int \sqrt{1 + \sinh^2 3x} dx$ $\therefore s = \int \cosh 3x dx$ $= \left[\frac{1}{3} \sinh 3x \right]_b^{a}$ $= \frac{1}{3} \sinh 3 \ln a = \frac{1}{6} [e^{3 \ln a} - e^{-3 \ln a}]$ $= \frac{1}{6} \left(a^3 - \frac{1}{a^3} \right) \quad (\text{so } k = 1/6)$	B1 M1 A1 M1 DM1 A1 (6 marks)

Notes

1B1: cao

1M1: Use of arc length formula, need both $\sqrt{\quad}$ and $\left(\frac{dy}{dx}\right)^2$.

1A1: $\int \cosh 3x dx$ cao

2M1: Attempt to integrate, getting a hyperbolic function o.e.

3M1: depends on previous M mark. Correct use of $\ln a$ and 0 as limits. Must see some exponentials.

2A1: cao

Question Number	Scheme	Marks
3. (a)	$\vec{AC} = 3\mathbf{i} + 6\mathbf{j} + 2\mathbf{k}, \quad \vec{BC} = -3\mathbf{i} + 4\mathbf{j} + 3\mathbf{k}$ $\vec{AC} \times \vec{BC} = 10\mathbf{i} - 15\mathbf{j} + 30\mathbf{k}$	B1, B1 M1 A1 (4)
(b)	$\text{Area of triangle } ABC = \frac{1}{2} 10\mathbf{i} - 15\mathbf{j} + 30\mathbf{k} = \frac{1}{2} \sqrt{1225} = 17.5$	M1 A1 (2)
(c)	Equation of plane is $10x - 15y + 30z = -20$ or $2x - 3y + 6z = -4$ So $\mathbf{r} \cdot (2\mathbf{i} - 3\mathbf{j} + 6\mathbf{k}) = -4$ or correct multiple	M1 A1 (2) (8 marks)

Notes

a1B1: $\vec{AC} = 3\mathbf{i} + 6\mathbf{j} + 2\mathbf{k}$ cao, any form

a2B1: $\vec{BC} = -3\mathbf{i} + 4\mathbf{j} + 3\mathbf{k}$ cao, any form

a1M1: Attempt to find cross product, modulus of one term correct.

a1A1: cao, any form.

b1M1: modulus of their answer to (a) – condone missing $\frac{1}{2}$ here. To finding area of triangle by correct method.

b1A1: cao.

c1M1: [Using their answer to (a) to] find **equation** of plane. Look for **a.n** or **b.n** or **c.n** for p.

c1A1: cao

Question Number	Scheme	Marks
4(a)	$I_n = \left[x^n \left(-\frac{1}{2} \cos 2x \right) \right]_0^{\frac{\pi}{4}} - \int_0^{\frac{\pi}{4}} -\frac{1}{2} n x^{n-1} \cos 2x dx$ <p>so</p> $I_n = \left(\left[x^n \left(-\frac{1}{2} \cos 2x \right) \right]_0^{\frac{\pi}{4}} \right) + \left[\frac{1}{4} n x^{n-1} \sin 2x \right]_0^{\frac{\pi}{4}} - \int_0^{\frac{\pi}{4}} \frac{1}{4} n(n-1) x^{n-2} \sin 2x dx$ <p>i.e. $I_n = \frac{1}{4} n \left(\frac{\pi}{4} \right)^{n-1} - \frac{1}{4} n(n-1) I_{n-2} *$</p>	<p>M1 A1</p> <p>M1 A1</p> <p>A1cso</p> <p>(5)</p>
(b)	$I_0 = \int_0^{\frac{\pi}{4}} \sin 2x dx = \left[-\frac{1}{2} \cos 2x \right]_0^{\frac{\pi}{4}} = \frac{1}{2}$ $I_2 = \frac{1}{4} \times 2 \times \left(\frac{\pi}{4} \right) - \frac{1}{4} \times 2 \times I_0, \text{ so } I_2 = \frac{\pi}{8} - \frac{1}{4}$	<p>M1 A1</p> <p>M1 A1</p> <p>(4)</p>
(c)	$I_4 = \left(\frac{\pi}{4} \right)^3 - \frac{1}{4} \times 4 \times 3 I_2 = \frac{\pi^3}{64} - 3 \left(\frac{\pi}{8} - \frac{1}{4} \right) = \frac{1}{64} (\pi^3 - 24\pi + 48) *$	<p>M1 A1cso</p> <p>(2)</p>

Notes

a1M1: Use of integration by parts, integrating $\sin 2x$, differentiating x^n .

a1A1: cao

a2M1: Second application of integration by parts, integrating $\cos 2x$, differentiating x^{n-1} .

a2A1: cao

a3A1: cso Including correct use of $\frac{\pi}{4}$ and 0 as limits.

b1M1: Integrating to find I_0 or setting up parts to find I_2 .

b1A1: cao (Accept $I_0 = \frac{1}{2}$ here for both marks)

b2M1: Finding I_2 in terms of π . If 'n's left in M0

b2A1: cao

c1M1: Finding I_4 in terms of I_2 then in terms of π . If 'n's left in M0

c1A1: cso

Question Number	Scheme	Marks
5. (a)	$\operatorname{ar sinh} 2x, +x \frac{2}{\sqrt{1+4x^2}}$	M1A1, A1 (3)
(b)	$\begin{aligned} \therefore \int_0^{\sqrt{2}} \operatorname{arsinh} 2x dx &= [x \operatorname{ar sinh} 2x]_0^{\sqrt{2}} - \int_0^{\sqrt{2}} \frac{2x}{\sqrt{1+4x^2}} dx \\ &= [x \operatorname{ar sinh} 2x]_0^{\sqrt{2}} - \left[\frac{1}{2} (1+4x^2)^{\frac{1}{2}} \right]_0^{\sqrt{2}} \\ &= \sqrt{2} \operatorname{arsinh} 2\sqrt{2} - \left[\frac{3}{2} - \frac{1}{2} \right] \\ &= \sqrt{2} \ln(3+2\sqrt{2}) - 1 \end{aligned}$	1M1 1A1ft 2M1 2A1 3DM1 4M1 3A1 (7) (10 marks)

Notes

a1M1: Differentiating getting an arsinh term **and** a term of the form $\frac{px}{\sqrt{1 \pm qx^2}}$

a1A1: cao $\operatorname{ar sinh} 2x$

a2A1: cao $+ \frac{2x}{\sqrt{1+4x^2}}$

b1M1: rearranging their answer to (a). **OR** setting up parts

b1A1: ft from their (a) **OR** setting up parts correctly

b2M1: Integrating getting an arsinh or arcosh term **and** a $(1 \pm ax^2)^{\frac{1}{2}}$ term o.e..

b2A1: cao

b3DM1: depends on previous M, correct use of $\sqrt{2}$ and 0 as limits.

b4M1: converting to log form.

b3A1: cao depends on all previous M marks.

Question Number	Scheme	Marks
6(a)	$\frac{2x}{a^2} + \frac{2y}{b^2} \frac{dy}{dx} = 0 \quad \text{and so} \quad \frac{dy}{dx} = -\frac{xb^2}{ya^2} = -\frac{b \cos \theta}{a \sin \theta}$ $\therefore y - b \sin \theta = -\frac{b \cos \theta}{a \sin \theta} (x - a \cos \theta)$ <p>Uses $\cos^2 \theta + \sin^2 \theta = 1$ to give $\frac{x \cos \theta}{a} + \frac{y \sin \theta}{b} = 1$ *</p>	<p>M1 A1</p> <p>M1</p> <p>A1cso</p> <p>(4)</p>
(b)	<p>Gradient of circle is $-\frac{\cos \theta}{\sin \theta}$ and equation of tangent is</p> $y - a \sin \theta = -\frac{\cos \theta}{\sin \theta} (x - a \cos \theta) \quad \text{or sets } a = b \text{ in previous answer}$ <p>So $y \sin \theta + x \cos \theta = a$</p>	<p>M1</p> <p>A1</p> <p>(2)</p>
(c)	<p>Eliminate x or y to give $y \sin \theta (\frac{a}{b} - 1) = 0$ or $x \cos \theta (\frac{b}{a} - 1) = b - a$</p> <p>$l_1$ and l_2 meet at $(\frac{a}{\cos \theta}, 0)$</p>	<p>M1</p> <p>A1, B1</p> <p>(3)</p>
(d)	<p>The locus of R is part of the line $y = 0$, such that $x \geq a$ and $x \leq -a$</p> <p>Or clearly labelled sketch.</p> <p>Accept "real axis"</p>	<p>B1, B1</p> <p>(2)</p> <p>(11 marks)</p>

Notes

a1M1: Finding gradient in terms of θ . Must use calculus.

a1A1: cao

a2M1: Finding equation of tangent

a2A1: cso (answer given). Need to get $\cos^2 \theta + \sin^2 \theta$ on the same side.

b1M1: Finding gradient and equation of tangent, **or** setting $a = b$.

b1A1: cao need not be simplified.

c1M1: As scheme

c1A1: $x = \frac{a}{\cos \theta}$, need not be simplified.

c1B1: $y = 0$, need not be simplified.

d1B1: Identifying locus as $y = 0$ or real/'x' axis.

d2B1: Depends on previous B mark, identifies correct parts of $y = 0$. Condone use of strict inequalities.

Question Number	Scheme	Marks
7(a)	$f(x) = 5\cosh x - 4\sinh x = 5 \times \frac{1}{2}(e^x + e^{-x}) - 4 \times \frac{1}{2}(e^x - e^{-x})$ $= \frac{1}{2}(e^x + 9e^{-x}) \quad *$	M1 A1cso (2)
(b)	$\frac{1}{2}(e^x + 9e^{-x}) = 5 \Rightarrow e^{2x} - 10e^x + 9 = 0$ <p>So $e^x = 9$ or 1 and $x = \ln 9$ or 0</p>	M1 A1 M1 A1 (4)
(c)	<p>Integral may be written $\int \frac{2e^x}{e^{2x} + 9} dx$</p> <p>This is $\frac{2}{3} \arctan\left(\frac{e^x}{3}\right)$</p> <p>Uses limits to give $\left[\frac{2}{3} \arctan 1 - \frac{2}{3} \arctan\left(\frac{1}{\sqrt{3}}\right)\right] = \left[\frac{2}{3} \times \frac{\pi}{4} - \frac{2}{3} \times \frac{\pi}{6}\right] = \frac{\pi}{18} *$</p>	B1 M1 A1 DM1 A1cso (5) (11 marks)

Notes

a1M1: Replacing both coshx and sinhx by terms in e^x and e^{-x} condone sign errors here.

a1A1: cso (answer given)

b1M1: Getting a three term quadratic in e^x

b1A1: cao

b2M1: solving to $x =$

b2A1: cao need $\ln 9$ (o.e) and 0 (not $\ln 1$)

c1B1: cao getting into suitable form, may substitute first.

c1M1: Integrating to give term in arctan

c1A1: cao

c2M1: Depends on previous M mark. Correct use of $\ln 3$ and $\frac{1}{2} \ln 3$ as limits.

c2A1: cso must see them subtracting two terms in π .

Question Number	Scheme	Marks
8. (a)	$\begin{vmatrix} 2-\lambda & 1 & 0 \\ 1 & 2-\lambda & 0 \\ -1 & 0 & 4-\lambda \end{vmatrix} = 0 \therefore (2-\lambda)(2-\lambda)(4-\lambda) - (4-\lambda) = 0$ <p>(4 - λ) = 0 verifies λ = 4 is an eigenvalue (can be seen anywhere)</p> <p>∴ (4 - λ){4 - 4λ + λ² - 1} = 0 ∴ (4 - λ){λ² - 4λ + 3} = 0</p> <p>∴ (4 - λ)(λ - 1)(λ - 3) = 0 and 3 and 1 are the other two eigenvalues</p>	<p>M1</p> <p>M1</p> <p>A1</p> <p>M1 A1</p> <p>(5)</p>
(b)	<p>Set $\begin{pmatrix} 2 & 1 & 0 \\ 1 & 2 & 0 \\ -1 & 0 & 4 \end{pmatrix} \begin{pmatrix} x \\ y \\ z \end{pmatrix} = 4 \begin{pmatrix} x \\ y \\ z \end{pmatrix}$ or $\begin{pmatrix} -2 & 1 & 0 \\ 1 & -2 & 0 \\ -1 & 0 & 0 \end{pmatrix} \begin{pmatrix} x \\ y \\ z \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \\ 0 \end{pmatrix}$</p> <p>Solve -2x + y = 0 and x - 2y = 0 and -x = 0 to obtain x = 0, y = 0, z = k</p> <p>Obtain eigenvector as k (or multiple)</p>	<p>M1</p> <p>M1</p> <p>A1 (3)</p>
(c)	<p>l_1 has equation which may be written $\begin{pmatrix} 3+\lambda \\ 2-\lambda \\ -2+2\lambda \end{pmatrix}$</p> <p>So l_2 is given by $\mathbf{r} = \begin{pmatrix} 2 & 1 & 0 \\ 1 & 2 & 0 \\ -1 & 0 & 4 \end{pmatrix} \begin{pmatrix} 3+\lambda \\ 2-\lambda \\ -2+2\lambda \end{pmatrix}$</p> <p>i.e. $\mathbf{r} = \begin{pmatrix} 8+\lambda \\ 7-\lambda \\ -11+7\lambda \end{pmatrix}$</p> <p>So $(\mathbf{r} - \mathbf{c}) \times \mathbf{d} = \mathbf{0}$ where $\mathbf{c} = 8\mathbf{i} + 7\mathbf{j} - 11\mathbf{k}$ and $\mathbf{d} = \mathbf{i} - \mathbf{j} + 7\mathbf{k}$</p>	<p>B1</p> <p>M1</p> <p>M1 A1</p> <p>A1ft (5) (13 marks)</p>

Notes

a1M1: Condone missing = 0. (They might expand the determinant using any row or column)

a2M1: Shows λ = 4 is an eigenvalue. Some working needed need to see = 0 at some stage.

a1A1: Three term quadratic factor cao, may be implicit (this A depends on 1st M only)

a2M1: Attempt at factorisation (usual rules), solving to λ = .

a2A1: cao. If they state λ = 1 and 3 please give the marks.

b1M1: Using $\mathbf{Ax} = 4\mathbf{x}$ o.e.

b2M1: Getting a pair of correct equations.

b1A1: cao

c1B1: Using **a** and **b**.

c1M1: Using $\mathbf{r} = \mathbf{M} \times$ their matrix in **a** and **b**.

c2M1: Getting an expression for l_2 with at least one component correct.

c1A1: cao all three components correct

c2A1ft: ft their vector, must have $\mathbf{r} =$ or $(\mathbf{r} - \mathbf{c}) \times \mathbf{d} = \mathbf{0}$ need both equation and r.




Mark Scheme (Results)

Summer 2013

GCE Further Pure Mathematics 3 (6669/01R)

Question Number	Scheme		Marks
	Foci (±5, 0), Directrices $x = \pm\frac{9}{5}$		
1.	$(\pm)ae = (\pm)5$ and $(\pm)\frac{a}{e} = (\pm)\frac{9}{5}$	Correct equations (ignore ±'s)	B1
	so $e = \frac{5}{a} \Rightarrow \frac{a^2}{5} = \frac{9}{5} \Rightarrow a^2 = 9$ or $a = \frac{5}{e} \Rightarrow \frac{5}{e^2} = \frac{9}{5} \Rightarrow e = \frac{5}{3} \Rightarrow a = 3$	M1: Solves using an appropriate method to find a^2 or a	M1A1
		A1: $a^2 = 9$ or $a = (\pm)3$	
	$b^2 = a^2e^2 - a^2 \Rightarrow b^2 = 25 - 9$ so $b^2 = 16 \quad (\Rightarrow b = 4)$ or $b^2 = a^2(e^2 - 1) \Rightarrow b^2 = 9\left(\frac{25}{9} - 1\right)$ $b^2 = 16 \quad (\Rightarrow b = 4)$	M1: Use of $b^2 = a^2(e^2 - 1)$ to obtain a numerical value for b^2 or b	M1 A1
	A1: $b^2 = 16$ or $b = (\pm)4$		
	So $\frac{x^2}{9} - \frac{y^2}{16} = 1$	M1: Use of $\frac{x^2}{a^2} - \frac{y^2}{b^2} = 1$ with their a^2 and b^2	M1 A1
		A1: Correct hyperbola in any form.	
		(7)	

Question Number	Scheme		Marks
3. (a)			<p>A closed curve approximately symmetrical about both axes. A vertical line to the right of the curve. A horizontal line from any point on the ellipse to the vertical line with both P and N clearly marked.</p> <p>B1 (1)</p>
3. (b)	M is $\left(\frac{x+8}{2}, y\right) = (X, Y)$ or $\left(\frac{6\cos\theta+8}{2}, 3\sin\theta\right) = (X, Y)$	M1: Finds the mid-point of PN	M1A1
	$\frac{(2X-8)^2}{36} + \frac{Y^2}{9} = 1$	M1: Attempt cartesian equation A1: Correct equation	
			(4)
The next 3 marks are dependent on having the equation of a circle.			
(c)	Circle because equation may be written $(x-4)^2 + y^2 = 3^2$	Convincing argument – allow follow through provided they do have a circle! Can be implied by their centre and radius.	B1ft
	The centre is (4, 0) and the radius is 3	M1: Use their circle equation to find centre and radius A1: Correct centre and radius	M1A1
			(3)
Total 8			
<p>Special Case: In (b) they assume the locus is a circle and find the intercepts on the x-axis as (1, 0) and (7, 0) and hence deduce the centre (4, 0) and radius 3. This approach scores no marks in (b) but allow recovery in (c).</p>			

Question Number	Scheme		Marks
4.	$\begin{pmatrix} 2 & 0 & 3 \\ 0 & 2 & -1 \\ 0 & 1 & 2 \end{pmatrix} \begin{pmatrix} 1+s+t \\ -1+s+2t \\ 2 & -2t \end{pmatrix}$	M1: Writes Π_1 as a single vector	M1A1
		A1: Correct statement	
	$\begin{pmatrix} 2 & 0 & 3 \\ 0 & 2 & -1 \\ 0 & 1 & 2 \end{pmatrix} \begin{pmatrix} 1+s+t \\ -1+s+2t \\ 2 & -2t \end{pmatrix} = \begin{pmatrix} 2+2s+2t+6-6t \\ -2+2s+4t-2+2t \\ -1+s+2t+4-4t \end{pmatrix}$		M1A1
	M1: Correct attempt to multiply A1: Correct vector in any form		
	$= \begin{pmatrix} 8+2s-4t \\ -4+2s+6t \\ 3+s-2t \end{pmatrix}$	Correct simplified vector	B1
	$\mathbf{r} = \begin{pmatrix} 8 \\ -4 \\ 3 \end{pmatrix} + s \begin{pmatrix} 2 \\ 2 \\ 1 \end{pmatrix} + t \begin{pmatrix} -4 \\ 6 \\ -2 \end{pmatrix}$		
	$\mathbf{n} = \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ 2 & 2 & 1 \\ -4 & 6 & -2 \end{vmatrix} = -10\mathbf{i} + 20\mathbf{k}$	M1: Attempts cross product of their direction vectors	M1A1
		A1: Any multiple of $-10\mathbf{i} + 20\mathbf{k}$	
	$(8\mathbf{i} - 4\mathbf{j} + 3\mathbf{k}) \cdot (\mathbf{i} - 2\mathbf{k}) = 8 - 6$	Attempt scalar product of their normal vector with their position vector	M1
	$\mathbf{r} \cdot (\mathbf{i} - 2\mathbf{k}) = 2$	Correct equation (accept any correct equivalent e.g. $\mathbf{r} \cdot (-10\mathbf{i} + 20\mathbf{k}) = -20$)	A1
			(9)

Question Number	Scheme		Marks
5(a)	$I_n = \left[x^n (2x-1)^{\frac{1}{2}} \right]_1^5 - \int_1^5 nx^{n-1} (2x-1)^{\frac{1}{2}} dx$	M1: Parts in the correct direction including a valid attempt to integrate $(2x-1)^{-\frac{1}{2}}$ A1: Fully correct application – may be un-simplified. (Ignore limits)	M1 A1
	$I_n = \underline{5^n \times 3 - 1} - \int_1^5 nx^{n-1} \underline{(2x-1)(2x-1)^{-\frac{1}{2}}} dx$	Obtains a correct (possibly un-simplified) expression using the limits 5 and 1 and writes $(2x-1)^{\frac{1}{2}}$ as $(2x-1)(2x-1)^{-\frac{1}{2}}$	B1
	$I_n = 5^n \times 3 - 1 - 2nI_n + nI_{n-1}$	Replaces $\int x^n (2x-1)^{-\frac{1}{2}} dx$ with I_n and $\int x^{n-1} (2x-1)^{-\frac{1}{2}} dx$ with I_{n-1}	dM1
	$(2n+1)I_n = nI_{n-1} + 3 \times 5^n - 1 *$	Correct completion to printed answer with no errors seen	A1cso
			(5)
(b)	$I_0 = \int_1^5 (2x-1)^{-\frac{1}{2}} dx = \left[(2x-1)^{\frac{1}{2}} \right]_1^5 = 2$	$I_0 = 2$	B1
	$5I_2 = 2I_1 + 74 \text{ and } 3I_1 = I_0 + 14$	M1: Correctly applies the given reduction formula twice A1: Correct <u>equations</u> for I_2 and I_1 (may be implied)	M1 A1
	$\text{So } I_1 = \frac{16}{3} \text{ and } I_2 = \dots \text{ or}$ $5I_2 = 2 \frac{I_0 + 14}{3} + 74 \text{ and } I_2 = \dots$	Completes to obtains a numerical expression for I_2	dM1
	$I_2 = \frac{254}{15}$		B1
			Total 10

Question Number	Scheme		Marks
6. (a)	$\begin{pmatrix} 4 & 2 & 3 \\ 2 & b & 0 \\ a & 1 & 8 \end{pmatrix} \begin{pmatrix} 1 \\ 2 \\ 0 \end{pmatrix} = \begin{pmatrix} 8 \\ \dots \\ \dots \end{pmatrix}, = \lambda \begin{pmatrix} 1 \\ 2 \\ 0 \end{pmatrix}, \lambda = 8$	M1: Multiplies the given matrix by the given eigenvector M1: Equates the x value to λ A1: $\lambda = 8$	M1, M1, A1
			(3)
(b)	$\begin{pmatrix} 8 \\ 2+2b \\ a+2 \end{pmatrix} = "8" \begin{pmatrix} 1 \\ 2 \\ 0 \end{pmatrix} \text{ So } a = -2 \text{ and } b = 7$	M1: Their $2 + 2b = 2\lambda$ or their $a + 2 = 0$ A1: $b = 7$ or $a = -2$ A1: $b = 7$ and $a = -2$	M1 A1 A1
			(3)
(c)	$\begin{vmatrix} 4-\lambda & 2 & 3 \\ 2 & 7-\lambda & 0 \\ -2 & 1 & 8-\lambda \end{vmatrix} = 0$ $\therefore (4-\lambda)(7-\lambda)(8-\lambda) - 2 \times 2(8-\lambda) + 3(2+2(7-\lambda)) = 0$		M1
Correct attempt to establish the Characteristic Equation. = 0 is required but may be implied by later work Allow this mark if the equation is in terms of a, b, c			
Attempts to factorise i.e. $(8-\lambda)(30-11\lambda+\lambda^2)$ or $(6-\lambda)(40-13\lambda+\lambda^2)$ or $(5-\lambda)(48-14\lambda+\lambda^2)$ (NB $240-118\lambda+19\lambda^2-\lambda^3=0$)			M1 A1
M1: Attempt to factorise their cubic – an attempt to identify a linear factor and processes to obtain a simplified quadratic factor A1: Correct factorisation into one linear and one quadratic factor			
Eigenvalues are 5 and 6		M1: Solves their equation to obtain the other eigenvalues A1: 5 and 6	M1 A1
			(5)
			Total 8

Question Number	Scheme		Marks
7(a)	Put $6\cosh x = 9 - 2\sinh x$		M1
	$6 \times \frac{1}{2}(e^x + e^{-x}) = 9 - 2 \times \frac{1}{2}(e^x - e^{-x})$	Replaces $\cosh x$ and $\sinh x$ by the correct exponential forms	M1
	$4e^x + 2e^{-x} - 9 = 0 \Rightarrow 4e^{2x} - 9e^x + 2 = 0$	M1: Multiplies by e^x A1: Correct quadratic in e^x in any form with terms collected	M1 A1
	So $e^x = \frac{1}{4}$ or 2 and $x = \ln 2$ or $\ln \frac{1}{4}$	M1: Solves their quadratic in e^x A1: Correct values of x (Any correct equivalent form)	M1 A1
(b)	Area is $\int (9 - 2\sinh x - 6\cosh x) dx$	$\int (9 - 2\sinh x - 6\cosh x) dx$ or $\int (6\cosh x - (9 - 2\sinh x)) dx$ or the equivalent in exponential form	M1
	$\pm(9x - 2\cosh x - 6\sinh x)$ or $\pm(9x - 4e^x + 2e^{-x})$	M1: Attempt to integrate A1: Correct integration	M1 A1
	$\pm\left(9\ln 2 - 2\cosh \ln 2 - 6\sinh \ln 2\right) - \left(9\ln \frac{1}{4} - 2\cosh \ln \frac{1}{4} - 6\sinh \ln \frac{1}{4}\right)$		dM1
	Complete substitution of their limits from part (a). Depends on both previous M's		
	$= \pm\left(9\ln\left(2 \div \frac{1}{4}\right) - (e^{\ln 2} + e^{-\ln 2}) - 3(e^{\ln 2} - e^{-\ln 2}) + (e^{\ln \frac{1}{4}} + e^{-\ln \frac{1}{4}}) + 3(e^{\ln \frac{1}{4}} - e^{-\ln \frac{1}{4}})\right)$		M1
	Combines logs correctly and uses cosh and sinh of ln correctly at least once		
	$\left(9\ln 8 - \frac{5}{2} - \frac{18}{4} + 4.25 - 11.25\right) = 9\ln 8 - 14$ or $27\ln 2 - 14$ Any correct equivalent		A1cao
Subtracting the wrong way round could score 5/6 max			
			(6)
			Total 12
<u>Note</u> If they use $4e^{2x} - 9e^x + 2$ in (b) to find the area – no marks			

Question Number	Scheme		Marks
8(a)	$\frac{dy}{dx} = x^{-\frac{1}{2}}$	Correct derivative (may be unsimplified)	B1
	$s = \int \sqrt{1+(x^{-\frac{1}{2}})^2} dx = \int_1^8 \sqrt{1+\frac{1}{x}} dx$	A correct formula quoted or implied. There must be some working before the printed answer.	B1
			(2)
(b)	$x = \sinh^2 u \Rightarrow \frac{dx}{du} = 2 \sinh u \cosh u$	Correct derivative	B1
	$(1 + \frac{1}{x}) = 1 + \operatorname{cosech}^2 u = \coth^2 u$	$(1 + \frac{1}{x}) = \coth^2 u$ or $(1 + \frac{1}{x}) = \frac{\cosh^2 u}{\sinh^2 u}$ (may be implied by later work)	B1
	$s = \int \coth u \cdot 2 \sinh u \cosh u du = \int 2 \cosh^2 u du$	M1: Complete substitution A1: $\int 2 \cosh^2 u du$	M1 A1
	$= u + \frac{1}{2} \sinh 2u$ or $\frac{1}{4} e^{2u} + u - \frac{1}{4} e^{-2u}$	M1: Uses $\cosh 2u = \pm 2 \cosh^2 u \pm 1$ or changes to exponentials in an attempt to integrate an expression of the form $k \cosh^2 u$ A1: Correct integration	dM1 A1
	$x = 8 \Rightarrow u = \operatorname{arsinh} \sqrt{8} = \ln(3 + 2\sqrt{2}), x = 1 \Rightarrow u = \operatorname{arsinh} 1 = \ln(1 + \sqrt{2})$		
	$\left[u + \frac{1}{2} \sinh 2u \right]_{\operatorname{arsinh} 1}^{\operatorname{arsinh} \sqrt{8}}$ $= \operatorname{arsinh} \sqrt{8} + \frac{1}{2} \sinh(2 \operatorname{arsinh} \sqrt{8}) - (\operatorname{arsinh} 1 + \frac{1}{2} \sinh(2 \operatorname{arsinh} 1))$ <p>or</p> $\left[\frac{1}{4} e^{2u} + u - \frac{1}{4} e^{-2u} \right]_{\operatorname{arsinh} 1}^{\operatorname{arsinh} \sqrt{8}}$ $= \frac{1}{4} e^{\operatorname{arsinh} \sqrt{8}} + \operatorname{arsinh} \sqrt{8} - \frac{1}{4} e^{-2 \operatorname{arsinh} 1}$ <p>or</p> $\left[\operatorname{arsinh} \sqrt{x} + \frac{1}{2} \sinh(2 \operatorname{arsinh} \sqrt{x}) \right]_1^8$ $= \operatorname{arsinh} \sqrt{8} + \frac{1}{2} \sinh(2 \operatorname{arsinh} \sqrt{8}) - (\operatorname{arsinh} 1 + \frac{1}{2} \sinh(2 \operatorname{arsinh} 1))$		ddM1A1
	M1: The limits $\operatorname{arsinh} \sqrt{8}$ and $\operatorname{arsinh} 1$ or their $\ln(3 + 2\sqrt{2})$ and $\ln(1 + \sqrt{2})$ used correctly in their $f(u)$ or the limits 8 and 1 used correctly if they revert to x Dependent on both previous M's A1: A completely correct expression		
	$\ln(1 + \sqrt{2}) + 5\sqrt{2}$		A1
			(9)
			Total 11



Mark Scheme (Results)

Summer 2013

GCE Further Pure Mathematics 3 (6669/01)

Question Number	Scheme		Marks	
Mark (a) and (b) together				
1. (a) & (b)	$ae = 13$ and $a^2(e^2 - 1) = 25$	Sight of both of these (can be implied by their work) (allow $\pm ae = \pm 13$ or $\pm ae = 13$ or $ae = \pm 13$)	B1	
	Solves to obtain $a^2 = \dots$ or $a = \dots$	Eliminates e to reach $a^2 = \dots$ or $a = \dots$	M1	
	$a = 12$	Cao (not ± 12) unless -12 is rejected	A1	
	$e = 13/ "12"$	Uses their a to find e or finds e by eliminating a (Ignore \pm here) (Can be implied by a correct answer)	M1	
	$x = (\pm) \frac{a}{e}, = \pm \frac{144}{13}$	M1: $(x =)(\pm) \frac{a}{e}$ \pm not needed for this mark nor is x and even allow $y = (\pm) \frac{a}{e}$ here – just look for use of $\frac{a}{e}$ with numerical a and e . A1: $x = \pm \frac{144}{13}$ oe but must be an <u>equation</u> (Do not allow $x = \pm \frac{12}{13/12}$)	M1, A1	
			Total 6	
	If they use the eccentricity equation for the ellipse ($b^2 = a^2(1 - e^2)$) allow the M's			

Question Number	Scheme	Marks
2. (a)	$k \operatorname{arsinh}\left(\frac{2x}{3}\right) (+c)$ or $k \ln\left[px + \sqrt{(p^2x^2 + \frac{9}{4}p^2)}\right] (+c)$	M1
	$\frac{1}{2} \operatorname{ar sinh}\left(\frac{2x}{3}\right) (+c)$ or $\frac{1}{2} \ln\left[px + \sqrt{(p^2x^2 + \frac{9}{4}p^2)}\right] (+c)$	A1
		(2)
(b)	So: $\frac{1}{2} \ln[6 + \sqrt{45}] - \frac{1}{2} \ln[-6 + \sqrt{45}] = \frac{1}{2} \ln\left[\frac{6 + \sqrt{45}}{-6 + \sqrt{45}}\right]$	M1
	Uses correct limits <u>and</u> combines logs	
	$= \frac{1}{2} \ln\left[\frac{6 + \sqrt{45}}{-6 + \sqrt{45}}\right] \left[\frac{6 + \sqrt{45}}{6 + \sqrt{45}}\right] = \frac{1}{2} \ln\left[\frac{(6 + \sqrt{45})^2}{9}\right]$	M1
	Correct method to rationalise denominator (may be implied) Method must be clear if answer does not follow their fraction	
	$= \ln[2 + \sqrt{5}]$ (or $\frac{1}{2} \ln[9 + 4\sqrt{5}]$)	A1 cso
	Note that the last 3 marks can be scored without the need to rationalise e.g. $2 \times \frac{1}{2} \left[\ln[2x + \sqrt{(4x^2 + 9)}] \right]_0^3 = \ln(6 + \sqrt{45}) - \ln 3 = \ln\left(\frac{6 + \sqrt{45}}{3}\right)$ M1: Uses the limits 0 and 3 and doubles M1: Combines Logs A1: $\ln[2 + \sqrt{5}]$ oe	
		(3)
Total 5		
Alternative for (a)	$x = \frac{3}{2} \sinh u \Rightarrow \int \frac{1}{\sqrt{9 \sinh^2 u + 9}} \cdot \frac{3}{2} \cosh u \, du = k \operatorname{arsinh}\left(\frac{2x}{3}\right) (+c)$	M1
	$\frac{1}{2} \operatorname{ar sinh}\left(\frac{2x}{3}\right) (+c)$	A1
Alternative for (b)	$\left[\frac{1}{2} \operatorname{arsinh}\left(\frac{2x}{3}\right)\right]_{-3}^3 = \frac{1}{2} \operatorname{arsinh} 2 - \frac{1}{2} \operatorname{arsinh} -2$	
	$\frac{1}{2} \ln(2 + \sqrt{5}) - \frac{1}{2} \ln(\sqrt{5} - 2) = \frac{1}{2} \ln\left(\frac{2 + \sqrt{5}}{\sqrt{5} - 2}\right)$	M1
	Uses correct limits <u>and</u> combines logs	
	$= \frac{1}{2} \ln\left(\frac{2 + \sqrt{5}}{\sqrt{5} - 2} \cdot \frac{\sqrt{5} + 2}{\sqrt{5} + 2}\right) = \frac{1}{2} \ln\left(\frac{2\sqrt{5} + 4 + 5 + 2\sqrt{5}}{5 - 4}\right)$	M1
	Correct method to rationalise denominator (may be implied) Method must be clear if answer does not follow their fraction	
	$= \frac{1}{2} \ln[9 + 4\sqrt{5}]$	A1 cso

Question Number	Scheme	Marks
3.	$\left(\frac{dx}{d\theta}\right) = 2 \sinh 2\theta \quad \text{and} \quad \left(\frac{dy}{d\theta}\right) = 4 \cosh \theta$ <p>Or equivalent correct derivatives</p>	B1
	$A = (2\pi) \int 4 \sinh \theta \sqrt{2 \sinh^2 \theta + 4 \cosh^2 \theta} d\theta$ <p>or</p> $A = (2\pi) \int 4 \sinh \theta \sqrt{\left(1 + \frac{4 \cosh^2 \theta}{2 \sinh^2 \theta}\right)^2} \cdot 2 \sinh 2\theta d\theta$	M1
	<p>Use of correct formula including replacing dx with "2 sinh 2θ" dθ if chain rule used. Allow the omission of the 2π here.</p>	
	$A = 32\pi \int \sinh \theta \cosh^2 \theta d\theta$ $A = 32\pi \int (\sinh \theta + \sinh^3 \theta) d\theta$	B1
	<p>Completely correct expression for A with the square root removed This mark may be recovered later if the 2π is introduced later</p>	
	$A = \frac{32\pi}{3} [\cosh^3 \theta]_0^1$	<p>M1: Valid attempt to integrate a correct expression or a multiple of a correct expression – dependent on the first M1</p> <p>A1: Correct expression</p>
	$= \frac{32\pi}{3} [\cosh^3 1 - 1]$	<p>M1: Uses the limits 0 and 1 correctly. Dependent on both previous M's</p> <p>A1: Cao and cso (no errors seen)</p>
		(7)
	<p>Example Alternative Integration for last 4 marks</p>	
	$\int \sinh \theta \cosh^2 \theta d\theta = \int \sinh \theta (1 + \sinh^2 \theta) d\theta = \int (\sinh \theta + \sinh^3 \theta) d\theta$ $\int \left(\sinh \theta + \frac{1}{4} \sinh 3\theta - \frac{3}{4} \sinh \theta\right) d\theta = \frac{1}{4} \int (\sinh \theta + \sinh 3\theta) d\theta$ $= \frac{1}{4} \cosh \theta + \frac{1}{12} \cosh 3\theta$ <p>dM1: $\int \sinh \theta \cosh^2 \theta d\theta = p \cosh \theta + q \cosh 3\theta$</p> <p>A1: $32\pi \left[\frac{1}{4} \cosh \theta + \frac{1}{12} \cosh 3\theta \right]$</p>	dM1A1
	$A = 8\pi \left[\cosh \theta + \frac{1}{3} \cosh 3\theta \right]_0^1$ $= 8\pi \left(\cosh 1 + \frac{1}{3} \cosh 3 - \cosh 0 - \frac{1}{3} \cosh 0 \right)$ <p>.....</p> $\frac{32\pi}{3} [\cosh^3 1 - 1]$	<p>M1: Uses the limits 0 and 1 correctly. Dependent on both previous M's</p> <p>A1: Cao</p>

Question Number	Scheme		Marks
3.	Alternative Cartesian Approach		
	$x = 1 + \frac{y^2}{8}$	Any correct Cartesian equation	B1
	$\frac{dx}{dy} = \frac{y}{4}$ or $\frac{dy}{dx} = \frac{\sqrt{2}}{(x-1)^{\frac{1}{2}}}$	Correct Derivative	B1
	$A = \int 2\pi \cdot y \sqrt{1 + \left(\frac{y}{4}\right)^2} dy$ or $A = \int 2\pi \cdot \sqrt{8}(x-1)^{\frac{1}{2}} \sqrt{1 + \left(\frac{2}{x-1}\right)} dx$		M1
	Use of a correct formula		
	$A = 2\pi \times \frac{2}{3} \times 8 \left(1 + \frac{y^2}{16}\right)^{\frac{3}{2}}$ or $A = \frac{4\pi\sqrt{8}}{3} x + 1^{\frac{3}{2}}$		dM1 A1
	M1: Convincing attempt to integrate a relevant expression – dependent on the first M1 but allow the omission of 2π		
	A1: Completely correct expression for A		
	$A = 2\pi \times \frac{2}{3} \times 8 \left[1 + \sinh^2 1\right]^{\frac{3}{2}} - 2\pi \times \frac{2}{3} \times 8$ or $2\pi \times \frac{2}{3} \times \sqrt{8} \left[1 + \cosh 2\right]^{\frac{3}{2}} - \frac{32\pi}{3}$		ddM1
	Correct use of limits (0 → 4sinh1 for y or 1 → cosh2 for x)		
	Use $1 + \sinh^2 1 = \cosh^2 1$ to give $\frac{32\pi}{3} [\cosh^3 1 - 1]$	Use $\cosh 2 = 2 \cosh^2 1 - 1$ to give $\frac{32\pi}{3} [\cosh^3 1 - 1]$	A1

Question Number	Scheme		Marks
4.	$\frac{dy}{dx} = \frac{40}{\sqrt{(x^2 - 1)}} - 9$	M1: $\frac{dy}{dx} = \frac{p}{\sqrt{(x^2 - 1)}} - q$	M1 A1
		A1: Cao	
	Put $\frac{dy}{dx} = 0$ and obtain $x^2 = \dots$ (Allow sign errors only)	e.g. $\left(\frac{1681}{81}\right)$	dM1
	$x = \frac{41}{9}$	M1: Square root	M1 A1
		A1: $x = \frac{41}{9}$ or exact equivalent (not $\pm \frac{41}{9}$)	
$y = 40 \ln \left\{ \left(\frac{41}{9} \right) + \sqrt{\left(\frac{41}{9} \right)^2 - 1} \right\} - 41$	Substitutes $x = \frac{41}{9}$ into the curve and uses the logarithmic form of arcosh	M1	
So $y = 80 \ln 3 - 41$	Cao	A1	
		Total 7	

Question Number	Scheme	Marks
5. (a) (i)&(ii)	$\begin{pmatrix} 1 & 1 & a \\ 2 & b & c \\ -1 & 0 & 1 \end{pmatrix} \begin{pmatrix} 0 \\ 1 \\ 1 \end{pmatrix} = \begin{pmatrix} 1+a \\ b+c \\ 1 \end{pmatrix} = \lambda_1 \begin{pmatrix} 0 \\ 1 \\ 1 \end{pmatrix}, \text{ and so } a = -1, \lambda_1 = 1$	M1, A1, A1
	<p>M1: Multiplies out matrix with first eigenvector and puts equal to λ_1 times eigenvector. A1 : Deduces $a = -1$. A1: Deduces $\lambda_1 = 1$</p>	
	$\begin{pmatrix} 1 & 1 & a \\ 2 & b & c \\ -1 & 0 & 1 \end{pmatrix} \begin{pmatrix} 1 \\ 0 \\ -1 \end{pmatrix} = \begin{pmatrix} 1-a \\ 2-c \\ -2 \end{pmatrix} = \lambda_2 \begin{pmatrix} 1 \\ 0 \\ -1 \end{pmatrix}, \text{ and so } c = 2, \lambda_2 = 2$	M1, A1, A1
	<p>M1: Multiplies out matrix with second eigenvector and puts equal to λ_2 times eigenvector. A1: Deduces $c = 2$. A1: Deduces $\lambda_2 = 2$</p>	
	$b + c = \lambda_1 \quad \text{so } b = -1$	<p>M1: Uses $b + c = \lambda_1$ with their λ_1 to find a value for b (They must have an equation in b and c from the first eigenvector to score this mark) A1: $b = -1$</p>
	$(a = -1, b = -1, c = 2, \lambda_1 = 1, \lambda_2 = 2)$	(8)
(b)(i)	$\det P = -d - 1$	<p>Allow $1 - d - 2$ or $1 - (2 + d)$ A correct (possibly un-simplified) determinant</p>
(ii)	$P^T = \begin{pmatrix} 1 & 2 & -1 \\ 1 & 1 & 0 \\ 0 & d & 1 \end{pmatrix} \text{ or minors } \begin{pmatrix} 1 & d+2 & 1 \\ 1 & 1 & 1 \\ d & d & -1 \end{pmatrix} \text{ or}$ $\text{cofactors } \begin{pmatrix} 1 & -2-d & 1 \\ -1 & 1 & -1 \\ d & -d & -1 \end{pmatrix} \text{ a correct first step}$	B1
	$\frac{1}{-d-1} \begin{pmatrix} 1 & -1 & d \\ -2-d & 1 & -d \\ 1 & -1 & -1 \end{pmatrix}$	<p>M1: Identifiable full attempt at inverse including reciprocal of determinant. Could be indicated by at least 6 correct elements. A1: Two rows or two columns correct (ignoring determinant) BUT M0A1A0 or M0A1A1 is not possible A1: Fully correct inverse</p>
		(5)
		Total 13

Question Number	Scheme		Marks
6(a)	$I_n = \int_0^4 x^{n-1} \times x(16-x^2)^{\frac{1}{2}} dx$	M1: Obtains $x(16-x^2)^{\frac{1}{2}}$ prior to integration	M1A1
		A1: Correct underlined expression (can be implied by their integration)	
	$I_n = \left[-\frac{1}{3} x^{n-1} (16-x^2)^{\frac{3}{2}} \right]_0^4 + \frac{n-1}{3} \int_0^4 x^{n-2} (16-x^2)^{\frac{3}{2}} dx$		dM1
	dM1: Parts in the correct direction (Ignore limits)		
	$\therefore I_n = \frac{n-1}{3} \int_0^4 x^{n-2} (16-x^2)(16-x^2)^{\frac{1}{2}} dx$		
	i.e. $I_n = \frac{16(n-1)}{3} I_{n-2} - \frac{n-1}{3} I_n$	Manipulates to obtain at least one integral in terms of I_n or I_{n-2} on the rhs.	M1
	$I_n \left(1 + \frac{n-1}{3}\right) = \frac{16(n-1)}{3} I_{n-2}$	Collects terms in I_n from both sides	M1
	$(n+2)I_n = 16(n-1)I_{n-2} *$	Printed answer with no errors	A1*cso
			(6)
Way 2	$\int_0^4 x^n (16-x^2)^{\frac{1}{2}} dx = \int_0^4 x^n \frac{(16-x^2)}{(16-x^2)^{\frac{1}{2}}} dx = \int_0^4 \frac{16x^n}{(16-x^2)^{\frac{1}{2}}} dx - \int_0^4 \frac{x^{n+2}}{(16-x^2)^{\frac{1}{2}}} dx$		
	$= \int_0^4 16x^{n-1} \times x(16-x^2)^{-\frac{1}{2}} dx - \int_0^4 x^{n+1} \times x(16-x^2)^{-\frac{1}{2}} dx$		M1A1
	M1: Obtains $x(16-x^2)^{-\frac{1}{2}}$ prior to integration A1: Correct expressions		
	$= \left[-16x^{n-1} (16-x^2)^{\frac{1}{2}} \right]_0^4 + 16(n-1) \int_0^4 x^{n-2} (16-x^2)^{\frac{1}{2}} dx$ $- \left[-x^{n+1} (16-x^2)^{\frac{1}{2}} \right]_0^4 + (n+1) \int_0^4 x^n (16-x^2)^{\frac{1}{2}} dx$		dM1
	dM1: Parts in the correct direction on both (Ignore limits)		
	$I_n = 16(n-1)I_{n-2} - (n+1)I_n$	Manipulates to obtain at least one integral in terms of I_n or I_{n-2} on the rhs.	M1
	$I_n (1+n+1) = 16(n-1)I_{n-2}$	Collects terms in I_n from both sides	M1
	$(n+2)I_n = 16(n-1)I_{n-2} *$	Printed answer with no errors	A1*
Way 3	$\int_0^4 x^n (16-x^2)^{\frac{1}{2}} dx = \int_0^4 x \times x^{n-1} \frac{(16-x^2)}{(16-x^2)^{\frac{1}{2}}} dx$	M1: Obtains $x(16-x^2)^{-\frac{1}{2}}$ prior to integration	M1A1
		A1: Correct expression	
	$= \left[-x^{n-1} (16-x^2)(16-x^2)^{\frac{1}{2}} \right]_0^4 + \int_0^4 (16(n-1)x^{n-2} - (n+1)x^n)(16-x^2)^{\frac{1}{2}} dx$		dM1
	dM1: Parts in the correct direction (Ignore limits)		
	$I_n = 16(n-1)I_{n-2} - (n+1)I_n$	Manipulates to obtain at least one integral in terms of I_n or I_{n-2} on the rhs.	M1
	$I_n (1+n+1) = 16(n-1)I_{n-2}$	Collects terms in I_n from both sides	M1
	$(n+2)I_n = 16(n-1)I_{n-2} *$	Printed answer with no errors	A1*

Question Number	Scheme		Marks
(b)	$I_1 = \int_0^4 x\sqrt{(16-x^2)}dx = \left[-\frac{1}{3}(16-x^2)^{\frac{3}{2}}\right]_0^4 = \frac{64}{3}$	M1: Correct integration to find I_1	M1 A1
		A1: $\frac{64}{3}$ or equivalent (May be implied by a later work – they are not asked explicitly for I_1)	
	$\frac{64}{3}$ must come from correct work		
	Using $x = 4\sin\theta$: $I_1 = \int_0^{\frac{\pi}{2}} 4\sin\theta\sqrt{(16-16\sin^2\theta)}4\cos\theta d\theta = \int_0^{\frac{\pi}{2}} 64\sin\theta\cos^2\theta d\theta$ $= \left[-\frac{64}{3}\cos^3\theta\right]_0^{\frac{\pi}{2}}$ M1: A <u>complete</u> substitution and attempt to substitute <u>changed</u> limits A1: $\frac{64}{3}$ or equivalent		
	$I_5 = \frac{64}{7}I_3, I_3 = \frac{32}{5}I_1$	Applies to apply reduction formula twice. First M1 for I_5 in terms of I_3 , second M1 for I_3 in terms of I_1 (Can be implied)	M1, M1
$I_5 = \frac{131072}{105}$	Any <u>exact</u> equivalent (Depends on all previous marks having been scored)	A1	
		(5)	
			Total 11

Question Number	Scheme		Marks
7(a)	$(\frac{dx}{d\theta} = -a \sin \theta \text{ and } \frac{dy}{d\theta} = b \cos \theta) \text{ so } \frac{dy}{dx} = \frac{b \cos \theta}{-a \sin \theta}$		M1 A1
	M1: Differentiates both x and y and divides correctly A1: Fully correct derivative		
	Alternative: M1: $\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1 \Rightarrow \frac{2x}{a^2} + \frac{2yy'}{b^2} = 0 \Rightarrow y' = -\frac{b^2 x}{a^2 y} = -\frac{b^2 a \cos \theta}{a^2 b \sin \theta}$ Differentiates implicitly and substitutes for x and y A1: $= -\frac{b \cos \theta}{a \sin \theta}$		
	Normal has gradient $\frac{a \sin \theta}{b \cos \theta}$ or $\frac{a^2 y}{b^2 x}$	Correct perpendicular gradient rule	M1
	$(y - b \sin \theta) = \frac{a \sin \theta}{b \cos \theta} (x - a \cos \theta)$	Correct straight line method using a „changed“ gradient which is a function of θ	M1
	If $y = mx + c$ is used need to find c for M1		
	$ax \sin \theta - by \cos \theta = (a^2 - b^2) \sin \theta \cos \theta$ *		A1
	Fully correct completion to printed answer		
			(5)
(b)	$x = \frac{(a^2 - b^2) \cos \theta}{a}$	Allow un-simplified	B1
	$y = -\frac{(a^2 - b^2) \sin \theta}{b}$	Allow un-simplified	B1
	$\left(= \frac{1}{2} \frac{(a^2 - b^2)^2 \cos \theta \sin \theta}{ab} \right) = \frac{1}{4} \frac{(a^2 - b^2)^2}{ab} \sin 2\theta$		M1A1
	M1: Area of triangle is $\frac{1}{2}$ "OA" x "OB" and uses double angle formula correctly A1: Correct expression for the area (must be positive)		
			(4)
(c)	Maximum area when $\sin 2\theta = 1$ so $\theta = \frac{\pi}{4}$ or 45	Correct value for θ (may be implied by correct coordinates)	B1
	So the point P is at $\left(\frac{a}{\sqrt{2}}, \frac{b}{\sqrt{2}} \right)$ oe $\left(a \cos \frac{\pi}{4}, b \sin \frac{\pi}{4} \right)$ scores B1M1A0	M1: Substitutes their value of θ where $0 < \theta < \frac{\pi}{2}$ or $0 < \theta < 90$ into their parametric coordinates A1: Correct exact coordinates	M1 A1
	Mark part (c) independently		
			(3)
			Total 12

Question Number	Scheme		Marks
8(a)	$(6\mathbf{i} + 2\mathbf{j} + 12\mathbf{k}) \cdot (3\mathbf{i} - 4\mathbf{j} + 2\mathbf{k}) = 34$	Attempt scalar product	M1
	$\frac{ (6\mathbf{i} + 2\mathbf{j} + 12\mathbf{k}) \cdot (3\mathbf{i} - 4\mathbf{j} + 2\mathbf{k}) - 5 }{\sqrt{3^2 + 4^2 + 2^2}}$	Use of correct formula	M1
	$\sqrt{29}$ (not $-\sqrt{29}$)	Correct distance (Allow $29/\sqrt{29}$)	A1
			(3)
(a) Way 2	$\mathbf{r} = (6\mathbf{i} + 2\mathbf{j} + 12\mathbf{k}) + \lambda(3\mathbf{i} - 4\mathbf{j} + 2\mathbf{k})$ $\therefore 6 + 3\lambda \quad 2 - 4\lambda \quad 12 + 2\lambda = 5$		M1
	Substitutes the parametric coordinates of the line through (6, 2, 12) perpendicular to the plane into the cartesian equation.		
	$\lambda = -1 \Rightarrow 3, 6, 10$ or $-3\mathbf{i} + 4\mathbf{j} - 2\mathbf{k}$	Solves for λ to obtain the required point or vector.	M1
	$\sqrt{29}$	Correct distance	A1
(a) Way 3	Parallel plane containing (6, 2, 12) is $\mathbf{r} \cdot (3\mathbf{i} - 4\mathbf{j} + 2\mathbf{k}) = 34$ $\Rightarrow \frac{\mathbf{r} \cdot (3\mathbf{i} - 4\mathbf{j} + 2\mathbf{k})}{\sqrt{29}} = \frac{34}{\sqrt{29}}$	Origin to this plane is $\frac{34}{\sqrt{29}}$	M1
	$\Rightarrow \frac{\mathbf{r} \cdot (3\mathbf{i} - 4\mathbf{j} + 2\mathbf{k})}{\sqrt{29}} = \frac{5}{\sqrt{29}}$	Origin to plane is $\frac{5}{\sqrt{29}}$	M1
	$\frac{34}{\sqrt{29}} - \frac{5}{\sqrt{29}} = \sqrt{29}$	Correct distance	A1
(b) For a cross product, if the method is unclear, 2 out of 3 components should be correct for M1	$\begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ 2 & 1 & 5 \\ 1 & -1 & -2 \end{vmatrix} = \begin{pmatrix} 3 \\ 9 \\ -3 \end{pmatrix}$	M1: Attempts $(2\mathbf{i} + 1\mathbf{j} + 5\mathbf{k}) \times (\mathbf{i} - \mathbf{j} - 2\mathbf{k})$ A1: Any multiple of $\mathbf{i} + 3\mathbf{j} - \mathbf{k}$	M1A1
	$(\cos \theta) = \frac{(3\mathbf{i} - 4\mathbf{j} + 2\mathbf{k}) \cdot (\mathbf{i} + 3\mathbf{j} - \mathbf{k})}{\sqrt{3^2 + 4^2 + 2^2} \sqrt{1^2 + 3^2 + 1^2}} \left(= \frac{-11}{\sqrt{29}\sqrt{11}} \right)$		M1
	Attempts scalar product of normal vectors including magnitudes		
	52	Obtains angle using arccos (dependent on previous M1)	dM1 A1
	Do not isw and mark the final answer e.g. $90 - 52 = 38$ loses the A1		(5)
(c)	$\begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ 1 & 3 & -1 \\ 3 & -4 & 2 \end{vmatrix} = \begin{pmatrix} 2 \\ -5 \\ -13 \end{pmatrix}$	M1: Attempt cross product of normal vectors A1: Correct vector	M1A1
	$x = 0: (0, \frac{5}{2}, \frac{15}{2}), y = 0: (1, 0, 1), z = 0: (\frac{15}{13}, \frac{-5}{13}, 0)$		M1A1
M1: Valid attempt at a point on both planes. A1: Correct coordinates May use way 3 to find a point on the line			
	$\mathbf{r} \times (-2\mathbf{i} + 5\mathbf{j} + 13\mathbf{k}) = -5\mathbf{i} - 15\mathbf{j} + 5\mathbf{k}$	M1: $\mathbf{r} \times \text{dir} = \text{pos. vector} \times \text{dir}$ (This way round) A1: Correct equation	M1A1
			(6)

Question Number	Scheme	Marks	
(c) Way 2	“ $x + 3y - z = 0$ ” and $3x - 4y + 2z = 5$ uses their cartesian form of and eliminate x , or y or z and substitutes back to obtain two of the variables in terms of the third	M1	
	$(x = 1 - \frac{2}{5}y \text{ and } z = 1 + \frac{13}{5}y) \text{ or } (y = \frac{5z-5}{13} \text{ and } x = \frac{15-2z}{13}) \text{ or}$ $(y = \frac{5-5x}{2} \text{ and } z = \frac{15-13x}{2})$	A1	
	Cartesian Equations: $x = \frac{y - \frac{5}{2}}{-\frac{5}{2}} = \frac{z - \frac{15}{2}}{-\frac{13}{2}} \text{ or } \frac{x-1}{-\frac{2}{5}} = y = \frac{z-1}{\frac{13}{5}} \text{ or } \frac{x - \frac{15}{13}}{-\frac{2}{13}} = \frac{y + \frac{5}{13}}{\frac{5}{13}} = z$		
	Points and Directions: Direction can be any multiple $(0, \frac{5}{2}, \frac{15}{2}), \mathbf{i} - \frac{5}{2}\mathbf{j} - \frac{13}{2}\mathbf{k}$ or $(1, 0, 1), -\frac{2}{5}\mathbf{i} + \mathbf{j} + \frac{13}{5}\mathbf{k}$ or $(\frac{15}{13}, -\frac{5}{13}, 0), -\frac{2}{13}\mathbf{i} + \frac{5}{13}\mathbf{j} + \mathbf{k}$	M1 A1	
	M1: Uses their Cartesian equations correctly to obtain a point and direction A1: Correct point and direction – it may not be clear which is which – i.e. look for the correct numbers either as points or vectors		
	Equation of line in required form: e.g. $\mathbf{r} \times (-2\mathbf{i} + 5\mathbf{j} + 13\mathbf{k}) = -5\mathbf{i} - 15\mathbf{j} + 5\mathbf{k}$ Or Equivalent	M1 A1	
		(6)	
		Total 14	
(c) Way 3	$\begin{pmatrix} 2\lambda + \mu \\ \lambda - \mu \\ 5\lambda - 2\mu \end{pmatrix} \cdot \begin{pmatrix} 3 \\ -4 \\ 2 \end{pmatrix} = 5 \Rightarrow 12\lambda + 3\mu = 5$	M1: Substitutes parametric form of Π_2 into the vector equation of Π_1	M1A1
		A1: Correct equation	
	$\mu = \frac{5}{3}, \lambda = 0$ gives $(\frac{5}{3}, -\frac{5}{3}, \frac{10}{3})$ $\mu = 0, \lambda = \frac{5}{12}$ gives $(\frac{5}{6}, \frac{5}{12}, \frac{25}{12})$ Direction $\begin{pmatrix} -2 \\ 5 \\ 13 \end{pmatrix}$	M1: Finds 2 points and direction	M1A1
	A1: Correct coordinates and direction		
	Equation of line in required form: e.g. $\mathbf{r} \times (-2\mathbf{i} + 5\mathbf{j} + 13\mathbf{k}) = -5\mathbf{i} - 15\mathbf{j} + 5\mathbf{k}$ Or Equivalent	M1A1	
	Do not allow ‘mixed’ methods – mark the best single attempt		
	NB for checking, a general point on the line will be of the form: $(1 - 2\lambda, 5\lambda, 1 + 13\lambda)$		