


June 2011  
Further Pure Mathematics FP3 6669  
Mark Scheme

Question Number	Scheme	Marks
<b>1.</b>	$\frac{dy}{dx} = 6x^2 \text{ and so surface area} = 2\pi \int 2x^3 \sqrt{1+(6x^2)^2} dx$ $= 4\pi \left[ \frac{2}{3 \times 36 \times 4} (1+36x^4)^{\frac{3}{2}} \right]$ <p>Use limits 2 and 0 to give <math>\frac{4\pi}{216} [13860.016 - 1] = 806</math> (to 3 sf)</p>	B1 M1 A1 DM1 A1  <b>5</b>
	<b>Notes:</b>	
<b>B1</b>	Both bits CAO but condone lack of $2\pi$	
<b>1M1</b>	Integrating $\int \left( y \sqrt{1 + \left( \text{their } \frac{dy}{dx} \right)^2} \right) dx$ , getting $k(1+36x^4)^{\frac{3}{2}}$ , condone lack of $2\pi$	
<b>1A1</b>	CAO	
<b>2DM1</b>	Correct use of 2 and 0 as limits	
<b>2A1</b>	CAO	
<b>2.</b>		
<b>(a) (i)</b>	$\frac{dy}{dx} = \frac{x}{\sqrt{1-x^2}} + \arcsin x$	M1 A1
<b>(ii)</b>	At given value derivative $= \frac{1}{\sqrt{3}} + \frac{\pi}{6} = \frac{2\sqrt{3} + \pi}{6}$	B1 (2)
		(1)
<b>(b)</b>	$\frac{dy}{dx} = \frac{6e^{2x}}{1+9e^{4x}}$ $= \frac{6}{e^{-2x} + 9e^{2x}}$ $= \frac{3}{\frac{5}{2}(e^{2x} + e^{-2x}) + \frac{4}{2}(e^{2x} - e^{-2x})}$ $\therefore \frac{dy}{dx} = \frac{3}{5 \cosh 2x + 4 \sinh 2x} \quad *$	1M1 A1 2M1 3M1 A1 cso  (5) <b>8</b>
	<b>Notes:</b>	
<b>(a) M1</b>	Differentiating getting an arcsinx term and a $\frac{1}{\sqrt{1 \pm x^2}}$ term	
<b>A1</b>	CAO	
<b>B1</b>	CAO any correct form	

Question Number	Scheme	Marks
<b>(b) 1M1</b> <b>1A1</b> <b>2M1</b> <b>3M1</b> <b>2A1</b>	Of correct form $\frac{ae^{2x}}{1 \pm be^{4x}}$ CAO Getting from expression in $e^{4x}$ to $e^{2x}$ and $e^{-2x}$ only Using $\sinh 2x$ and $\cosh 2x$ in terms of $(e^{2x} + e^{-2x})$ and $(e^{2x} - e^{-2x})$ CSO – answer given	
<b>3.</b> <b>(a)</b>	$x^2 - 10x + 34 = (x-5)^2 + 9 \quad \text{so} \quad \frac{1}{x^2 - 10x + 34} = \frac{1}{(x-5)^2 + 9} = \frac{1}{u^2 + 9}$ (mark can be earned in either part (a) or (b)) $I = \int \frac{1}{u^2 + 9} du = \left[ \frac{1}{3} \arctan\left(\frac{u}{3}\right) \right] \quad \left  \quad I = \int \frac{1}{(x-5)^2 + 9} du = \left[ \frac{1}{3} \arctan\left(\frac{x-5}{3}\right) \right] \right.$ Uses limits 3 and 0 to give $\frac{\pi}{12}$ Uses limits 8 and 5 to give $\frac{\pi}{12}$	B1 M1 A1 DM1 A1 (5)
<b>(b) Alt 1</b>   <b>(b) Alt 2</b>   <b>(b) Alt 3</b>	$I = \ln\left(\left(\frac{x-5}{3}\right) + \sqrt{\left(\frac{x-5}{3}\right)^2 + 1}\right) \quad \text{or} \quad I = \ln\left(\frac{x-5 + \sqrt{(x-5)^2 + 9}}{3}\right)$ $\text{or} \quad I = \ln\left((x-5) + \sqrt{(x-5)^2 + 9}\right)$ Uses limits 5 and 8 to give $\ln(1 + \sqrt{2})$ . Uses $u = x-5$ to get $I = \int \frac{1}{\sqrt{u^2 + 9}} du = \left[ \operatorname{arsinh}\left(\frac{u}{3}\right) \right] = \ln\left\{u + \sqrt{u^2 + 9}\right\}$ Uses limits 3 and 0 and $\ln$ expression to give $\ln(1 + \sqrt{2})$ . Use substitution $x-5 = 3 \tan \theta$ , $\frac{dx}{d\theta} = 3 \sec^2 \theta$ and so $I = \int \sec \theta d\theta = \ln(\sec \theta + \tan \theta)$ Uses limits 0 and $\frac{\pi}{4}$ to get $\ln(1 + \sqrt{2})$ .	M1 A1 DM1 A1 (4) <b>9</b> M1 A1 DM1 A1 (4) M1 A1 DM1 A1 (4)
<b>(a) B1</b> <b>1M1</b> <b>1A1</b> <b>2DM1</b> <b>2A1</b>	<p style="text-align: center;"><b>Notes:</b></p> CAO allow recovery in (b) Integrating getting k arctan term CAO Correctly using limits. CAO	

Question Number	Scheme	Marks
<p><b>(b) 1M1</b> Integrating to get a ln or hyperbolic term  <b>1A1</b> CAO  <b>2DM1</b> Correctly using limits.  <b>2A1</b> CAO</p>		
<p><b>4.</b></p> <p><b>(a)</b></p> $I_n = \left[ \frac{x^3}{3} (\ln x)^n \right] - \int \frac{x^3}{3} \times \frac{n(\ln x)^{n-1}}{x} dx$ $= \left[ \frac{x^3}{3} (\ln x)^n \right]_1^e - \int_1^e \frac{nx^2 (\ln x)^{n-1}}{3} dx$ $\therefore I_n = \frac{e^3}{3} - \frac{n}{3} I_{n-1} \quad *$		<p>M1 A1</p> <p>DM1</p> <p>A1cso</p> <p>(4)</p>
<p><b>(b)</b></p> $I_0 = \int_1^e x^2 dx = \left[ \frac{x^3}{3} \right]_1^e = \frac{e^3}{3} - \frac{1}{3} \text{ or } I_1 = \frac{e^3}{3} - \frac{1}{3} \left( \frac{e^3}{3} - \frac{1}{3} \right) = \frac{2e^3}{9} + \frac{1}{9}$ $I_1 = \frac{e^3}{3} - \frac{1}{3} I_0, \quad I_2 = \frac{e^3}{3} - \frac{2}{3} I_1 \text{ and } I_3 = \frac{e^3}{3} - \frac{3}{3} I_2 \text{ so } I_3 = \frac{4e^3}{27} + \frac{2}{27}$ <p style="text-align: center;"><b>Notes:</b></p> <p><b>(a)1M1</b> Using integration by parts, integrating <math>x^2</math>, differentiating <math>(\ln x)^n</math>  <b>1A1</b> CAO  <b>2DM1</b> Correctly using limits 1 and e  <b>2A1</b> CSO answer given</p> <p><b>(b)1M1</b> Evaluating <math>I_0</math> or <math>I_1</math> by an attempt to integrate something  <b>1A1</b> CAO  <b>2M1</b> Finding <math>I_3</math> (also probably <math>I_1</math> and <math>I_2</math>) If 'n's left in M0  <b>2A1</b> <math>I_3</math> CAO</p>		<p>M1 A1</p> <p>M1 A1</p> <p>(4)</p> <p><b>8</b></p>

Question Number	Scheme	Marks
<p>5. (a)</p>	 <p>Graph of <math>y = 3\sinh 2x</math></p> <p>Shape of <math>-e^{2x}</math> graph</p> <p>Asymptote: <math>y = 13</math></p> <p>Value 10 on y axis and value 0.7 or <math>\frac{1}{2} \ln\left(\frac{13}{3}\right)</math> on x axis</p>	<p>B1</p> <p>B1</p> <p>B1</p> <p>B1</p> <p>(4)</p>
<p>(b)</p>	<p>Use definition <math>\frac{3}{2}(e^{2x} - e^{-2x}) = 13 - 3e^{2x} \rightarrow 9e^{4x} - 26e^{2x} - 3 = 0</math> to form quadratic</p> <p><math>\therefore e^{2x} = -\frac{1}{9}</math> or 3</p> <p><math>\therefore x = \frac{1}{2} \ln(3)</math></p>	<p>M1 A1</p> <p>DM1 A1</p> <p>B1</p> <p>(5)</p> <p><b>9</b></p>
<p style="text-align: center;"><b>Notes:</b></p> <p>(a) <b>1B1</b> <math>y = 3\sinh 2x</math> first and third quadrant.  <b>2B1</b> Shape of <math>y = -e^{2x}</math> correct intersects on positive axes.  <b>3B1</b> <b>Equation</b> of asymptote, <math>y = 13</math>, given. Penalise 'extra' asymptotes here  <b>4B1</b> Intercepts correct both</p> <p>(b) <b>1M1</b> Getting a three terms quadratic in <math>e^{2x}</math>  <b>1A1</b> Correct three term quadratic  <b>2DM1</b> Solving for <math>e^{2x}</math>  <b>2A1</b> CAO for <math>e^{2x}</math> condone omission of negative value.  <b>B1</b> CAO one answer only</p>		

Question Number	Scheme	Marks
<b>6.</b> <b>(a)</b>	$\mathbf{n} = (2\mathbf{j} - \mathbf{k}) \times (3\mathbf{i} + 2\mathbf{j} + 2\mathbf{k}) = 6\mathbf{i} - 3\mathbf{j} - 6\mathbf{k}$ o.a.e. (e.g. $2\mathbf{i} - \mathbf{j} - 2\mathbf{k}$ )	M1 A1 (2)
<b>(b)</b>	Line $l$ has direction $2\mathbf{i} - 2\mathbf{j} - \mathbf{k}$ Angle between line $l$ and normal is given by $(\cos \beta \text{ or } \sin \alpha) = \frac{4+2+2}{\sqrt{9}\sqrt{9}} = \frac{8}{9}$ $\alpha = 90 - \beta = 63$ degrees to nearest degree.	B1 M1 A1ft A1 awrt (4)
<b>(c) Alt 1</b>	Plane $P$ has equation $\mathbf{r} \cdot (2\mathbf{i} - \mathbf{j} - 2\mathbf{k}) = 1$ Perpendicular distance is $\frac{1 - (-7)}{\sqrt{9}} = \frac{8}{3}$	M1 A1 M1 A1 (4) <b>10</b>
<b>(c) Alt 2</b>	Parallel plane through A has equation $\mathbf{r} \cdot \frac{2\mathbf{i} - \mathbf{j} - 2\mathbf{k}}{3} = \frac{-7}{3}$ Plane P has equation $\mathbf{r} \cdot \frac{2\mathbf{i} - \mathbf{j} - 2\mathbf{k}}{3} = \frac{1}{3}$ So O lies between the two and perpendicular distance is $\frac{1}{3} + \frac{7}{3} = \frac{8}{3}$	M1 A1 M1 A1 (4)
<b>(c) Alt 3</b>	Distance A to $(3,1,2) = \sqrt{2^2 + 2^2 + 1^2} = 3$ Perpendicular distance is '3' $\sin \alpha = 3 \times \frac{8}{9} = \frac{8}{3}$	M1A1 M1A1 (4)
<b>(c) Alt 4</b>	Finding Cartesian equation of plane P: $2x - y - 2z - 1 = 0$ $d = \frac{ n_1\alpha + n_2\beta + n_3\gamma + d }{\sqrt{n_1^2 + n_2^2 + n_3^2}} = \frac{ 2(1) - 1(3) - 2(3) - 1 }{\sqrt{2^2 + 1^2 + 2^2}} = \frac{8}{3}$	M1 A1 M1A1 (4)
	<b>Notes:</b> <b>(a) M1</b> Cross product of the correct vectors <b>A1</b> CAO o.e. <b>(b) B1</b> CAO <b>M1</b> Angle between ' $2\mathbf{i} - \mathbf{j} - 2\mathbf{k}$ ' and $2\mathbf{i} - 2\mathbf{j} - \mathbf{k}$ , formula of correct form <b>1A1ft</b> 8/9ft <b>2A1</b> CAO awrt <b>(c) 1M1</b> Eqn of plane using $2\mathbf{i} - \mathbf{j} - 2\mathbf{k}$ or dist of A from O or finding length of AP <b>1A1</b> Correct equation (must have =) or A to $(3,1,2) = 3$ <b>2M1</b> Using correct method to find perpendicular distance <b>2A1</b> CAO	

Question Number	Scheme	Marks
7. (a)	$\text{Det } \mathbf{M} = k(0 - 2) + 1(1 + 3) + 1(-2 - 0) = -2k + 4 - 2 = 2 - 2k$	M1 A1 (2)
(b)	$\mathbf{M}^T = \begin{pmatrix} k & 1 & 3 \\ -1 & 0 & -2 \\ 1 & -1 & 1 \end{pmatrix} \text{ so cofactors} = \begin{pmatrix} -2 & -1 & 1 \\ -4 & k-3 & k+1 \\ -2 & 2k-3 & 1 \end{pmatrix}$ <p>(-1 A mark for each term wrong)</p> $\mathbf{M}^{-1} = \frac{1}{2-2k} \begin{pmatrix} -2 & -1 & 1 \\ -4 & k-3 & k+1 \\ -2 & 2k-3 & 1 \end{pmatrix}$	M1  M1 A3 (5)
(c)	<p>Let <math>(x, y, z)</math> be on <math>l_1</math>. Equation of <math>l_2</math> can be written as <math>\begin{pmatrix} x' \\ y' \\ z' \end{pmatrix} = \begin{pmatrix} 4 \\ 1 \\ 7 \end{pmatrix} + \lambda \begin{pmatrix} 4 \\ 1 \\ 3 \end{pmatrix}</math>.</p> <p>Use <math>\begin{pmatrix} x \\ y \\ z \end{pmatrix} = \mathbf{M}^{-1} \begin{pmatrix} x' \\ y' \\ z' \end{pmatrix}</math> with <math>k = 2</math>. i.e. <math>\begin{pmatrix} x \\ y \\ z \end{pmatrix} = \frac{1}{-2} \begin{pmatrix} -2 &amp; -1 &amp; 1 \\ -4 &amp; -1 &amp; 3 \\ -2 &amp; 1 &amp; 1 \end{pmatrix} \begin{pmatrix} 4+4\lambda \\ 1+\lambda \\ 7+3\lambda \end{pmatrix}</math></p> $\therefore \begin{pmatrix} x \\ y \\ z \end{pmatrix} = \begin{pmatrix} 3\lambda + 1 \\ 4\lambda - 2 \\ 2\lambda \end{pmatrix}$ <p>and so <math>(\mathbf{r} - \mathbf{a}) \times \mathbf{b} = \mathbf{0}</math> where <math>\mathbf{a} = \mathbf{i} - 2\mathbf{j}</math> and <math>\mathbf{b} = 3\mathbf{i} + 4\mathbf{j} + 2\mathbf{k}</math> or equivalent  <b>or</b> <math>\mathbf{r} = \mathbf{a} + \lambda \mathbf{b}</math> where <math>\mathbf{a} = \mathbf{i} - 2\mathbf{j}</math> and <math>\mathbf{b} = 3\mathbf{i} + 4\mathbf{j} + 2\mathbf{k}</math> or equivalent</p>	B1  M1  M1 A1  B1ft (5) <b>12</b>
(a) M1 A1	<p style="text-align: center;"><b>Notes:</b></p> <p>Finding determinant at least one component correct.                  CAO</p>	
(b) 1M1 2M1 1A1 2A1 3A1	<p>Finding matrix of cofactors or its transpose                  Finding inverse matrix, 1/(det) cofactors + transpose                  At least seven terms correct (so at most 2 incorrect) condone missing det                  At least eight terms correct (so at most 1 incorrect) condone missing det                  All nine terms correct, condone missing det</p>	
(c) 1B1 1M1 2M1 A1 2B1	<p>Equation of <math>l_2</math>                  Using inverse transformation matrix correctly                  Finding general point in terms of <math>\lambda</math>.                  CAO for general point in terms of one parameter                  ft for vector equation of their <math>l_1</math></p>	

Question Number	Scheme	Marks
<p><b>8.</b></p> <p><b>(a)</b></p>	<p>Uses <math>\frac{dy}{dx} = \frac{\frac{dy}{d\theta}}{\frac{dx}{d\theta}} = \frac{b \cosh \theta}{a \sinh \theta}</math>      <b>or</b>      <math>\frac{2x}{a^2} - \frac{2yy'}{b^2} = 0 \rightarrow y' = \frac{xb^2}{ya^2} = \frac{b \cosh \theta}{a \sinh \theta}</math></p> <p>So <math>y - b \sinh \theta = \frac{b \cosh \theta}{a \sinh \theta} (x - a \cosh \theta)</math></p> <p><math>\therefore ab(\cosh^2 \theta - \sinh^2 \theta) = xb \cosh \theta - ya \sinh \theta</math> and as <math>(\cosh^2 \theta - \sinh^2 \theta) = 1</math></p> <p><math>xb \cosh \theta - ya \sinh \theta = ab</math>      *</p>	<p>M1 A1</p> <p>M1</p> <p>A1cso</p> <p>(4)</p>
<p><b>(b)</b></p>	<p><math>P</math> is the point <math>(\frac{a}{\cosh \theta}, 0)</math></p>	<p>M1 A1</p> <p>(2)</p>
<p><b>(c)</b></p>	<p><math>l_2</math> has equation <math>x = a</math> and meets <math>l_1</math> at <math>Q(a, \frac{b(\cosh \theta - 1)}{\sinh \theta})</math></p>	<p>M1 A1</p> <p>(2)</p>
<p><b>(d) Alt 1</b></p> <p><b>(d) Alt 2</b></p>	<p>The mid point of <math>PQ</math> is given by <math>X = \frac{a(\cosh \theta + 1)}{2 \cosh \theta}</math>, <math>Y = \frac{b(\cosh \theta - 1)}{2 \sinh \theta}</math></p> <p><math>4Y^2 + b^2 = b^2 \left( \frac{\cosh^2 \theta + 1 - 2 \cosh \theta + \sinh^2 \theta}{\sinh^2 \theta} \right)</math></p> <p><math>= b^2 \left( \frac{2 \cosh^2 \theta - 2 \cosh \theta}{\sinh^2 \theta} \right)</math></p> <p><math>X(4Y^2 + b^2) = ab^2 \left( \frac{(\cosh \theta + 1)(\cosh \theta - 1)2 \cosh \theta}{2 \cosh \theta \sinh^2 \theta} \right)</math></p> <p>Simplify fraction by using <math>\cosh^2 \theta - \sinh^2 \theta = 1</math> to give <math>x(4y^2 + b^2) = ab^2</math>      *</p> <p>First line of solution as before</p> <p><math>4Y^2 + b^2 = b^2 (\coth^2 \theta + \operatorname{cosech}^2 \theta - 2 \coth \theta \operatorname{cosech} \theta + 1)</math></p> <p><math>= b^2 (2 \coth^2 \theta - 2 \coth \theta \operatorname{cosech} \theta)</math></p> <p><math>X(4Y^2 + b^2) = ab^2 (\coth \theta (\coth \theta - \operatorname{cosech} \theta) (1 + \operatorname{sech} \theta))</math></p> <p>Simplify expansion by using <math>\coth^2 \theta - \operatorname{cosech}^2 \theta = 1</math> to give <math>x(4y^2 + b^2) = ab^2</math>      *</p>	<p>1M1 A1ft</p> <p>2M1</p> <p>3M1</p> <p>4M1</p> <p>A1cso</p> <p>(6)</p> <p>1M1A1ft</p> <p>2M1</p> <p>3M1</p> <p>4M1</p> <p>A1cso</p> <p>(6)</p> <p><b>14</b></p>

Question Number	Scheme	Marks
<b>8.</b> <b>(a)1M1</b> Finding gradient in terms of $\theta$ <b>1A1</b> CAO <b>2M1</b> Finding equation of tangent <b>2A1</b> CSO (answer given) look for $\pm(\cosh^2\theta - \sinh^2\theta)$  <b>(b)M1</b> Putting $y = 0$ into their tangent <b>A1ft</b> P found, ft for their tangent o.e.  <b>(c) M1</b> Putting $x = a$ into their tangent. <b>A1</b> CAO Q found o.e.  <b>(d)</b> For Alt 1 and 2 <b>1M1</b> Finding expressions, in terms of $\sinh \theta$ and $\cosh \theta$ but must be adding <b>1A1</b> Ft on their P and Q, <b>2M1</b> Finding $4y^2 + b^2$ <b>3M1</b> Simplified, factorised, maximum of 2 terms per bracket <b>4M1</b> Finding $x(4y^2 + b^2)$ , completely factorised, maximum of 2 terms per bracket <b>2A1</b> CSO  <b>(d)</b> For Alts 3, 4 and 5 <b>1M1</b> Finding expressions, in terms of $\sinh \theta$ and $\cosh \theta$ but must be adding <b>1A1</b> Ft on their P and Q <b>2M1</b> Getting $\cosh \theta$ in terms of x <b>3M1</b> y or $y^2$ in terms of $\cosh \theta$ <b>or</b> $\sinh \theta$ in terms of x and y <b>4M1</b> Getting equation in terms of x and y only. No square roots. <b>2A1</b> CSO		



Question Number	Scheme	Marks
<p><b>8(d)</b> <b>Alt 3</b></p>	$X = \frac{a(\cosh \theta + 1)}{2 \cosh \theta}, \quad Y = \frac{b(\cosh \theta - 1)}{2 \sinh \theta}$ $\cosh \theta = \frac{a}{2x - a}$ $\sinh \theta = \frac{b(\cosh \theta - 1)}{2y} = \frac{b(a - x)}{(2x - a)y}$ $\left( \frac{a}{2x - a} \right)^2 - \left( \frac{b(a - x)}{(2x - a)y} \right)^2 = 1$ <p>Simplifies to give required equation  <math>[y^2 4x(a - x) = b^2(a - x)^2, \quad x(4y^2 + b^2) = ab^2]</math></p>	<p>As main scheme 1M1 A1ft</p> <p>cosh <math>\theta</math> in terms of x 2M1</p> <p>sinh <math>\theta</math> in terms of x and y 3M1</p> <p>Using <math>\cosh^2 \theta - \sinh^2 \theta = 1</math> 4M1</p> <p>A1cso (6)</p>
<p><b>Alt 4</b></p>	$X = \frac{a(\cosh \theta + 1)}{2 \cosh \theta}, \quad Y = \frac{b(\cosh \theta - 1)}{2 \sinh \theta}$ $\cosh \theta = \frac{a}{2x - a}$ $y^2 = \frac{b^2(\cosh \theta - 1)^2}{4(\cosh^2 \theta - 1)} = \frac{b^2(\cosh \theta - 1)}{4(\cosh \theta + 1)}$ $y^2 = \frac{b^2 \left( \frac{2a - 2x}{2x - a} \right)^2}{4 \left( \frac{2x}{2x - a} \right)} \text{ o.e.}$ <p>Simplifies to give required equation</p>	<p>As main scheme 1M1 A1ft</p> <p>cosh <math>\theta</math> in terms of x 2M1</p> <p><math>y^2</math> in terms of cosh <math>\theta</math> only 3M1</p> <p>Forms equation in x and y only 4M1</p> <p>A1 cso (6)</p>
<p><b>Alt 5</b></p>	$X = \frac{a(\cosh \theta + 1)}{2 \cosh \theta}, \quad Y = \frac{b(\cosh \theta - 1)}{2 \sinh \theta}$ $\cosh \theta = \frac{a}{2x - a}$ $y = \left( \frac{b(\cosh \theta - 1)}{2 \sinh \theta} \right) = \left( \frac{b(\cosh \theta - 1)}{2 \sqrt{\cosh^2 \theta - 1}} \right)$ <p>Eliminate <math>\sqrt{\quad}</math> and forms equation in x and y Simplifies to give required equation</p>	<p>As main scheme 1M1 A1ft</p> <p>cosh <math>\theta</math> in terms of x 2M1</p> <p>y in terms of cosh <math>\theta</math> only 3M1</p> <p>4M1 A1cso</p>